



wwPDB X-ray Structure Validation Summary Report ⓘ

Apr 25, 2026 – 02:25 AM EDT

PDB ID : 1TQY / pdb_00001tqy
Title : The Actinorhodin Ketosynthase/Chain Length Factor
Authors : Keatinge-Clay, A.T.; Maltby, D.A.; Medzihradsky, K.F.; Khosla, C.; Stroud, R.M.
Deposited on : 2004-06-18
Resolution : 2.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	10052 (2.00-2.00)
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
Sidechain outliers	187428	11029 (2.00-2.00)
RSRZ outliers	180081	10067 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	424	7% 72% 25% ..
1	C	424	11% 75% 22% ..
1	E	424	21% 73% 25% ..
1	G	424	7% 73% 22% ..
2	B	415	6% 71% 23% ..

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	D	415	4% 75% 18%
2	F	415	6% 72% 22%
2	H	415	5% 74% 21%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	ACE	E	633	-	-	X	-
4	ACE	G	633	-	-	X	-

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 25535 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Actinorhodin polyketide putative beta-ketoacyl synthase 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	421	Total	C	N	O	S	0	0	0
			3140	1936	570	618	16			
1	C	421	Total	C	N	O	S	0	0	0
			3140	1936	570	618	16			
1	E	421	Total	C	N	O	S	0	0	0
			3140	1936	570	618	16			
1	G	421	Total	C	N	O	S	0	0	0
			3140	1936	570	618	16			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MET	-	initiating methionine	UNP Q02059
C	1	MET	-	initiating methionine	UNP Q02059
E	1	MET	-	initiating methionine	UNP Q02059
G	1	MET	-	initiating methionine	UNP Q02059

- Molecule 2 is a protein called Actinorhodin polyketide putative beta-ketoacyl synthase 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	402	Total	C	N	O	S	0	0	0
			2970	1863	537	565	5			
2	D	402	Total	C	N	O	S	0	0	0
			2970	1863	537	565	5			
2	F	402	Total	C	N	O	S	0	0	0
			2970	1863	537	565	5			
2	H	402	Total	C	N	O	S	0	0	0
			2970	1863	537	565	5			

There are 36 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	-7	MET	-	initiating methionine	UNP Q02062
B	-6	ASP	-	expression tag	UNP Q02062
B	-5	TYR	-	expression tag	UNP Q02062
B	-4	LYS	-	expression tag	UNP Q02062
B	-3	ASP	-	expression tag	UNP Q02062
B	-2	ASP	-	expression tag	UNP Q02062
B	-1	ASP	-	expression tag	UNP Q02062
B	0	ASP	-	expression tag	UNP Q02062
B	1	LYS	-	expression tag	UNP Q02062
D	-7	MET	-	initiating methionine	UNP Q02062
D	-6	ASP	-	expression tag	UNP Q02062
D	-5	TYR	-	expression tag	UNP Q02062
D	-4	LYS	-	expression tag	UNP Q02062
D	-3	ASP	-	expression tag	UNP Q02062
D	-2	ASP	-	expression tag	UNP Q02062
D	-1	ASP	-	expression tag	UNP Q02062
D	0	ASP	-	expression tag	UNP Q02062
D	1	LYS	-	expression tag	UNP Q02062
F	-7	MET	-	initiating methionine	UNP Q02062
F	-6	ASP	-	expression tag	UNP Q02062
F	-5	TYR	-	expression tag	UNP Q02062
F	-4	LYS	-	expression tag	UNP Q02062
F	-3	ASP	-	expression tag	UNP Q02062
F	-2	ASP	-	expression tag	UNP Q02062
F	-1	ASP	-	expression tag	UNP Q02062
F	0	ASP	-	expression tag	UNP Q02062
F	1	LYS	-	expression tag	UNP Q02062
H	-7	MET	-	initiating methionine	UNP Q02062
H	-6	ASP	-	expression tag	UNP Q02062
H	-5	TYR	-	expression tag	UNP Q02062
H	-4	LYS	-	expression tag	UNP Q02062
H	-3	ASP	-	expression tag	UNP Q02062
H	-2	ASP	-	expression tag	UNP Q02062
H	-1	ASP	-	expression tag	UNP Q02062
H	0	ASP	-	expression tag	UNP Q02062
H	1	LYS	-	expression tag	UNP Q02062

- Molecule 3 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total Mg 1 1	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	C	1	Total	Mg	0	0
			1	1		
3	E	1	Total	Mg	0	0
			1	1		
3	G	1	Total	Mg	0	0
			1	1		

- Molecule 4 is ACETYL GROUP (CCD ID: ACE) (formula: C₂H₄O).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			3	2	1		
4	C	1	Total	C	O	0	0
			3	2	1		
4	E	1	Total	C	O	0	0
			3	2	1		
4	G	1	Total	C	O	0	0
			3	2	1		

- Molecule 5 is SODIUM ION (CCD ID: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	1	Total	Na	0	0
			1	1		
5	D	1	Total	Na	0	0
			1	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	F	1	Total 1	Na 1	0	0
5	H	1	Total 1	Na 1	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	99	Total 99	O 99	0	0
6	B	144	Total 144	O 144	0	0
6	C	105	Total 105	O 105	0	0
6	D	187	Total 187	O 187	0	0
6	E	62	Total 62	O 62	0	0
6	F	157	Total 157	O 157	0	0
6	G	110	Total 110	O 110	0	0
6	H	211	Total 211	O 211	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Actinorhodin polyketide putative beta-ketoacyl synthase 1




- Molecule 1: Actinorhodin polyketide putative beta-ketoacyl synthase 1




- Molecule 1: Actinorhodin polyketide putative beta-ketoacyl synthase 1




• Molecule 1: Actinorhodin polyketide putative beta-ketoacyl synthase 1





• Molecule 2: Actinorhodin polyketide putative beta-ketoacyl synthase 2







• Molecule 2: Actinorhodin polyketide putative beta-ketoacyl synthase 2




• Molecule 2: Actinorhodin polyketide putative beta-ketoacyl synthase 2



• Molecule 2: Actinorhodin polyketide putative beta-ketoacyl synthase 2





4 Data and refinement statistics

Property	Value	Source
Space group	H 3	Depositor
Cell constants a, b, c, α , β , γ	262.70Å 262.70Å 224.03Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	50.00 – 2.00 50.00 – 2.00	Depositor EDS
% Data completeness (in resolution range)	98.7 (50.00-2.00) 98.7 (50.00-2.00)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.70 (at 2.00Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.233 , 0.253 0.227 , 0.246	Depositor DCC
R_{free} test set	38543 reflections (10.00%)	wwPDB-VP
Wilson B-factor (Å ²)	25.2	Xtriage
Anisotropy	0.059	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 37.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.016 for h,-h-k,-l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	25535	wwPDB-VP
Average B, all atoms (Å ²)	30.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 11.06% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MG, NA, ACE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.47	0/3199	1.00	16/4346 (0.4%)
1	C	0.48	0/3199	0.96	12/4346 (0.3%)
1	E	0.43	0/3199	0.92	14/4346 (0.3%)
1	G	0.47	0/3199	0.98	16/4346 (0.4%)
2	B	0.47	0/3041	0.96	6/4151 (0.1%)
2	D	0.52	0/3041	1.01	13/4151 (0.3%)
2	F	0.47	0/3041	1.01	9/4151 (0.2%)
2	H	0.51	0/3041	1.03	16/4151 (0.4%)
All	All	0.48	0/24960	0.98	102/33988 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	B	0	1
2	F	0	2
All	All	0	3

There are no bond length outliers.

The worst 5 of 102 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	F	314	ARG	NE-CZ-NH2	-17.02	103.88	119.20
2	F	314	ARG	NE-CZ-NH1	14.05	135.55	121.50
2	H	314	ARG	NE-CZ-NH2	-14.01	106.59	119.20
2	H	314	ARG	NE-CZ-NH1	11.33	132.83	121.50
1	A	144	VAL	CA-C-N	8.98	128.62	119.82

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	B	137	TYR	Sidechain
2	F	137	TYR	Sidechain
2	F	314	ARG	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3140	0	3032	84	0
1	C	3140	0	3032	76	0
1	E	3140	0	3032	79	0
1	G	3140	0	3032	83	0
2	B	2970	0	2903	79	0
2	D	2970	0	2903	67	0
2	F	2970	0	2903	78	0
2	H	2970	0	2903	69	0
3	A	1	0	0	0	0
3	C	1	0	0	0	0
3	E	1	0	0	0	0
3	G	1	0	0	0	0
4	A	3	0	3	1	0
4	C	3	0	3	1	0
4	E	3	0	3	2	0
4	G	3	0	3	2	0
5	B	1	0	0	0	0
5	D	1	0	0	0	0
5	F	1	0	0	0	0
5	H	1	0	0	0	0
6	A	99	0	0	1	0
6	B	144	0	0	2	1
6	C	105	0	0	9	0
6	D	187	0	0	11	0
6	E	62	0	0	1	0
6	F	157	0	0	7	1
6	G	110	0	0	6	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	H	211	0	0	4	0
All	All	25535	0	23752	570	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

The worst 5 of 570 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:402:ARG:HG3	6:D:1237:HOH:O	1.39	1.21
1:G:376:THR:HG23	6:G:1143:HOH:O	1.47	1.13
1:C:392:GLU:HB2	6:C:1124:HOH:O	1.50	1.11
1:E:275:MET:HE3	2:F:146:ILE:HD12	1.36	1.07
1:C:146:SER:HB2	6:D:1189:HOH:O	1.54	1.05

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
6:B:1122:HOH:O	6:F:1120:HOH:O[9_554]	2.14	0.06

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	419/424 (99%)	397 (95%)	22 (5%)	0	100	100
1	C	419/424 (99%)	393 (94%)	25 (6%)	1 (0%)	43	42
1	E	419/424 (99%)	394 (94%)	24 (6%)	1 (0%)	43	42
1	G	419/424 (99%)	399 (95%)	17 (4%)	3 (1%)	18	14
2	B	400/415 (96%)	379 (95%)	21 (5%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	D	400/415 (96%)	383 (96%)	16 (4%)	1 (0%)	36	35
2	F	400/415 (96%)	385 (96%)	15 (4%)	0	100	100
2	H	400/415 (96%)	381 (95%)	18 (4%)	1 (0%)	36	35
All	All	3276/3356 (98%)	3111 (95%)	158 (5%)	7 (0%)	43	42

5 of 7 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	G	310	GLY
1	G	388	LEU
1	E	388	LEU
1	G	109	ALA
1	C	146	SER

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	319/321 (99%)	308 (97%)	11 (3%)	32	33
1	C	319/321 (99%)	309 (97%)	10 (3%)	35	37
1	E	319/321 (99%)	312 (98%)	7 (2%)	45	50
1	G	319/321 (99%)	304 (95%)	15 (5%)	23	22
2	B	297/309 (96%)	277 (93%)	20 (7%)	15	11
2	D	297/309 (96%)	272 (92%)	25 (8%)	10	7
2	F	297/309 (96%)	278 (94%)	19 (6%)	16	12
2	H	297/309 (96%)	279 (94%)	18 (6%)	17	14
All	All	2464/2520 (98%)	2339 (95%)	125 (5%)	21	19

5 of 125 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	D	269	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	H	136	PHE
1	E	389	GLU
2	H	18	LEU
2	H	269	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 22 such sidechains are listed below:

Mol	Chain	Res	Type
2	D	395	ASN
2	F	248	HIS
2	F	55	HIS
2	F	395	ASN
1	C	260	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 12 ligands modelled in this entry, 8 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	ACE	C	633	1	1,2,2	1.05	0	0,1,1	-	-
4	ACE	E	633	1	1,2,2	1.04	0	0,1,1	-	-
4	ACE	G	633	1	1,2,2	1.07	0	0,1,1	-	-
4	ACE	A	633	1	1,2,2	1.06	0	0,1,1	-	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	C	633	ACE	1	0
4	E	633	ACE	2	0
4	G	633	ACE	2	0
4	A	633	ACE	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	421/424 (99%)	0.54	28 (6%) 24 22	20, 30, 45, 63	0
1	C	421/424 (99%)	0.68	45 (10%) 11 10	18, 29, 49, 79	0
1	E	421/424 (99%)	1.16	88 (20%) 2 2	20, 37, 56, 66	0
1	G	421/424 (99%)	0.47	31 (7%) 20 19	18, 28, 46, 62	0
2	B	402/415 (96%)	0.35	25 (6%) 26 25	18, 27, 43, 56	0
2	D	402/415 (96%)	0.07	18 (4%) 38 37	16, 23, 39, 56	0
2	F	402/415 (96%)	0.51	26 (6%) 25 23	19, 28, 45, 62	0
2	H	402/415 (96%)	0.17	21 (5%) 33 31	17, 24, 41, 59	0
All	All	3292/3356 (98%)	0.50	282 (8%) 16 15	16, 28, 48, 79	0

The worst 5 of 282 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	F	403	PHE	7.4
1	C	423	ALA	5.3
1	C	421	GLY	5.2
2	H	403	PHE	5.1
1	E	54	ASP	5.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	ACE	G	633	3/3	0.70	0.28	46,46,47,47	0
3	MG	A	1095	1/1	0.71	0.23	51,51,51,51	0
4	ACE	E	633	3/3	0.73	0.25	47,47,48,48	0
4	ACE	A	633	3/3	0.76	0.26	52,52,52,52	0
4	ACE	C	633	3/3	0.79	0.22	52,52,53,53	0
5	NA	B	1091	1/1	0.83	0.12	39,39,39,39	0
3	MG	E	1097	1/1	0.87	0.20	47,47,47,47	0
3	MG	C	1096	1/1	0.87	0.14	39,39,39,39	0
5	NA	D	1092	1/1	0.92	0.08	38,38,38,38	0
3	MG	G	1098	1/1	0.94	0.12	35,35,35,35	0
5	NA	H	1094	1/1	0.95	0.09	36,36,36,36	0
5	NA	F	1093	1/1	0.96	0.07	45,45,45,45	0

6.5 Other polymers [i](#)

There are no such residues in this entry.