



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 12, 2026 – 02:35 PM UTC

PDB ID : 3TQO / pdb_00003tqo
Title : Structure of the cysteinyl-tRNA synthetase (cysS) from Coxiella burnetii.
Authors : Rudolph, M.; Cheung, J.; Franklin, M.C.; Cassidy, M.; Gary, E.; Burshteyn, F.; Love, J.
Deposited on : 2011-09-09
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6319 (2.30-2.30)
Clashscore	190562	6919 (2.30-2.30)
Ramachandran outliers	187476	6854 (2.30-2.30)
Sidechain outliers	187428	6854 (2.30-2.30)
RSRZ outliers	180081	6325 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	462	5% 71% 12% 16%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 3310 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CysteinyI-tRNA synthetase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	387	Total	C	N	O	S	Se	0	0	0
			3159	2027	545	574	3	10			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	ASN	-	expression tag	UNP Q83BL7
A	0	ALA	-	expression tag	UNP Q83BL7

- Molecule 2 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Zn	0	0
			1	1		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	150	Total	O	0	0
			150	150		

4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, α , β , γ	97.63Å 97.63Å 108.14Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	36.23 – 2.30 36.23 – 2.30	Depositor EDS
% Data completeness (in resolution range)	99.1 (36.23-2.30) 98.9 (36.23-2.30)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.58 (at 2.29Å)	Xtriage
Refinement program	PHENIX 1.6.4_486	Depositor
R, R_{free}	0.208 , 0.260 0.214 , 0.256	Depositor DCC
R_{free} test set	1222 reflections (5.13%)	wwPDB-VP
Wilson B-factor (Å ²)	31.7	Xtriage
Anisotropy	0.529	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 46.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	3310	wwPDB-VP
Average B, all atoms (Å ²)	37.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.92% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.42	6/3223 (0.2%)	0.66	0/4324

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	50	MSE	SE-CE	-5.54	1.78	1.95
1	A	37	MSE	SE-CE	-5.41	1.79	1.95
1	A	57	MSE	SE-CE	-5.38	1.79	1.95
1	A	31	MSE	SE-CE	-5.24	1.79	1.95
1	A	213	MSE	SE-CE	-5.16	1.79	1.95
1	A	256	MSE	SE-CE	-5.02	1.80	1.95

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3159	0	3137	45	0
2	A	1	0	0	0	0
3	A	150	0	0	3	0
All	All	3310	0	3137	45	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (45) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:194:LYS:HG3	1:A:201:GLU:HB3	1.56	0.88
1:A:236:HIS:HD2	1:A:257:HIS:NE2	1.89	0.71
1:A:57:MSE:SE	1:A:391:ILE:HD11	2.42	0.68
1:A:263:ILE:O	1:A:266:GLU:HG2	1.97	0.65
1:A:50:MSE:HE3	1:A:350:PHE:HB3	1.81	0.62
1:A:114:ARG:O	1:A:117:GLN:HG2	2.02	0.59
1:A:50:MSE:CE	1:A:350:PHE:HB3	2.33	0.57
1:A:311:GLU:HA	1:A:314:ARG:NH1	2.20	0.56
1:A:29:CYS:SG	1:A:236:HIS:HE1	2.29	0.56
1:A:42:GLY:HA3	1:A:278:ILE:HD11	1.88	0.56
1:A:114:ARG:H	1:A:117:GLN:NE2	2.05	0.54
1:A:336:SER:HB3	1:A:384:GLU:OE2	2.09	0.52
1:A:351:ASN:OD1	1:A:354:ILE:HD13	2.09	0.52
1:A:38:HIS:HD2	1:A:40:GLY:N	2.08	0.51
1:A:114:ARG:H	1:A:117:GLN:HE21	1.59	0.50
1:A:311:GLU:HA	1:A:314:ARG:HH11	1.78	0.49
1:A:38:HIS:HA	1:A:277:SER:HA	1.95	0.49
1:A:137:THR:HG22	1:A:138:GLY:O	2.15	0.47
1:A:42:GLY:O	1:A:46:ILE:HG12	2.14	0.47
1:A:70:ASP:CG	1:A:116:THR:HG23	2.40	0.46
1:A:38:HIS:HD2	1:A:40:GLY:H	1.63	0.46
1:A:54:TYR:HA	1:A:391:ILE:HD13	1.96	0.46
1:A:152:ASP:HB3	1:A:155:LYS:HE3	1.97	0.45
1:A:305:TYR:CZ	1:A:310:LEU:HD11	2.52	0.45
1:A:362:MSE:O	1:A:366:ILE:HG13	2.17	0.45
1:A:194:LYS:CG	1:A:201:GLU:HB3	2.39	0.44
1:A:194:LYS:HG2	1:A:195:TRP:N	2.32	0.44
1:A:352:THR:OG1	1:A:353:PRO:HD3	2.18	0.44
1:A:253:LYS:HD3	3:A:558:HOH:O	2.18	0.43
1:A:194:LYS:HE3	1:A:201:GLU:CD	2.43	0.43
1:A:324:LEU:O	1:A:325:ARG:C	2.61	0.43
1:A:313:GLY:HA3	3:A:533:HOH:O	2.17	0.43
1:A:325:ARG:HA	1:A:399:PRO:HB2	2.00	0.43
1:A:132:ASN:O	1:A:133:GLN:HB2	2.19	0.42
1:A:183:VAL:HG11	1:A:186:LYS:HB3	2.02	0.42
1:A:137:THR:HA	1:A:138:GLY:HA3	1.85	0.42
1:A:266:GLU:HG3	1:A:274:ASN:ND2	2.35	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:325:ARG:HA	1:A:399:PRO:CB	2.50	0.42
1:A:351:ASN:CG	1:A:354:ILE:HD13	2.46	0.41
1:A:182:PHE:HE1	1:A:242:GLN:HE21	1.67	0.41
1:A:233:PHE:CG	1:A:234:PRO:HA	2.56	0.41
1:A:400:GLU:HG3	3:A:528:HOH:O	2.20	0.41
1:A:57:MSE:SE	1:A:391:ILE:CD1	3.16	0.41
1:A:11:LYS:HD3	1:A:158:HIS:CE1	2.56	0.40
1:A:124:LYS:HB3	1:A:124:LYS:HE2	1.80	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	381/462 (82%)	374 (98%)	6 (2%)	1 (0%)	36	46

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	325	ARG

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	339/392 (86%)	336 (99%)	3 (1%)	70 84

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	194	LYS
1	A	336	SER
1	A	391	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	38	HIS
1	A	117	GLN
1	A	236	HIS
1	A	298	HIS
1	A	367	ASN
1	A	390	ASN
1	A	396	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	377/462 (81%)	0.37	24 (6%)	25 27	18, 34, 59, 83	0

All (24) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	161	LEU	5.5
1	A	139	GLN	4.9
1	A	138	GLY	4.5
1	A	-1	ASN	4.1
1	A	400	GLU	3.4
1	A	162	ASP	3.2
1	A	325	ARG	3.0
1	A	160	HIS	2.9
1	A	290	LEU	2.9
1	A	175	SER	2.7
1	A	329	VAL	2.7
1	A	375	ILE	2.6
1	A	368	ARG	2.5
1	A	326	GLY	2.5
1	A	311	GLU	2.5
1	A	314	ARG	2.5
1	A	403	LEU	2.5
1	A	266	GLU	2.4
1	A	391	ILE	2.4
1	A	376	GLU	2.3
1	A	83	LYS	2.3
1	A	137	THR	2.2
1	A	147	VAL	2.1
1	A	312	ASN	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	ZN	A	461	1/1	0.99	0.05	25,25,25,25	0

6.5 Other polymers [i](#)

There are no such residues in this entry.