



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 03:17 PM UTC

PDB ID : 5TQ6 / pdb_00005tq6
Title : Design and Synthesis of a pan-JAK Kinase Inhibitor Clinical Candidate (PF-06263276) Suitable for Inhaled and Topical Delivery for the Treatment of Inflammatory Diseases of the Lungs and Skin
Authors : Chrencik, J.; Jones, P.
Deposited on : 2016-10-23
Resolution : 2.06 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.06 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3774 (2.08-2.04)
Clashscore	190562	3883 (2.08-2.04)
Ramachandran outliers	187476	3860 (2.08-2.04)
Sidechain outliers	187428	3860 (2.08-2.04)
RSRZ outliers	180081	3775 (2.08-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	304	
1	B	304	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 4846 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Tyrosine-protein kinase JAK2.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	276	Total	C	N	O	P	S	0	0	0
			2296	1459	394	428	2	13			
1	B	273	Total	C	N	O	P	S	0	0	0
			2277	1449	391	422	2	13			

There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	829	MET	-	initiating methionine	UNP O60674
A	830	GLY	-	expression tag	UNP O60674
A	831	HIS	-	expression tag	UNP O60674
A	832	HIS	-	expression tag	UNP O60674
A	833	HIS	-	expression tag	UNP O60674
A	834	HIS	-	expression tag	UNP O60674
A	835	HIS	-	expression tag	UNP O60674
A	836	HIS	-	expression tag	UNP O60674
A	1073	SER	MET	engineered mutation	UNP O60674
A	1076	THR	PHE	engineered mutation	UNP O60674
B	829	MET	-	initiating methionine	UNP O60674
B	830	GLY	-	expression tag	UNP O60674
B	831	HIS	-	expression tag	UNP O60674
B	832	HIS	-	expression tag	UNP O60674
B	833	HIS	-	expression tag	UNP O60674
B	834	HIS	-	expression tag	UNP O60674
B	835	HIS	-	expression tag	UNP O60674
B	836	HIS	-	expression tag	UNP O60674
B	1073	SER	MET	engineered mutation	UNP O60674
B	1076	THR	PHE	engineered mutation	UNP O60674

- Molecule 2 is {(3R,4R)-4-methyl-3-[methyl(7H-pyrrolo[2,3-d]pyrimidin-4-yl)amino]piperidin-1-yl}(pyrrolidin-1-yl)methanone (CCD ID: 7GV) (formula: C₁₈H₂₆N₆O).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			25	18	6	1		
2	B	1	Total	C	N	O	0	0
			25	18	6	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	109	Total	O	0	0
			109	109		
3	B	114	Total	O	0	0
			114	114		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Tyrosine-protein kinase JAK2



• Molecule 1: Tyrosine-protein kinase JAK2



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	36.98Å 183.40Å 52.58Å 90.00° 110.59° 90.00°	Depositor
Resolution (Å)	47.54 – 2.06 47.54 – 2.06	Depositor EDS
% Data completeness (in resolution range)	92.6 (47.54-2.06) 92.9 (47.54-2.06)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.54 (at 2.05Å)	Xtriage
Refinement program	REFMAC 5.8.0073	Depositor
R, R_{free}	0.257 , 0.289 0.259 , 0.291	Depositor DCC
R_{free} test set	1906 reflections (4.70%)	wwPDB-VP
Wilson B-factor (Å ²)	18.6	Xtriage
Anisotropy	0.103	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 27.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.42$, $\langle L^2 \rangle = 0.24$	Xtriage
Estimated twinning fraction	0.486 for h,-k,-h-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	4846	wwPDB-VP
Average B, all atoms (Å ²)	20.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 14.64% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PTR, 7GV

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.80	0/2308	0.95	0/3101
1	B	0.79	0/2289	0.95	4/3077 (0.1%)
All	All	0.80	0/4597	0.95	4/6178 (0.1%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	860	PHE	N-CA-C	6.46	122.28	113.37
1	B	1018	ILE	N-CA-C	5.64	117.05	110.62
1	B	1022	ALA	CA-C-N	5.08	124.69	119.05
1	B	1022	ALA	C-N-CA	5.08	124.69	119.05

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2296	0	2275	56	0
1	B	2277	0	2261	42	0
2	A	25	0	0	1	0
2	B	25	0	0	0	0
3	A	109	0	0	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	114	0	0	5	0
All	All	4846	0	4536	98	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

All (98) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:985:GLU:HG3	1:B:991:LYS:HE3	1.65	0.78
1:B:941:LEU:O	1:B:945:LYS:HB3	1.85	0.75
1:B:894:ASP:OD1	1:B:897:ARG:NH2	2.21	0.73
1:B:1096:ASP:O	1:B:1100:MET:HG3	1.88	0.73
1:A:1052:GLU:HG3	3:B:4104:HOH:O	1.89	0.73
1:B:854:GLN:NE2	1:B:856:GLY:O	2.22	0.72
1:A:904:SER:O	1:A:970:LYS:NZ	2.24	0.70
1:A:883:LYS:HD3	1:A:926:LYS:HD3	1.73	0.68
1:B:846:GLU:HB2	1:B:917:CYS:SG	2.34	0.68
1:A:1054:SER:O	1:A:1063:ARG:NH2	2.27	0.67
1:A:1051:ILE:HG22	1:A:1051:ILE:O	1.95	0.66
1:B:905:LEU:HD13	1:B:910:ILE:HD13	1.78	0.66
1:B:938:ARG:HA	1:B:1051:ILE:HD13	1.78	0.66
1:A:947:ARG:HH11	1:A:987:GLU:HG3	1.61	0.65
1:B:985:GLU:HG3	1:B:991:LYS:CE	2.26	0.65
1:A:850:LYS:HE3	1:A:869:ASP:HB3	1.80	0.64
1:A:1097:GLU:HG3	1:A:1126:ILE:HG12	1.81	0.62
1:B:914:LYS:HE3	1:B:930:GLU:HA	1.79	0.62
1:A:865:MET:HE3	1:A:931:TYR:CE2	2.34	0.62
1:A:1009:LYS:HE3	1:A:1028:GLU:O	2.04	0.58
1:B:1004:ASP:HB3	1:B:1005:LYS:HE3	1.86	0.58
1:B:1080:GLU:HG2	3:B:4109:HOH:O	2.03	0.58
1:A:1073:SER:O	1:A:1077:HIS:HD2	1.87	0.58
1:A:907:HIS:HE1	1:A:909:ASN:HD22	1.53	0.56
1:A:1011:LYS:HD2	1:A:1029:SER:OG	2.05	0.56
1:A:941:LEU:HD23	1:A:1051:ILE:CD1	2.36	0.56
1:B:893:ARG:HD3	3:B:4129:HOH:O	2.07	0.55
1:B:1010:VAL:O	1:B:1029:SER:HB3	2.06	0.55
1:A:940:TYR:OH	1:A:947:ARG:NH1	2.39	0.55
1:A:985:GLU:HB3	1:A:989:ARG:O	2.07	0.55
1:A:1028:GLU:HG2	3:A:4145:HOH:O	2.07	0.55
1:B:1034:ALA:HB3	1:B:1110:VAL:HG22	1.87	0.54

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:846:GLU:O	1:B:846:GLU:HG3	2.07	0.54
1:A:850:LYS:NZ	1:A:872:GLN:OE1	2.35	0.53
1:A:941:LEU:HD23	1:A:1051:ILE:HD13	1.91	0.53
1:B:848:HIS:HB3	1:B:869:ASP:C	2.34	0.52
1:B:854:GLN:OE1	1:B:864:GLU:HG2	2.09	0.52
1:B:1048:PHE:CZ	1:B:1098:ILE:HD13	2.44	0.52
1:B:850:LYS:HD3	1:B:869:ASP:HB2	1.92	0.52
1:A:1057:PRO:HB2	1:A:1058:PRO:HD3	1.91	0.51
1:A:1002:PRO:HD3	1:A:1008:PTR:HD2	1.92	0.51
1:A:1005:LYS:HE2	1:A:1007:PTR:O3P	2.11	0.51
1:A:932:LEU:HD22	1:A:985:GLU:HA	1.92	0.50
1:B:890:GLU:HB3	3:B:4190:HOH:O	2.11	0.50
1:A:1005:LYS:NZ	1:A:1007:PTR:O3P	2.45	0.49
1:A:855:LEU:HD11	1:A:931:TYR:CE1	2.47	0.49
1:B:914:LYS:HE3	1:B:930:GLU:CA	2.43	0.48
1:B:938:ARG:HA	1:B:1051:ILE:CD1	2.43	0.48
1:A:854:GLN:NE2	1:A:856:GLY:O	2.46	0.47
1:A:974:HIS:O	1:A:975:ARG:HB2	2.14	0.47
1:A:1090:ARG:NH2	1:A:1096:ASP:CG	2.72	0.47
1:B:1061:PHE:HA	1:B:1064:MET:HE3	1.96	0.47
1:B:1034:ALA:CB	1:B:1110:VAL:HG22	2.45	0.47
1:B:1073:SER:O	1:B:1077:HIS:ND1	2.37	0.47
1:A:958:SER:O	1:A:962:LYS:HG3	2.15	0.47
1:A:1099:TYR:O	1:A:1103:THR:HG23	2.15	0.47
1:B:932:LEU:HD13	1:B:985:GLU:HG2	1.96	0.47
1:A:1004:ASP:OD1	1:A:1005:LYS:HG3	2.16	0.46
1:A:1027:THR:HG22	1:A:1079:ILE:HD13	1.96	0.46
1:A:901:ILE:HD12	1:A:972:TYR:CZ	2.50	0.46
1:A:966:TYR:O	1:A:969:THR:HG22	2.16	0.45
1:A:912:LYS:HD3	1:A:930:GLU:CD	2.41	0.45
1:B:944:HIS:CD2	1:B:947:ARG:NH1	2.84	0.45
1:A:846:GLU:OE2	1:A:919:SER:HB3	2.16	0.45
1:B:850:LYS:HD3	1:B:869:ASP:CB	2.46	0.45
1:A:901:ILE:HD12	1:A:972:TYR:CE1	2.52	0.45
1:A:855:LEU:HD11	1:A:931:TYR:HE1	1.82	0.44
1:B:1104:GLU:CD	1:B:1122:ARG:HH22	2.25	0.44
1:A:1038:TRP:CE3	1:A:1106:TRP:HA	2.51	0.44
1:A:952:LYS:NZ	1:A:987:GLU:OE2	2.51	0.44
1:A:867:ARG:HD3	1:A:874:ASN:HA	2.00	0.44
1:A:913:TYR:CZ	1:A:915:GLY:HA2	2.53	0.44
1:A:858:GLY:HA3	2:A:4000:7GV:C23	2.48	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:954:LEU:HB2	1:B:1127:ARG:HD2	1.98	0.44
1:B:1100:MET:O	1:B:1104:GLU:HG3	2.18	0.43
1:B:852:LEU:HD23	1:B:865:MET:HE2	1.99	0.43
1:B:975:ARG:HD3	1:B:997:LEU:O	2.17	0.43
1:B:882:LYS:HB3	1:B:927:LEU:HB3	2.00	0.43
1:B:1008:PTR:HD1	1:B:1008:PTR:C	2.47	0.43
1:B:1038:TRP:CE3	1:B:1106:TRP:HA	2.53	0.43
1:A:976:ASP:OD1	1:A:980:ARG:NH2	2.51	0.43
1:A:1005:LYS:CE	1:A:1007:PTR:O3P	2.66	0.43
1:A:899:ILE:HG12	1:A:927:LEU:HD13	2.01	0.43
1:A:1039:SER:O	1:A:1043:VAL:HG23	2.19	0.42
1:A:903:LYS:HD2	3:A:4115:HOH:O	2.19	0.42
1:A:945:LYS:HA	1:A:948:ILE:HD12	2.01	0.42
1:A:1051:ILE:O	1:A:1051:ILE:CG2	2.64	0.42
1:B:848:HIS:O	1:B:869:ASP:N	2.39	0.42
1:A:975:ARG:HD3	1:A:997:LEU:O	2.19	0.42
1:A:893:ARG:HG2	1:A:897:ARG:CZ	2.49	0.42
1:A:1010:VAL:O	1:A:1029:SER:HB3	2.20	0.42
1:B:1000:VAL:HG22	3:B:4125:HOH:O	2.19	0.41
1:B:1057:PRO:HB2	1:B:1058:PRO:HD3	2.02	0.41
1:A:848:HIS:CG	1:A:870:PRO:HA	2.55	0.41
1:B:1051:ILE:HG22	1:B:1051:ILE:O	2.20	0.41
1:B:888:THR:HG22	1:B:889:GLU:N	2.35	0.40
1:A:971:ARG:NH2	1:A:1005:LYS:O	2.54	0.40
1:A:1098:ILE:HG22	1:A:1102:MET:HE3	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	266/304 (88%)	257 (97%)	9 (3%)	0	100 100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	263/304 (86%)	250 (95%)	12 (5%)	1 (0%)	30	23
All	All	529/608 (87%)	507 (96%)	21 (4%)	1 (0%)	43	38

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	1002	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	253/275 (92%)	232 (92%)	21 (8%)	10	4
1	B	252/275 (92%)	237 (94%)	15 (6%)	17	10
All	All	505/550 (92%)	469 (93%)	36 (7%)	13	7

All (36) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	857	LYS
1	A	871	LEU
1	A	889	GLU
1	A	896	GLU
1	A	918	TYR
1	A	925	LEU
1	A	937	LEU
1	A	946	GLU
1	A	962	LYS
1	A	967	LEU
1	A	969	THR
1	A	985	GLU
1	A	1003	GLN
1	A	1006	GLU
1	A	1010	VAL

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	1052	GLU
1	A	1073	SER
1	A	1080	GLU
1	A	1090	ARG
1	A	1110	VAL
1	A	1121	LEU
1	B	843	GLN
1	B	846	GLU
1	B	852	LEU
1	B	918	TYR
1	B	937	LEU
1	B	943	LYS
1	B	945	LYS
1	B	967	LEU
1	B	985	GLU
1	B	1010	VAL
1	B	1016	SER
1	B	1083	LYS
1	B	1087	ARG
1	B	1097	GLU
1	B	1110	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (16) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	853	GLN
1	A	859	ASN
1	A	885	GLN
1	A	906	GLN
1	A	909	ASN
1	A	950	HIS
1	A	955	GLN
1	A	988	ASN
1	A	1003	GLN
1	A	1077	HIS
1	A	1125	GLN
1	B	885	GLN
1	B	909	ASN
1	B	942	GLN
1	B	944	HIS
1	B	955	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	PTR	A	1008	1	15,16,17	0.69	0	17,22,24	0.80	0
1	PTR	B	1008	1	15,16,17	0.70	1 (6%)	17,22,24	0.86	1 (5%)
1	PTR	B	1007	1	15,16,17	0.84	0	17,22,24	0.91	0
1	PTR	A	1007	1	15,16,17	0.64	0	17,22,24	0.90	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	PTR	A	1008	1	-	1/10/11/13	0/1/1/1
1	PTR	B	1008	1	-	1/10/11/13	0/1/1/1
1	PTR	B	1007	1	-	1/10/11/13	0/1/1/1
1	PTR	A	1007	1	-	2/10/11/13	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	1008	PTR	OH-CZ	-2.04	1.36	1.40

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	1007	PTR	OH-P-O1P	-2.35	101.64	109.48

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	1008	PTR	OH-P-O1P	-2.26	101.94	109.48

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	1007	PTR	CZ-OH-P-O1P
1	B	1008	PTR	CZ-OH-P-O1P
1	A	1008	PTR	CZ-OH-P-O1P
1	B	1007	PTR	C-CA-CB-CG
1	A	1007	PTR	CZ-OH-P-O2P

There are no ring outliers.

3 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	1008	PTR	1	0
1	B	1008	PTR	1	0
1	A	1007	PTR	3	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	7GV	A	4000	-	27,28,28	0.66	1 (3%)	30,40,40	1.23	3 (10%)
2	7GV	B	4001	-	27,28,28	0.88	1 (3%)	30,40,40	1.43	5 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	7GV	A	4000	-	-	1/16/36/36	0/4/4/4
2	7GV	B	4001	-	-	2/16/36/36	0/4/4/4

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	4001	7GV	C11-C15	3.51	1.45	1.41
2	A	4000	7GV	C11-C15	2.66	1.44	1.41

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	4001	7GV	C11-C15-N14	-3.74	106.20	108.33
2	A	4000	7GV	C11-C15-N14	-3.51	106.33	108.33
2	A	4000	7GV	C15-N14-C13	3.03	109.86	108.11
2	B	4001	7GV	N18-C10-N8	-2.82	113.55	116.82
2	B	4001	7GV	C15-N14-C13	2.67	109.65	108.11
2	B	4001	7GV	C17-N18-C10	2.51	117.96	111.83
2	A	4000	7GV	C17-N18-C10	2.34	117.55	111.83
2	B	4001	7GV	C3-C2-C7	2.09	114.58	110.26

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	4001	7GV	N5-C19-N21-C25
2	B	4001	7GV	O20-C19-N21-C25
2	A	4000	7GV	O20-C19-N5-C4


There are no ring outliers.


1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	4000	7GV	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In

addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	274/304 (90%)	-1.59	0 100 100	6, 19, 34, 41	0
1	B	271/304 (89%)	-1.58	0 100 100	8, 18, 37, 42	0
All	All	545/608 (89%)	-1.59	0 100 100	6, 18, 36, 42	0

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	PTR	A	1007	16/17	0.99	0.03	26,29,35,35	4
1	PTR	A	1008	16/17	0.99	0.04	26,32,36,36	5
1	PTR	B	1007	16/17	0.99	0.04	25,29,30,31	4
1	PTR	B	1008	16/17	0.99	0.03	25,29,36,37	4

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.


6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	7GV	A	4000	25/25	0.99	0.03	18,20,22,23	0
2	7GV	B	4001	25/25	0.99	0.02	16,17,18,18	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers [i](#)

There are no such residues in this entry.