



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 09:28 PM UTC

PDB ID : 6TQ0 / pdb_00006tq0
Title : Crystal structure of the human Arc N-lobe bound to repeat peptide 5 from GKAP
Authors : Hallin, E.I.; Bramham, C.R.; Kursula, P.
Deposited on : 2019-12-15
Resolution : 1.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3494 (1.96-1.96)
Clashscore	190562	3612 (1.96-1.96)
Ramachandran outliers	187476	3587 (1.96-1.96)
Sidechain outliers	187428	3587 (1.96-1.96)
RSRZ outliers	180081	3495 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	75	3% 85% 11%
1	C	75	4% 80% 8% 11%
1	E	75	7% 73% 16% 9%
1	G	75	3% 77% 9% 12%
1	I	75	3% 81% 8% 11%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	K	75	4% 76% 12% 12%
1	M	75	% 81% 7% 11%
1	O	75	4% 76% 11% 13%
2	B	8	25% 62% 38%
2	D	8	38% 75% 25%
2	F	8	25% 50% 25% 12% 12%
2	H	8	38% 75% 12% 12%
2	J	8	12% 75% 25%
2	L	8	50% 75% 12% 12%
2	N	8	25% 62% 12% 25%
2	P	8	25% 88% 12%

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 9873 atoms, of which 4741 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Activity-regulated cytoskeleton-associated protein.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	67	Total	C	H	N	O	S	0	0	0
			1119	376	538	96	108	1			
1	C	67	Total	C	H	N	O	S	0	0	0
			1119	376	538	96	108	1			
1	E	68	Total	C	H	N	O	S	0	0	0
			1131	380	542	97	111	1			
1	G	66	Total	C	H	N	O	S	0	0	0
			1105	372	531	95	106	1			
1	I	67	Total	C	H	N	O	S	0	0	0
			1119	376	538	96	108	1			
1	K	66	Total	C	H	N	O	S	0	0	0
			1105	372	531	95	106	1			
1	M	67	Total	C	H	N	O	S	0	0	0
			1119	376	538	96	108	1			
1	O	65	Total	C	H	N	O	S	0	0	0
			1097	370	528	94	104	1			

There are 32 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	203	GLY	-	expression tag	UNP Q7LC44
A	204	ALA	-	expression tag	UNP Q7LC44
A	205	MET	-	expression tag	UNP Q7LC44
A	206	GLY	-	expression tag	UNP Q7LC44
C	203	GLY	-	expression tag	UNP Q7LC44
C	204	ALA	-	expression tag	UNP Q7LC44
C	205	MET	-	expression tag	UNP Q7LC44
C	206	GLY	-	expression tag	UNP Q7LC44
E	203	GLY	-	expression tag	UNP Q7LC44
E	204	ALA	-	expression tag	UNP Q7LC44
E	205	MET	-	expression tag	UNP Q7LC44
E	206	GLY	-	expression tag	UNP Q7LC44
G	203	GLY	-	expression tag	UNP Q7LC44

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
G	204	ALA	-	expression tag	UNP Q7LC44
G	205	MET	-	expression tag	UNP Q7LC44
G	206	GLY	-	expression tag	UNP Q7LC44
I	203	GLY	-	expression tag	UNP Q7LC44
I	204	ALA	-	expression tag	UNP Q7LC44
I	205	MET	-	expression tag	UNP Q7LC44
I	206	GLY	-	expression tag	UNP Q7LC44
K	203	GLY	-	expression tag	UNP Q7LC44
K	204	ALA	-	expression tag	UNP Q7LC44
K	205	MET	-	expression tag	UNP Q7LC44
K	206	GLY	-	expression tag	UNP Q7LC44
M	203	GLY	-	expression tag	UNP Q7LC44
M	204	ALA	-	expression tag	UNP Q7LC44
M	205	MET	-	expression tag	UNP Q7LC44
M	206	GLY	-	expression tag	UNP Q7LC44
O	203	GLY	-	expression tag	UNP Q7LC44
O	204	ALA	-	expression tag	UNP Q7LC44
O	205	MET	-	expression tag	UNP Q7LC44
O	206	GLY	-	expression tag	UNP Q7LC44

- Molecule 2 is a protein called repeat peptide 5 from GKAP.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	B	8	Total	C	H	N	O	S	0	0	0
			134	41	67	14	9	3			
2	D	6	Total	C	H	N	O	S	0	0	0
			93	30	46	9	6	2			
2	F	7	Total	C	H	N	O	S	0	0	0
			118	36	59	13	8	2			
2	H	7	Total	C	H	N	O	S	0	0	0
			118	36	59	13	8	2			
2	J	8	Total	C	H	N	O	S	0	0	0
			134	41	67	14	9	3			
2	L	7	Total	C	H	N	O	S	0	0	0
			118	36	59	13	8	2			
2	N	6	Total	C	H	N	O	S	0	0	0
			93	30	46	9	6	2			
2	P	7	Total	C	H	N	O	S	0	0	0
			109	35	54	10	7	3			



- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	12	Total 12	O 12	0	0
3	C	7	Total 7	O 7	0	0
3	D	1	Total 1	O 1	0	0
3	E	2	Total 2	O 2	0	0
3	G	5	Total 5	O 5	0	0
3	I	6	Total 6	O 6	0	0
3	K	5	Total 5	O 5	0	0
3	M	2	Total 2	O 2	0	0
3	O	2	Total 2	O 2	0	0



3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Activity-regulated cytoskeleton-associated protein




- Molecule 1: Activity-regulated cytoskeleton-associated protein




- Molecule 1: Activity-regulated cytoskeleton-associated protein




- Molecule 1: Activity-regulated cytoskeleton-associated protein




- Molecule 1: Activity-regulated cytoskeleton-associated protein




- Molecule 1: Activity-regulated cytoskeleton-associated protein




- Molecule 1: Activity-regulated cytoskeleton-associated protein





- Molecule 1: Activity-regulated cytoskeleton-associated protein




- Molecule 2: repeat peptide 5 from GKAP





- Molecule 2: repeat peptide 5 from GKAP




- Molecule 2: repeat peptide 5 from GKAP





- Molecule 2: repeat peptide 5 from GKAP



- Molecule 2: repeat peptide 5 from GKAP





- Molecule 2: repeat peptide 5 from GKAP



- Molecule 2: repeat peptide 5 from GKAP



- Molecule 2: repeat peptide 5 from GKAP



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	39.31Å 192.30Å 39.56Å 90.00° 94.80° 90.00°	Depositor
Resolution (Å)	100.00 – 1.95 96.15 – 1.95	Depositor EDS
% Data completeness (in resolution range)	98.2 (100.00-1.95) 98.3 (96.15-1.95)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.04	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.24 (at 1.95Å)	Xtriage
Refinement program	PHENIX 1.16-3549	Depositor
R, R_{free}	0.221 , 0.263 0.224 , 0.268	Depositor DCC
R_{free} test set	1998 reflections (4.79%)	wwPDB-VP
Wilson B-factor (Å ²)	45.0	Xtriage
Anisotropy	0.291	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.41 , 48.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.088 for l,-k,h	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	9873	wwPDB-VP
Average B, all atoms (Å ²)	72.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.82% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.69	0/600	0.55	0/806
1	C	0.61	0/600	0.52	0/806
1	E	0.56	0/608	0.45	0/817
1	G	0.56	0/593	0.45	0/796
1	I	0.64	0/600	0.53	0/806
1	K	0.53	0/593	0.44	0/796
1	M	0.53	0/600	0.46	0/806
1	O	0.51	0/588	0.42	0/791
2	B	0.45	0/68	0.49	0/86
2	D	0.45	0/48	0.45	0/61
2	F	0.67	0/60	0.59	0/75
2	H	0.41	0/60	0.53	0/75
2	J	0.52	0/68	0.52	0/86
2	L	0.27	0/60	0.32	0/75
2	N	0.46	0/48	0.42	0/61
2	P	0.31	0/56	0.36	0/72
All	All	0.57	0/5250	0.48	0/7015

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	581	538	538	1	0
1	C	581	538	538	5	0
1	E	589	542	542	7	0
1	G	574	531	531	3	0
1	I	581	538	538	4	0
1	K	574	531	531	2	1
1	M	581	538	538	2	0
1	O	569	528	528	2	1
2	B	67	67	67	1	0
2	D	47	46	46	0	0
2	F	59	59	59	3	0
2	H	59	59	59	0	0
2	J	67	67	67	1	0
2	L	59	59	59	0	0
2	N	47	46	46	1	0
2	P	55	54	54	0	0
3	A	12	0	0	0	0
3	C	7	0	0	1	0
3	D	1	0	0	0	0
3	E	2	0	0	0	0
3	G	5	0	0	0	0
3	I	6	0	0	0	0
3	K	5	0	0	0	0
3	M	2	0	0	0	0
3	O	2	0	0	0	0
All	All	5132	4741	4741	24	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (24) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:229:ARG:NH1	1:G:216:ASP:OD2	2.20	0.75
1:M:218:ARG:NE	1:M:276:GLU:OE2	2.31	0.64
1:K:251:LYS:NZ	1:K:255:GLU:OE1	2.32	0.62
1:A:277:GLY:O	1:O:234:SER:OG	2.15	0.61
1:E:227:TYR:O	1:E:231:VAL:HG13	2.08	0.54
1:G:231:VAL:HG22	1:I:253:TRP:HA	1.89	0.54
1:E:231:VAL:HG23	1:E:232:GLY:O	2.09	0.52
1:I:231:VAL:HG22	1:K:253:TRP:HA	1.90	0.52
1:C:265:VAL:O	3:C:301:HOH:O	2.19	0.51

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:F:231:MET:O	2:F:232:ARG:HB2	2.15	0.47
1:E:218:ARG:NE	1:E:276:GLU:OE2	2.47	0.46
1:O:221:LEU:HD13	1:O:268:LYS:HD2	1.98	0.45
1:I:274:TYR:O	2:J:231:MET:HE1	2.16	0.45
1:G:242:ILE:CG2	1:G:246:MET:HE2	2.46	0.45
1:I:218:ARG:NE	1:I:276:GLU:OE2	2.49	0.45
2:B:225:MET:N	2:B:226:PRO:HD2	2.32	0.45
1:C:227:TYR:CE2	1:C:231:VAL:HG21	2.53	0.44
1:E:228:LEU:HB3	1:E:238:TRP:CE2	2.53	0.44
1:E:274:TYR:OH	2:F:231:MET:HG3	2.18	0.43
1:C:234:SER:OG	1:E:277:GLY:O	2.29	0.43
1:C:211:THR:O	1:C:211:THR:OG1	2.36	0.42
2:N:231:MET:HA	2:N:231:MET:HE3	2.01	0.42
1:M:227:TYR:O	1:M:231:VAL:HG13	2.20	0.41
1:E:274:TYR:CZ	2:F:231:MET:HG3	2.56	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:K:215:GLU:OE2	1:O:252:LYS:NZ[1_454]	2.13	0.07

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	65/75 (87%)	65 (100%)	0	0	100	100
1	C	65/75 (87%)	65 (100%)	0	0	100	100
1	E	66/75 (88%)	65 (98%)	0	1 (2%)	8	2
1	G	64/75 (85%)	64 (100%)	0	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	I	65/75 (87%)	65 (100%)	0	0	100	100
1	K	64/75 (85%)	64 (100%)	0	0	100	100
1	M	65/75 (87%)	65 (100%)	0	0	100	100
1	O	63/75 (84%)	63 (100%)	0	0	100	100
2	B	6/8 (75%)	6 (100%)	0	0	100	100
2	D	4/8 (50%)	4 (100%)	0	0	100	100
2	F	5/8 (62%)	5 (100%)	0	0	100	100
2	H	5/8 (62%)	4 (80%)	1 (20%)	0	100	100
2	J	6/8 (75%)	5 (83%)	1 (17%)	0	100	100
2	L	5/8 (62%)	5 (100%)	0	0	100	100
2	N	4/8 (50%)	4 (100%)	0	0	100	100
2	P	5/8 (62%)	5 (100%)	0	0	100	100
All	All	557/664 (84%)	554 (100%)	2 (0%)	1 (0%)	43	36

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	E	211	THR

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	61/65 (94%)	59 (97%)	2 (3%)	33	24
1	C	61/65 (94%)	59 (97%)	2 (3%)	33	24
1	E	62/65 (95%)	58 (94%)	4 (6%)	15	6
1	G	60/65 (92%)	55 (92%)	5 (8%)	10	3
1	I	61/65 (94%)	60 (98%)	1 (2%)	55	52
1	K	60/65 (92%)	55 (92%)	5 (8%)	10	3

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	M	61/65 (94%)	58 (95%)	3 (5%)	22	11
1	O	60/65 (92%)	56 (93%)	4 (7%)	15	5
2	B	7/7 (100%)	6 (86%)	1 (14%)	3	0
2	D	5/7 (71%)	5 (100%)	0	100	100
2	F	6/7 (86%)	4 (67%)	2 (33%)	0	0
2	H	6/7 (86%)	5 (83%)	1 (17%)	2	0
2	J	7/7 (100%)	6 (86%)	1 (14%)	3	0
2	L	6/7 (86%)	5 (83%)	1 (17%)	2	0
2	N	5/7 (71%)	5 (100%)	0	100	100
2	P	6/7 (86%)	6 (100%)	0	100	100
All	All	534/576 (93%)	502 (94%)	32 (6%)	17	7

All (32) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	211	THR
1	A	213	ILE
2	B	231	MET
1	C	231	VAL
1	C	257	LYS
1	E	210	ASP
1	E	231	VAL
1	E	257	LYS
1	E	258	GLN
2	F	230	ARG
2	F	231	MET
1	G	213	ILE
1	G	231	VAL
1	G	255	GLU
1	G	257	LYS
1	G	273	GLN
2	H	231	MET
1	I	213	ILE
2	J	232	ARG
1	K	225	GLU
1	K	230	GLN
1	K	236	GLU
1	K	257	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	K	260	SER
2	L	231	MET
1	M	226	GLU
1	M	231	VAL
1	M	257	LYS
1	O	213	ILE
1	O	247	ASN
1	O	257	LYS
1	O	262	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (15) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	244	ASN
1	A	245	HIS
1	A	247	ASN
1	A	273	GLN
1	C	263	ASN
1	C	273	GLN
1	E	243	GLN
1	G	230	GLN
1	G	244	ASN
1	I	273	GLN
1	K	223	HIS
1	K	230	GLN
1	K	273	GLN
1	M	243	GLN
1	O	241	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

There are no ligands in this entry.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	67/75 (89%)	0.51	2 (2%) 52 59	42, 60, 86, 104	0
1	C	67/75 (89%)	0.51	3 (4%) 38 44	44, 65, 91, 100	0
1	E	68/75 (90%)	0.73	5 (7%) 20 23	49, 70, 93, 101	0
1	G	66/75 (88%)	0.55	2 (3%) 52 59	49, 71, 101, 127	0
1	I	67/75 (89%)	0.54	2 (2%) 52 59	45, 61, 89, 127	0
1	K	66/75 (88%)	0.71	3 (4%) 38 44	51, 74, 101, 142	0
1	M	67/75 (89%)	0.61	1 (1%) 72 78	51, 70, 86, 106	0
1	O	65/75 (86%)	0.63	3 (4%) 37 43	52, 73, 108, 120	0
2	B	8/8 (100%)	1.46	2 (25%) 2 1	61, 82, 105, 121	0
2	D	6/8 (75%)	2.19	3 (50%) 0 0	76, 91, 101, 108	0
2	F	7/8 (87%)	1.64	2 (28%) 1 1	61, 74, 96, 114	0
2	H	7/8 (87%)	1.83	3 (42%) 0 0	93, 103, 123, 158	0
2	J	8/8 (100%)	1.10	1 (12%) 8 9	61, 88, 100, 109	0
2	L	7/8 (87%)	2.14	4 (57%) 0 0	99, 112, 136, 191	0
2	N	6/8 (75%)	1.81	2 (33%) 1 0	69, 84, 101, 121	0
2	P	7/8 (87%)	1.49	2 (28%) 1 1	83, 86, 131, 134	0
All	All	589/664 (88%)	0.70	40 (6%) 23 27	42, 70, 106, 191	0

All (40) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	M	256	PHE	4.7
2	F	226	PRO	4.0
1	I	277	GLY	3.9
2	L	229	PHE	3.6
1	O	256	PHE	3.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	277	GLY	3.4
2	H	226	PRO	3.4
2	D	226	PRO	3.3
2	D	227	GLY	3.2
1	A	277	GLY	3.1
1	G	213	ILE	3.0
1	E	238	TRP	2.9
1	E	211	THR	2.8
2	L	231	MET	2.8
2	B	225	MET	2.7
1	K	213	ILE	2.6
1	E	231	VAL	2.5
2	L	226	PRO	2.5
2	N	227	GLY	2.5
1	E	256	PHE	2.4
1	I	214	PHE	2.4
1	O	253	TRP	2.4
2	P	229	PHE	2.4
1	K	214	PHE	2.3
2	N	226	PRO	2.3
2	D	228	CYS	2.2
2	P	231	MET	2.1
2	L	228	CYS	2.1
1	E	228	LEU	2.1
2	B	232	ARG	2.1
2	H	230	ARG	2.1
1	O	214	PHE	2.1
2	F	227	GLY	2.1
1	K	277	GLY	2.1
1	A	213	ILE	2.0
2	H	229	PHE	2.0
1	C	223	HIS	2.0
1	C	211	THR	2.0
1	G	212	GLN	2.0
2	J	226	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.