



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 17, 2026 – 06:07 PM UTC

PDB ID : 1TRB / pdb_00001trb
Title : CONVERGENT EVOLUTION OF SIMILAR FUNCTION IN TWO STRUCTURALLY DIVERGENT ENZYMES
Authors : Kuriyan, J.; Krishna, T.S.R.
Deposited on : 1991-09-07
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
Sidechain outliers	187428	11029 (2.00-2.00)
RSRZ outliers	180081	10067 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	320	

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 2611 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called THIOREDOXIN REDUCTASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	316	2394	1491	424	469	10	0	0	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	138	SER	CYS	conflict	UNP P0A9P4

- Molecule 2 is FLAVIN-ADENINE DINUCLEOTIDE (CCD ID: FAD) (formula: $C_{27}H_{33}N_9O_{15}P_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
2	A	1	53	27	9	15	2	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	164	Total 164	O 164	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: THIOREDOXIN REDUCTASE



4 Data and refinement statistics

Property	Value	Source
Space group	P 63 2 2	Depositor
Cell constants a, b, c, α , β , γ	123.10Å 123.10Å 81.40Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	(Not available) – 2.00 106.61 – 2.00	Depositor EDS
% Data completeness (in resolution range)	(Not available) ((Not available)-2.00) 87.1 (106.61-2.00)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	15.18 (at 2.00Å)	Xtriage
Refinement program	X-PLOR	Depositor
R, R_{free}	0.177 , (Not available) 0.174 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	20.4	Xtriage
Anisotropy	0.282	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 72.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.35$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	2611	wwPDB-VP
Average B, all atoms (Å ²)	26.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.59% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.25	15/2433 (0.6%)	1.87	53/3295 (1.6%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (15) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	202	HIS	CD2-NE2	-7.27	1.29	1.37
1	A	241	VAL	CA-CB	7.08	1.62	1.54
1	A	270	HIS	CD2-NE2	-6.63	1.30	1.37
1	A	172	HIS	CD2-NE2	-6.49	1.30	1.37
1	A	22	VAL	CA-CB	6.21	1.62	1.54
1	A	71	HIS	CD2-NE2	-6.04	1.31	1.37
1	A	175	HIS	CD2-NE2	-5.97	1.31	1.37
1	A	202	HIS	CG-ND1	-5.80	1.31	1.38
1	A	5	HIS	CG-ND1	-5.75	1.31	1.38
1	A	311	ARG	NE-CZ	5.54	1.39	1.33
1	A	69	HIS	CD2-NE2	-5.48	1.31	1.37
1	A	251	ILE	CB-CG1	-5.30	1.42	1.53
1	A	5	HIS	CD2-NE2	-5.26	1.32	1.37
1	A	251	ILE	CA-CB	5.23	1.63	1.54
1	A	270	HIS	CG-ND1	-5.08	1.32	1.38

All (53) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	198	ASN	N-CA-C	13.64	127.65	111.82
1	A	224	ASP	CA-CB-CG	12.71	125.31	112.60
1	A	67	ARG	NE-CZ-NH2	-12.07	108.34	119.20
1	A	196	ASN	CA-CB-CG	10.47	123.07	112.60
1	A	67	ARG	CB-CG-CD	-8.74	91.20	111.30
1	A	251	ILE	CA-CB-CG2	8.71	125.31	110.50
1	A	213	ASP	CA-CB-CG	7.21	119.81	112.60
1	A	251	ILE	CA-CB-CG1	-7.17	98.22	110.40
1	A	230	ASN	N-CA-C	7.14	118.71	111.07
1	A	251	ILE	CB-CG1-CD1	-6.91	99.29	113.80
1	A	67	ARG	NE-CZ-NH1	6.81	128.31	121.50
1	A	172	HIS	CB-CG-CD2	-6.75	122.43	131.20
1	A	146	GLN	OE1-CD-NE2	-6.66	115.94	122.60
1	A	197	GLY	O-C-N	6.65	131.35	122.70
1	A	311	ARG	NE-CZ-NH2	-6.62	113.24	119.20
1	A	35	THR	N-CA-C	6.56	119.43	111.82
1	A	50	GLU	CA-CB-CG	6.37	126.83	114.10
1	A	155	ASN	CA-CB-CG	-6.21	106.39	112.60
1	A	197	GLY	CA-C-N	6.18	129.70	120.31
1	A	197	GLY	C-N-CA	6.18	129.70	120.31
1	A	90	GLN	OE1-CD-NE2	-6.13	116.47	122.60
1	A	220	VAL	N-CA-CB	-6.11	101.34	111.36
1	A	251	ILE	N-CA-CB	-6.09	98.72	110.78
1	A	232	GLU	CA-CB-CG	5.93	125.95	114.10
1	A	240	PHE	CA-CB-CG	-5.75	108.05	113.80
1	A	311	ARG	NE-CZ-NH1	5.75	127.25	121.50
1	A	236	VAL	N-CA-CB	-5.73	100.74	111.62
1	A	76	GLU	N-CA-C	5.47	119.25	112.24
1	A	146	GLN	CG-CD-NE2	5.44	124.56	116.40
1	A	196	ASN	OD1-CG-ND2	-5.43	117.17	122.60
1	A	226	GLN	OE1-CD-NE2	-5.42	117.18	122.60
1	A	240	PHE	CA-C-N	-5.40	115.61	122.37
1	A	240	PHE	C-N-CA	-5.40	115.61	122.37
1	A	22	VAL	N-CA-CB	5.36	118.63	110.58
1	A	87	VAL	N-CA-CB	-5.36	101.89	111.92
1	A	172	HIS	CB-CG-ND1	5.35	130.72	122.70
1	A	229	ASP	CA-C-N	5.32	127.36	120.44
1	A	229	ASP	C-N-CA	5.32	127.36	120.44
1	A	87	VAL	CG1-CB-CG2	5.32	122.49	110.80
1	A	192	ASP	CA-CB-CG	5.28	117.88	112.60
1	A	241	VAL	CB-CA-C	5.27	117.34	110.96
1	A	148	VAL	N-CA-CB	-5.26	102.09	111.92
1	A	291	ILE	N-CA-C	5.25	116.08	111.56

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	189	ARG	NE-CZ-NH2	5.24	123.91	119.20
1	A	231	ILE	N-CA-C	5.21	120.17	109.34
1	A	6	SER	CA-CB-OG	-5.19	100.72	111.10
1	A	226	GLN	CB-CG-CD	5.17	121.38	112.60
1	A	162	LEU	CD1-CG-CD2	5.15	122.13	110.80
1	A	202	HIS	CB-CG-CD2	-5.14	124.51	131.20
1	A	254	GLY	N-CA-C	-5.14	108.52	115.36
1	A	148	VAL	CA-CB-CG2	-5.13	101.68	110.40
1	A	236	VAL	CB-CA-C	5.04	120.25	111.18
1	A	270	HIS	CB-CG-CD2	-5.04	124.65	131.20

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	311	ARG	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2394	0	2356	41	0
2	A	53	0	31	0	0
3	A	164	0	0	6	0
All	All	2611	0	2387	41	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (41) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:184:LYS:HD2	1:A:185:ILE:H	1.61	0.66
1:A:87:VAL:HG13	1:A:89:LEU:HD13	1.79	0.64
1:A:184:LYS:HD2	1:A:185:ILE:N	2.13	0.63

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:272:ASN:H	1:A:311:ARG:HH22	1.46	0.62
1:A:183:GLU:HB3	1:A:184:LYS:HZ3	1.66	0.61
1:A:2:THR:HG22	3:A:644:HOH:O	1.99	0.61
1:A:85:ASN:HA	1:A:251:ILE:HD11	1.82	0.61
1:A:162:LEU:HG	1:A:189:ARG:NH1	2.15	0.61
1:A:155:ASN:HD22	1:A:186:LEU:HD22	1.66	0.61
1:A:183:GLU:HB3	1:A:184:LYS:NZ	2.15	0.60
1:A:153:GLY:O	1:A:181:ARG:HG3	2.02	0.59
1:A:184:LYS:HG2	3:A:724:HOH:O	2.04	0.58
1:A:162:LEU:HG	1:A:189:ARG:CZ	2.33	0.58
1:A:155:ASN:ND2	1:A:186:LEU:HD22	2.19	0.57
1:A:87:VAL:HG22	1:A:96:LEU:HG	1.87	0.57
1:A:207:LEU:HD11	1:A:220:VAL:HG22	1.88	0.55
1:A:245:HIS:HE1	3:A:653:HOH:O	1.90	0.55
1:A:42:GLN:HG2	3:A:712:HOH:O	2.06	0.54
1:A:3:THR:HG23	1:A:102:GLU:HG2	1.89	0.54
1:A:223:ARG:NH1	1:A:224:ASP:HB2	2.24	0.53
1:A:204:ASN:HB3	1:A:225:THR:HA	1.90	0.53
1:A:189:ARG:HH11	1:A:193:LYS:HE3	1.74	0.52
1:A:6:SER:O	1:A:105:CYS:HA	2.09	0.52
1:A:223:ARG:O	1:A:224:ASP:HB3	2.09	0.52
1:A:208:GLU:HG3	1:A:223:ARG:HG2	1.93	0.51
1:A:158:VAL:O	1:A:162:LEU:HB2	2.10	0.51
1:A:174:ILE:HD13	1:A:222:LEU:HD11	1.95	0.48
1:A:135:CYS:HG	1:A:138:SER:HG	0.49	0.47
1:A:158:VAL:HG12	1:A:162:LEU:HD22	1.97	0.46
1:A:59:LEU:HD13	1:A:67:ARG:NH2	2.29	0.46
1:A:220:VAL:HG13	1:A:222:LEU:HD13	1.96	0.46
1:A:94:PHE:O	1:A:104:THR:HA	2.16	0.46
1:A:220:VAL:HG12	1:A:234:LEU:HB2	1.97	0.46
1:A:125:GLU:HG2	3:A:674:HOH:O	2.17	0.45
1:A:39:LYS:NZ	1:A:81:PHE:HB3	2.34	0.43
1:A:162:LEU:CG	1:A:189:ARG:NH1	2.81	0.42
1:A:159:GLU:HG3	1:A:186:LEU:HD21	2.01	0.42
1:A:224:ASP:O	1:A:225:THR:HB	2.19	0.42
1:A:43:LEU:HB2	3:A:615:HOH:O	2.21	0.41
1:A:156:THR:O	1:A:160:GLU:HG2	2.21	0.40
1:A:162:LEU:HG	1:A:189:ARG:HD2	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	314/320 (98%)	296 (94%)	16 (5%)	2 (1%)	21	17

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	224	ASP
1	A	231	ILE

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	251/253 (99%)	218 (87%)	33 (13%)	4	2

All (33) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	2	THR
1	A	11	LEU
1	A	22	VAL
1	A	43	LEU
1	A	64	LEU
1	A	86	LYS
1	A	87	VAL
1	A	89	LEU
1	A	125	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	148	VAL
1	A	162	LEU
1	A	164	LEU
1	A	173	LEU
1	A	181	ARG
1	A	184	LYS
1	A	186	LEU
1	A	189	ARG
1	A	196	ASN
1	A	198	ASN
1	A	201	LEU
1	A	205	ARG
1	A	207	LEU
1	A	215	MET
1	A	220	VAL
1	A	222	LEU
1	A	225	THR
1	A	230	ASN
1	A	234	LEU
1	A	236	VAL
1	A	241	VAL
1	A	251	ILE
1	A	258	LEU
1	A	291	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	91	ASN
1	A	145	ASN
1	A	155	ASN
1	A	166	ASN
1	A	198	ASN
1	A	245	HIS
1	A	248	ASN
1	A	272	ASN
1	A	275	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	FAD	A	500	-	58,58,58	1.12	3 (5%)	85,89,89	0.95	4 (4%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FAD	A	500	-	-	3/34/50/50	0/6/6/6

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	500	FAD	PA-O3P	-2.97	1.56	1.59
2	A	500	FAD	C5A-C4A	-2.64	1.34	1.39
2	A	500	FAD	C4-N3	-2.00	1.35	1.38

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	500	FAD	O3P-PA-O1A	-2.34	103.66	110.70

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	500	FAD	O2A-PA-O3P	2.27	113.41	107.27
2	A	500	FAD	C1B-N9A-C8A	2.15	131.88	127.09
2	A	500	FAD	O4'-C4'-C3'	-2.00	104.56	109.25

There are no chirality outliers.


All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	500	FAD	P-O3P-PA-O1A
2	A	500	FAD	PA-O3P-P-O5'
2	A	500	FAD	P-O3P-PA-O2A

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	316/320 (98%)	0.06	17 (5%) 31 30	8, 21, 55, 96	0

All (17) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	231	ILE	6.0
1	A	226	GLN	5.1
1	A	225	THR	4.8
1	A	189	ARG	4.7
1	A	228	SER	4.5
1	A	224	ASP	4.5
1	A	229	ASP	4.4
1	A	196	ASN	3.9
1	A	227	ASN	3.8
1	A	100	ASN	3.5
1	A	215	MET	3.0
1	A	223	ARG	3.0
1	A	178	ASP	2.8
1	A	181	ARG	2.6
1	A	179	GLY	2.2
1	A	195	GLU	2.1
1	A	198	ASN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	FAD	A	500	53/53	0.98	0.04	8,12,20,21	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.