



Full wwPDB NMR Structure Validation Report ⓘ

Mar 9, 2026 – 01:07 PM UTC

PDB ID : 1TRL / pdb_00001trl
Title : NMR SOLUTION STRUCTURE OF THE C-TERMINAL FRAGMENT 255-316 OF THERMOLYSIN: A DIMER FORMED BY SUBUNITS HAVING THE NATIVE STRUCTURE
Authors : Rico, M.; Jimenez, M.A.; Gonzalez, C.; De Filippis, V.; Fontana, A.
Deposited on : 1994-09-02

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
wwPDB-RCI	:	v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV	:	Wang et al. (2010)
wwPDB-ShiftChecker	:	v1.2
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment was not calculated.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	229148	14424
Ramachandran outliers	224038	12848
Sidechain outliers	223484	12823

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	62	
1	B	62	

2 Ensemble composition and analysis

This entry contains 8 models. Model 4 is the overall representative, medoid model (most similar to other models).

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:262-A:313, B:262-B:313 (104)	0.69	4

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 2 clusters and 1 single-model cluster was found.

Cluster number	Models
1	1, 2, 3, 4, 6
2	7, 8
Single-model clusters	5

3 Entry composition

There is only 1 type of molecule in this entry. The entry contains 1168 atoms, of which 232 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called THERMOLYSIN FRAGMENT 255 - 316.


Mol	Chain	Residues	Atoms					Trace
1	A	62	Total	C	H	N	O	0
			584	296	116	81	91	
1	B	62	Total	C	H	N	O	0
			584	296	116	81	91	

4 Residue-property plots [i](#)


4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: THERMOLYSIN FRAGMENT 255 - 316



- Molecule 1: THERMOLYSIN FRAGMENT 255 - 316




4.2 Scores per residue for each member of the ensemble


Colouring as in section 4.1 above.

4.2.1 Score per residue for model 1

- Molecule 1: THERMOLYSIN FRAGMENT 255 - 316



- Molecule 1: THERMOLYSIN FRAGMENT 255 - 316




4.2.2 Score per residue for model 2

- Molecule 1: THERMOLYSIN FRAGMENT 255 - 316




- Molecule 1: THERMOLYSIN FRAGMENT 255 - 316




4.2.3 Score per residue for model 3

- Molecule 1: THERMOLYSIN FRAGMENT 255 - 316




- Molecule 1: THERMOLYSIN FRAGMENT 255 - 316




4.2.4 Score per residue for model 4 (medoid)

- Molecule 1: THERMOLYSIN FRAGMENT 255 - 316




- Molecule 1: THERMOLYSIN FRAGMENT 255 - 316




4.2.5 Score per residue for model 5

- Molecule 1: THERMOLYSIN FRAGMENT 255 - 316




- Molecule 1: THERMOLYSIN FRAGMENT 255 - 316




4.2.6 Score per residue for model 6

- Molecule 1: THERMOLYSIN FRAGMENT 255 - 316




- Molecule 1: THERMOLYSIN FRAGMENT 255 - 316




4.2.7 Score per residue for model 7

- Molecule 1: THERMOLYSIN FRAGMENT 255 - 316




- Molecule 1: THERMOLYSIN FRAGMENT 255 - 316




4.2.8 Score per residue for model 8

- Molecule 1: THERMOLYSIN FRAGMENT 255 - 316



- Molecule 1: THERMOLYSIN FRAGMENT 255 - 316



5 Refinement protocol and experimental data overview

Of the ? calculated structures, 8 were deposited, based on the following criterion: ?.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
GROMOS	refinement	

No chemical shift data was provided.

6 Model quality i

6.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	0.91±0.02	0±0/405 (0.0± 0.0%)	1.63±0.10	2±2/550 (0.5± 0.3%)
1	B	0.91±0.02	0±0/405 (0.0± 0.0%)	1.62±0.06	3±2/550 (0.6± 0.3%)
All	All	0.91	0/6480 (0.0%)	1.63	45/8800 (0.5%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
1	A	0.0±0.0	3.1±1.5
1	B	0.0±0.0	2.6±1.6
All	All	0	46

There are no bond-length outliers.

All unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	301	GLN	CG-CD-NE2	-13.50	96.14	116.40	4	1
1	A	301	GLN	CG-CD-OE1	10.75	142.31	120.80	4	1
1	A	308	GLN	OE1-CD-NE2	-8.98	113.62	122.60	8	1
1	A	298	SER	N-CA-C	7.89	119.67	111.14	4	4
1	B	298	SER	N-CA-C	7.62	119.49	111.03	4	4
1	B	267	PHE	CA-CB-CG	-7.35	106.45	113.80	6	6
1	A	267	PHE	CA-CB-CG	-7.30	106.50	113.80	3	6
1	B	280	ASN	OD1-CG-ND2	-7.21	115.39	122.60	7	1
1	B	285	ARG	CD-NE-CZ	6.72	133.80	124.40	4	1
1	A	267	PHE	N-CA-C	6.59	118.55	111.36	3	2
1	B	284	LEU	N-CA-CB	-5.95	101.39	110.01	5	3
1	B	285	ARG	CG-CD-NE	-5.78	99.29	112.00	4	1
1	B	280	ASN	CA-C-N	5.75	127.92	120.44	7	1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	B	280	ASN	C-N-CA	5.75	127.92	120.44	7	1
1	B	296	TYR	N-CA-C	5.63	119.33	111.56	3	1
1	B	294	ASP	N-CA-C	5.50	117.28	111.28	5	1
1	A	263	LEU	CA-C-N	5.38	125.95	119.98	3	1
1	A	263	LEU	C-N-CA	5.38	125.95	119.98	3	1
1	A	285	ARG	NE-CZ-NH2	5.30	123.97	119.20	7	1
1	B	308	GLN	N-CA-CB	-5.27	102.08	109.94	4	1
1	A	279	SER	N-CA-C	5.27	118.15	109.72	3	2
1	B	279	SER	N-CA-C	5.26	117.65	110.55	3	1
1	B	267	PHE	N-CA-C	5.16	116.90	111.28	2	3

There are no chirality outliers.

All unique planar outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Group	Models (Total)
1	A	268	TYR	Sidechain	5
1	A	296	TYR	Sidechain	5
1	B	274	TYR	Sidechain	5
1	B	296	TYR	Sidechain	5
1	B	268	TYR	Sidechain	4
1	B	281	PHE	Sidechain	3
1	A	281	PHE	Sidechain	3
1	A	310	PHE	Sidechain	2
1	A	274	TYR	Sidechain	2
1	A	279	SER	Mainchain	1
1	A	267	PHE	Sidechain	1
1	A	306	VAL	Mainchain	1
1	B	277	PRO	Mainchain	1
1	B	312	ALA	Mainchain	1
1	A	286	ALA	Mainchain	1
1	B	310	PHE	Sidechain	1
1	B	313	VAL	Mainchain	1
1	A	278	THR	Mainchain	1
1	A	308	GLN	Sidechain	1
1	A	309	ALA	Mainchain	1
1	A	313	VAL	Mainchain	1

6.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	398	96	399	4±2
1	B	398	96	399	3±1
All	All	6368	1536	6384	41

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:264:GLY:HA2	1:B:263:LEU:HB3	0.69	1.64	4	2
1:A:264:GLY:HA2	1:B:263:LEU:HB2	0.66	1.64	8	1
1:A:263:LEU:HA	1:A:266:ILE:HD12	0.62	1.70	6	6
1:B:269:ARG:CZ	1:B:273:GLN:HE22	0.59	2.09	6	1
1:A:263:LEU:HB2	1:B:264:GLY:HA2	0.58	1.76	8	1
1:A:293:THR:HG23	1:A:298:SER:HB3	0.57	1.76	5	3
1:B:293:THR:HG23	1:B:298:SER:HB3	0.54	1.80	5	2
1:A:263:LEU:HD22	1:B:264:GLY:HA2	0.52	1.80	6	3
1:B:265:LYS:HB3	1:B:295:LEU:HD13	0.50	1.82	1	1
1:A:300:SER:HB3	1:A:303:VAL:HG23	0.49	1.84	4	1
1:A:281:PHE:HB2	1:B:279:SER:HB2	0.49	1.84	5	1
1:B:263:LEU:HA	1:B:266:ILE:HD12	0.47	1.85	3	4
1:A:309:ALA:O	1:A:313:VAL:HG23	0.47	2.10	1	2
1:B:266:ILE:HD11	1:B:302:GLU:HG2	0.46	1.85	2	1
1:A:305:SER:HA	1:A:308:GLN:OE1	0.44	2.12	8	1
1:A:266:ILE:HG21	1:A:306:VAL:HG21	0.44	1.90	5	1
1:A:275:LEU:HB3	1:B:313:VAL:HG13	0.43	1.89	1	1
1:A:303:VAL:HG12	1:A:307:LYS:HE3	0.43	1.90	4	1
1:A:297:GLY:O	1:A:300:SER:HB2	0.43	2.13	2	1
1:B:271:LEU:HA	1:B:275:LEU:HB2	0.42	1.90	7	1
1:A:264:GLY:HA2	1:B:263:LEU:HD12	0.41	1.90	5	1
1:B:269:ARG:NH2	1:B:273:GLN:HE22	0.41	2.12	6	1
1:A:275:LEU:HG	1:A:279:SER:OG	0.41	2.15	6	1
1:B:298:SER:HA	1:B:303:VAL:HG21	0.41	1.91	7	1
1:A:293:THR:HG22	1:A:298:SER:HA	0.41	1.91	8	1
1:A:271:LEU:HA	1:A:275:LEU:HB2	0.41	1.93	4	1

6.3 Torsion angles

6.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	52/62 (84%)	48±2 (92±3%)	3±2 (6±3%)	1±1 (2±2%)	9	53
1	B	52/62 (84%)	48±1 (91±2%)	4±1 (8±2%)	0±0 (1±1%)	20	70
All	All	832/992 (84%)	764 (92%)	58 (7%)	10 (1%)	13	61

All 7 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	299	THR	3
1	B	297	GLY	2
1	A	296	TYR	1
1	A	297	GLY	1
1	A	298	SER	1
1	A	279	SER	1
1	B	264	GLY	1

6.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	42/49 (86%)	38±1 (91±2%)	4±1 (9±2%)	12	58
1	B	42/49 (86%)	39±1 (93±3%)	3±1 (7±3%)	16	65
All	All	672/784 (86%)	620 (92%)	52 (8%)	14	61

All 30 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	271	LEU	5
1	B	283	GLN	4
1	A	265	LYS	4
1	A	301	GLN	3
1	A	275	LEU	3
1	A	279	SER	2
1	B	271	LEU	2
1	B	290	GLN	2
1	B	298	SER	2
1	A	277	PRO	2
1	B	263	LEU	2
1	B	275	LEU	2
1	B	273	GLN	2
1	A	283	GLN	1
1	A	262	LYS	1
1	B	285	ARG	1
1	A	280	ASN	1
1	B	294	ASP	1
1	A	269	ARG	1
1	A	276	THR	1
1	A	299	THR	1
1	A	302	GLU	1
1	A	273	GLN	1
1	B	305	SER	1
1	A	263	LEU	1
1	B	301	GLN	1
1	B	302	GLU	1
1	B	307	LYS	1
1	A	293	THR	1
1	B	299	THR	1

6.3.3 RNA ⓘ

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

6.6 Ligand geometry

There are no ligands in this entry.

6.7 Other polymers

There are no such molecules in this entry.

6.8 Polymer linkage issues

There are no chain breaks in this entry.

7 Chemical shift validation

No chemical shift data were provided