



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 1, 2026 – 02:19 AM UTC

PDB ID : 4TRR / pdb_00004trr
Title : Crystal structure of a putative Putative D-beta-hydroxybutyrate dehydrogenase from Burkholderia cenocepacia J2315
Authors : Seattle Structural Genomics Center for Infectious Disease (SSGCID)
Deposited on : 2014-06-17
Resolution : 1.90 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7789 (1.90-1.90)
Clashscore	190562	8410 (1.90-1.90)
Ramachandran outliers	187476	8333 (1.90-1.90)
Sidechain outliers	187428	8333 (1.90-1.90)
RSRZ outliers	180081	7790 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	269	89% 6% 5%
1	B	269	88% 8% .
1	C	269	91% . 5%
1	D	269	88% 7% 5%
1	E	269	86% . 10%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	269	 84%10%6%
1	G	269	 78%6%16%
1	H	269	 87%8%5%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 15759 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Putative D-beta-hydroxybutyrate dehydrogenase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	256	Total	C	N	O	S	0	5	0
			1880	1193	322	358	7			
1	B	260	Total	C	N	O	S	0	6	0
			1930	1219	332	372	7			
1	C	256	Total	C	N	O	S	0	5	0
			1871	1186	322	356	7			
1	D	256	Total	C	N	O	S	0	6	0
			1895	1202	326	360	7			
1	E	242	Total	C	N	O	S	0	0	0
			1703	1073	294	329	7			
1	F	254	Total	C	N	O	S	0	1	0
			1796	1134	312	343	7			
1	G	226	Total	C	N	O	S	0	1	0
			1564	986	274	298	6			
1	H	256	Total	C	N	O	S	0	3	0
			1834	1158	314	355	7			

There are 64 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-7	MET	-	initiating methionine	UNP B4EG08
A	-6	ALA	-	expression tag	UNP B4EG08
A	-5	HIS	-	expression tag	UNP B4EG08
A	-4	HIS	-	expression tag	UNP B4EG08
A	-3	HIS	-	expression tag	UNP B4EG08
A	-2	HIS	-	expression tag	UNP B4EG08
A	-1	HIS	-	expression tag	UNP B4EG08
A	0	HIS	-	expression tag	UNP B4EG08
B	-7	MET	-	initiating methionine	UNP B4EG08
B	-6	ALA	-	expression tag	UNP B4EG08
B	-5	HIS	-	expression tag	UNP B4EG08
B	-4	HIS	-	expression tag	UNP B4EG08
B	-3	HIS	-	expression tag	UNP B4EG08

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
B	-2	HIS	-	expression tag	UNP B4EG08
B	-1	HIS	-	expression tag	UNP B4EG08
B	0	HIS	-	expression tag	UNP B4EG08
C	-7	MET	-	initiating methionine	UNP B4EG08
C	-6	ALA	-	expression tag	UNP B4EG08
C	-5	HIS	-	expression tag	UNP B4EG08
C	-4	HIS	-	expression tag	UNP B4EG08
C	-3	HIS	-	expression tag	UNP B4EG08
C	-2	HIS	-	expression tag	UNP B4EG08
C	-1	HIS	-	expression tag	UNP B4EG08
C	0	HIS	-	expression tag	UNP B4EG08
D	-7	MET	-	initiating methionine	UNP B4EG08
D	-6	ALA	-	expression tag	UNP B4EG08
D	-5	HIS	-	expression tag	UNP B4EG08
D	-4	HIS	-	expression tag	UNP B4EG08
D	-3	HIS	-	expression tag	UNP B4EG08
D	-2	HIS	-	expression tag	UNP B4EG08
D	-1	HIS	-	expression tag	UNP B4EG08
D	0	HIS	-	expression tag	UNP B4EG08
E	-7	MET	-	initiating methionine	UNP B4EG08
E	-6	ALA	-	expression tag	UNP B4EG08
E	-5	HIS	-	expression tag	UNP B4EG08
E	-4	HIS	-	expression tag	UNP B4EG08
E	-3	HIS	-	expression tag	UNP B4EG08
E	-2	HIS	-	expression tag	UNP B4EG08
E	-1	HIS	-	expression tag	UNP B4EG08
E	0	HIS	-	expression tag	UNP B4EG08
F	-7	MET	-	initiating methionine	UNP B4EG08
F	-6	ALA	-	expression tag	UNP B4EG08
F	-5	HIS	-	expression tag	UNP B4EG08
F	-4	HIS	-	expression tag	UNP B4EG08
F	-3	HIS	-	expression tag	UNP B4EG08
F	-2	HIS	-	expression tag	UNP B4EG08
F	-1	HIS	-	expression tag	UNP B4EG08
F	0	HIS	-	expression tag	UNP B4EG08
G	-7	MET	-	initiating methionine	UNP B4EG08
G	-6	ALA	-	expression tag	UNP B4EG08
G	-5	HIS	-	expression tag	UNP B4EG08
G	-4	HIS	-	expression tag	UNP B4EG08
G	-3	HIS	-	expression tag	UNP B4EG08
G	-2	HIS	-	expression tag	UNP B4EG08
G	-1	HIS	-	expression tag	UNP B4EG08

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
G	0	HIS	-	expression tag	UNP B4EG08
H	-7	MET	-	initiating methionine	UNP B4EG08
H	-6	ALA	-	expression tag	UNP B4EG08
H	-5	HIS	-	expression tag	UNP B4EG08
H	-4	HIS	-	expression tag	UNP B4EG08
H	-3	HIS	-	expression tag	UNP B4EG08
H	-2	HIS	-	expression tag	UNP B4EG08
H	-1	HIS	-	expression tag	UNP B4EG08
H	0	HIS	-	expression tag	UNP B4EG08

- Molecule 2 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	C	1	Total	O	S	0	0
			5	4	1		
2	D	1	Total	O	S	0	0
			5	4	1		
2	E	1	Total	O	S	0	0
			5	4	1		
2	H	1	Total	O	S	0	0
			5	4	1		

- Molecule 3 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			4	2	2		
3	D	1	Total	C	O	0	0
			4	2	2		
3	D	1	Total	C	O	0	0
			4	2	2		
3	F	1	Total	C	O	0	0
			4	2	2		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	249	Total	O	0	0
			249	249		
4	B	230	Total	O	0	0
			230	230		
4	C	215	Total	O	0	0
			215	215		
4	D	203	Total	O	0	0
			203	203		
4	E	93	Total	O	0	0
			93	93		
4	F	66	Total	O	0	0
			66	66		
4	G	42	Total	O	0	0
			42	42		
4	H	142	Total	O	0	0
			142	142		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Putative D-beta-hydroxybutyrate dehydrogenase

Chain A: 



- Molecule 1: Putative D-beta-hydroxybutyrate dehydrogenase

Chain B: 



- Molecule 1: Putative D-beta-hydroxybutyrate dehydrogenase

Chain C: 



- Molecule 1: Putative D-beta-hydroxybutyrate dehydrogenase

Chain D: 




- Molecule 1: Putative D-beta-hydroxybutyrate dehydrogenase

Chain E: 





- Molecule 1: Putative D-beta-hydroxybutyrate dehydrogenase


Chain F: 



● Molecule 1: Putative D-beta-hydroxybutyrate dehydrogenase



● Molecule 1: Putative D-beta-hydroxybutyrate dehydrogenase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	103.22Å 91.95Å 116.56Å 90.00° 92.94° 90.00°	Depositor
Resolution (Å)	48.05 – 1.90 48.05 – 1.90	Depositor EDS
% Data completeness (in resolution range)	98.6 (48.05-1.90) 98.6 (48.05-1.90)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.61 (at 1.90Å)	Xtriage
Refinement program	PHENIX (phenix.refine: dev_1702)	Depositor
R, R_{free}	0.183 , 0.222 0.185 , 0.224	Depositor DCC
R_{free} test set	8452 reflections (4.93%)	wwPDB-VP
Wilson B-factor (Å ²)	19.5	Xtriage
Anisotropy	0.504	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 33.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.43$, $\langle L^2 \rangle = 0.25$	Xtriage
Estimated twinning fraction	0.095 for h,-k,-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	15759	wwPDB-VP
Average B, all atoms (Å ²)	33.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.22% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.54	0/1925	0.79	0/2612
1	B	0.56	0/1977	0.77	0/2678
1	C	0.54	0/1916	0.77	0/2603
1	D	0.52	0/1942	0.77	0/2632
1	E	0.38	0/1729	0.75	0/2354
1	F	0.38	0/1828	0.76	0/2492
1	G	0.35	0/1593	0.73	0/2173
1	H	0.41	0/1873	0.76	0/2553
All	All	0.47	0/14783	0.76	0/20097

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1880	0	1879	11	0
1	B	1930	0	1933	16	0
1	C	1871	0	1849	5	0
1	D	1895	0	1894	12	0
1	E	1703	0	1630	6	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	F	1796	0	1723	17	0
1	G	1564	0	1460	8	0
1	H	1834	0	1758	13	0
2	A	5	0	0	0	0
2	B	5	0	0	0	0
2	C	5	0	0	0	0
2	D	5	0	0	0	0
2	E	5	0	0	0	0
2	H	5	0	0	0	0
3	A	4	0	6	0	0
3	D	8	0	12	0	0
3	F	4	0	6	0	0
4	A	249	0	0	3	0
4	B	230	0	0	6	0
4	C	215	0	0	1	0
4	D	203	0	0	1	0
4	E	93	0	0	1	0
4	F	66	0	0	1	0
4	G	42	0	0	1	0
4	H	142	0	0	2	0
All	All	15759	0	14150	83	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 83 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:G:191:THR:HB	1:G:192:PRO:HD2	1.66	0.77
1:F:134:ARG:NH1	4:F:430:HOH:O	2.20	0.74
1:B:195:ASP:OD2	4:B:401:HOH:O	2.11	0.68
1:E:40:GLN:HB2	1:E:61:ALA:HB2	1.75	0.68
1:B:231:ASP:OD2	4:B:555:HOH:O	2.11	0.67

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	257/269 (96%)	253 (98%)	4 (2%)	0	100	100
1	B	263/269 (98%)	258 (98%)	4 (2%)	1 (0%)	30	22
1	C	257/269 (96%)	253 (98%)	4 (2%)	0	100	100
1	D	257/269 (96%)	253 (98%)	3 (1%)	1 (0%)	30	22
1	E	238/269 (88%)	233 (98%)	4 (2%)	1 (0%)	30	22
1	F	251/269 (93%)	244 (97%)	6 (2%)	1 (0%)	30	22
1	G	221/269 (82%)	214 (97%)	5 (2%)	2 (1%)	14	6
1	H	255/269 (95%)	251 (98%)	4 (2%)	0	100	100
All	All	1999/2152 (93%)	1959 (98%)	34 (2%)	6 (0%)	36	29

5 of 6 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	G	259	PHE
1	B	259	PHE
1	D	259	PHE
1	E	259	PHE
1	F	259	PHE

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	192/207 (93%)	190 (99%)	2 (1%)	68	70

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	199/207 (96%)	198 (100%)	1 (0%)	81	84
1	C	188/207 (91%)	185 (98%)	3 (2%)	55	54
1	D	193/207 (93%)	191 (99%)	2 (1%)	68	70
1	E	163/207 (79%)	162 (99%)	1 (1%)	78	81
1	F	173/207 (84%)	173 (100%)	0	100	100
1	G	142/207 (69%)	140 (99%)	2 (1%)	59	59
1	H	179/207 (86%)	178 (99%)	1 (1%)	78	81
All	All	1429/1656 (86%)	1417 (99%)	12 (1%)	73	75

5 of 12 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	127	LYS
1	E	220	ASN
1	H	195	ASP
1	G	103	SER
1	C	103	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 11 such sidechains are listed below:

Mol	Chain	Res	Type
1	E	111	GLN
1	H	40	GLN
1	H	147	HIS
1	H	111	GLN
1	C	261	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

10 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	EDO	D	302	-	3,3,3	0.49	0	2,2,2	0.39	0
3	EDO	F	300	-	3,3,3	0.44	0	2,2,2	0.34	0
2	SO4	A	301	-	4,4,4	0.27	0	6,6,6	0.09	0
3	EDO	D	300	-	3,3,3	0.45	0	2,2,2	0.34	0
2	SO4	E	301	-	4,4,4	0.25	0	6,6,6	0.11	0
2	SO4	B	301	-	4,4,4	0.29	0	6,6,6	0.24	0
2	SO4	C	301	-	4,4,4	0.29	0	6,6,6	0.12	0
2	SO4	H	301	-	4,4,4	0.30	0	6,6,6	0.12	0
2	SO4	D	301	-	4,4,4	0.25	0	6,6,6	0.15	0
3	EDO	A	302	-	3,3,3	0.41	0	2,2,2	0.30	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	EDO	D	302	-	-	0/1/1/1	-
3	EDO	D	300	-	-	0/1/1/1	-
3	EDO	F	300	-	-	0/1/1/1	-
3	EDO	A	302	-	-	1/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	302	EDO	O1-C1-C2-O2

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	256/269 (95%)	-1.46	0 100 100	7, 20, 48, 84	5 (1%)
1	B	260/269 (96%)	-1.50	0 100 100	8, 19, 42, 60	6 (2%)
1	C	256/269 (95%)	-1.43	0 100 100	7, 22, 49, 69	5 (1%)
1	D	256/269 (95%)	-1.47	0 100 100	8, 21, 45, 69	6 (2%)
1	E	242/269 (89%)	-1.04	0 100 100	26, 39, 60, 82	0
1	F	254/269 (94%)	-0.85	0 100 100	21, 44, 62, 77	1 (0%)
1	G	226/269 (84%)	-0.64	0 100 100	25, 50, 80, 97	1 (0%)
1	H	256/269 (95%)	-1.17	0 100 100	16, 34, 58, 74	3 (1%)
All	All	2006/2152 (93%)	-1.21	0 100 100	7, 32, 63, 97	27 (1%)

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	SO4	E	301	5/5	0.98	0.04	46,54,58,64	0
3	EDO	F	300	4/4	0.98	0.06	58,59,59,60	0
2	SO4	D	301	5/5	0.99	0.07	31,34,38,39	0
2	SO4	A	301	5/5	0.99	0.03	43,48,50,51	0
2	SO4	H	301	5/5	0.99	0.06	40,42,47,49	0
3	EDO	A	302	4/4	0.99	0.04	28,29,34,37	0
3	EDO	D	300	4/4	0.99	0.04	27,28,32,34	0
3	EDO	D	302	4/4	0.99	0.04	34,34,41,42	0
2	SO4	C	301	5/5	0.99	0.05	41,41,48,49	0
2	SO4	B	301	5/5	1.00	0.03	27,29,31,33	0

6.5 Other polymers [i](#)

There are no such residues in this entry.