



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 8, 2026 – 05:20 PM UTC

PDB ID : 6TRH / pdb_00006trh
Title : Structure of E70A mutant of Rex8A from *Paenibacillus barcinonensis* complexed with 3(3)-alpha-L-arabinofuranosyl-xylotetraose.
Authors : Jimenez-Ortega, E.; Ramirez-Escudero, M.; Sanz-Aparicio, J.
Deposited on : 2019-12-18
Resolution : 1.86 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.86 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3428 (1.86-1.86)
Clashscore	190562	3579 (1.86-1.86)
Ramachandran outliers	187476	3553 (1.86-1.86)
Sidechain outliers	187428	3553 (1.86-1.86)
RSRZ outliers	180081	3429 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	385	92% 7%
1	B	385	91% 8%
1	C	385	91% 8%
1	D	385	93% 5%
2	E	5	100%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	F	5	 100%
2	G	5	 100%
2	H	5	 100%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 12933 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Reducing-end xylose-releasing exo-oligoxylanase Rex8A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	381	Total	C	N	O	S	0	0	0
			3128	1995	536	582	15			
1	B	380	Total	C	N	O	S	0	1	0
			3132	1998	538	581	15			
1	C	385	Total	C	N	O	S	0	2	0
			3173	2023	545	589	16			
1	D	380	Total	C	N	O	S	0	0	0
			3124	1993	535	581	15			

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	70	ALA	GLU	engineered mutation	UNP A0A0S2UQQ5
A	381	GLN	-	expression tag	UNP A0A0S2UQQ5
A	382	GLN	-	expression tag	UNP A0A0S2UQQ5
A	383	ALA	-	expression tag	UNP A0A0S2UQQ5
A	384	ALA	-	expression tag	UNP A0A0S2UQQ5
A	385	ALA	-	expression tag	UNP A0A0S2UQQ5
B	70	ALA	GLU	engineered mutation	UNP A0A0S2UQQ5
B	381	GLN	-	expression tag	UNP A0A0S2UQQ5
B	382	GLN	-	expression tag	UNP A0A0S2UQQ5
B	383	ALA	-	expression tag	UNP A0A0S2UQQ5
B	384	ALA	-	expression tag	UNP A0A0S2UQQ5
B	385	ALA	-	expression tag	UNP A0A0S2UQQ5
C	70	ALA	GLU	engineered mutation	UNP A0A0S2UQQ5
C	381	GLN	-	expression tag	UNP A0A0S2UQQ5
C	382	GLN	-	expression tag	UNP A0A0S2UQQ5
C	383	ALA	-	expression tag	UNP A0A0S2UQQ5
C	384	ALA	-	expression tag	UNP A0A0S2UQQ5
C	385	ALA	-	expression tag	UNP A0A0S2UQQ5
D	70	ALA	GLU	engineered mutation	UNP A0A0S2UQQ5
D	381	GLN	-	expression tag	UNP A0A0S2UQQ5
D	382	GLN	-	expression tag	UNP A0A0S2UQQ5

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	383	ALA	-	expression tag	UNP A0A0S2UQQ5
D	384	ALA	-	expression tag	UNP A0A0S2UQQ5
D	385	ALA	-	expression tag	UNP A0A0S2UQQ5

- Molecule 2 is an oligosaccharide called alpha-L-arabinofuranose-(1-3)-[beta-D-xylopyranose-(1-4)]beta-D-xylopyranose-(1-4)-beta-D-xylopyranose-(1-4)-beta-D-xylopyranose.



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
2	E	5	Total	C	O	0	0	0
			46	25	21			
2	F	5	Total	C	O	0	0	0
			46	25	21			
2	G	5	Total	C	O	0	0	0
			46	25	21			
2	H	5	Total	C	O	0	0	0
			46	25	21			

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	50	Total	O	0	0
			50	50		
3	B	46	Total	O	0	0
			46	46		
3	C	41	Total	O	0	0
			41	41		
3	D	55	Total	O	0	0
			55	55		

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Reducing-end xylose-releasing exo-oligoxylanase Rex8A

Chain A: 




- Molecule 1: Reducing-end xylose-releasing exo-oligoxylanase Rex8A

Chain B: 




- Molecule 1: Reducing-end xylose-releasing exo-oligoxylanase Rex8A

Chain C: 




- Molecule 1: Reducing-end xylose-releasing exo-oligoxylanase Rex8A

Chain D: 



- Molecule 2: alpha-L-arabinofuranose-(1-3)-[beta-D-xylopyranose-(1-4)]beta-D-xylopyranose-(1-4)-beta-D-xylopyranose-(1-4)-beta-D-xylopyranose

Chain E: 

XYP1
XYP2
XYP3
AHR4
XYP5

- Molecule 2: alpha-L-arabinofuranose-(1-3)-[beta-D-xylopyranose-(1-4)]beta-D-xylopyranose-(1-4)-beta-D-xylopyranose-(1-4)-beta-D-xylopyranose

Chain F:  100%

XYP1
XYP2
XYP3
AHR4
XYP5

- Molecule 2: alpha-L-arabinofuranose-(1-3)-[beta-D-xylopyranose-(1-4)]beta-D-xylopyranose-(1-4)-beta-D-xylopyranose-(1-4)-beta-D-xylopyranose

Chain G:  100%

XYP1
XYP2
XYP3
AHR4
XYP5

- Molecule 2: alpha-L-arabinofuranose-(1-3)-[beta-D-xylopyranose-(1-4)]beta-D-xylopyranose-(1-4)-beta-D-xylopyranose-(1-4)-beta-D-xylopyranose

Chain H:  100%

XYP1
XYP2
XYP3
AHR4
XYP5

4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	82.30Å 82.72Å 465.32Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.62 – 1.86 46.62 – 1.86	Depositor EDS
% Data completeness (in resolution range)	95.7 (46.62-1.86) 94.0 (46.62-1.86)	Depositor EDS
R_{merge}	0.15	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.17 (at 1.86Å)	Xtriage
Refinement program	REFMAC 5.8.0257	Depositor
R, R_{free}	0.238 , 0.263 (Not available) , 0.291	Depositor DCC
R_{free} test set	6472 reflections (4.84%)	wwPDB-VP
Wilson B-factor (Å ²)	32.4	Xtriage
Anisotropy	0.369	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 15.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.35$, $\langle L^2 \rangle = 0.17$	Xtriage
Estimated twinning fraction	0.428 for -k,-h,-l	Xtriage
Reported twinning fraction	0.520 for H, K, L 0.480 for -K, -H, -L	Depositor
Outliers	0 of 129753 reflections	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	12933	wwPDB-VP
Average B, all atoms (Å ²)	36.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.91% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: AHR, XYP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.00	0/3235	1.47	1/4394 (0.0%)
1	B	1.00	0/3242	1.47	0/4403
1	C	1.00	0/3286	1.47	1/4462 (0.0%)
1	D	1.00	0/3231	1.47	3/4389 (0.1%)
All	All	1.00	0/12994	1.47	5/17648 (0.0%)

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	173	GLY	CA-C-O	-6.44	117.67	122.37
1	A	173	GLY	CA-C-O	-6.02	117.45	122.33
1	C	173	GLY	CA-C-O	-5.87	117.58	122.33
1	D	25	GLU	N-CA-CB	5.42	117.88	110.01
1	D	24	ARG	CA-C-O	-5.20	115.49	121.47

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3128	0	2867	15	0
1	B	3132	0	2877	19	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	3173	0	2924	26	0
1	D	3124	0	2864	10	0
2	E	46	0	0	0	0
2	F	46	0	0	0	0
2	G	46	0	0	0	0
2	H	46	0	0	0	0
3	A	50	0	0	2	0
3	B	46	0	0	1	0
3	C	41	0	0	2	0
3	D	55	0	0	1	0
All	All	12933	0	11532	66	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 66 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:115:LYS:HG3	1:C:121:LEU:HD11	1.45	0.97
1:C:39:LEU:HD21	1:C:363:ASN:HB3	1.55	0.89
1:A:39:LEU:HD21	1:A:363:ASN:HB3	1.55	0.88
1:D:39:LEU:HD21	1:D:363:ASN:HB3	1.57	0.87
1:C:114:CYS:HA	1:C:121:LEU:HD13	1.57	0.86

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	379/385 (98%)	366 (97%)	13 (3%)	0	100	100
1	B	379/385 (98%)	365 (96%)	14 (4%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	C	385/385 (100%)	371 (96%)	12 (3%)	2 (0%)	24	13
1	D	378/385 (98%)	366 (97%)	12 (3%)	0	100	100
All	All	1521/1540 (99%)	1468 (96%)	51 (3%)	2 (0%)	48	35

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	3	ILE
1	C	6	LYS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	314/318 (99%)	313 (100%)	1 (0%)	86	85
1	B	315/318 (99%)	311 (99%)	4 (1%)	61	51
1	C	320/318 (101%)	319 (100%)	1 (0%)	86	85
1	D	314/318 (99%)	314 (100%)	0	100	100
All	All	1263/1272 (99%)	1257 (100%)	6 (0%)	81	77

5 of 6 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	169	GLU
1	B	299	ASP
1	C	4	THR
1	B	20	ARG
1	A	172	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 12 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	350	ASN
1	C	245	ASN
1	D	350	ASN
1	D	162	HIS
1	A	245	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

20 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	XYP	E	1	2	10,10,10	1.32	2 (20%)	14,14,14	1.46	2 (14%)
2	XYP	E	2	2	9,9,10	1.61	2 (22%)	10,12,14	2.42	4 (40%)
2	XYP	E	3	2	9,9,10	1.96	3 (33%)	10,12,14	2.02	3 (30%)
2	AHR	E	4	2	9,9,10	1.84	3 (33%)	11,12,14	1.62	3 (27%)
2	XYP	E	5	2	9,9,10	2.09	2 (22%)	10,12,14	3.74	5 (50%)
2	XYP	F	1	2	10,10,10	1.49	2 (20%)	14,14,14	1.29	2 (14%)
2	XYP	F	2	2	9,9,10	1.46	2 (22%)	10,12,14	2.50	4 (40%)
2	XYP	F	3	2	9,9,10	1.89	3 (33%)	10,12,14	2.20	3 (30%)
2	AHR	F	4	2	9,9,10	1.98	3 (33%)	11,12,14	1.35	2 (18%)
2	XYP	F	5	2	9,9,10	2.10	4 (44%)	10,12,14	3.60	6 (60%)
2	XYP	G	1	2	10,10,10	1.62	2 (20%)	14,14,14	1.25	2 (14%)
2	XYP	G	2	2	9,9,10	1.64	3 (33%)	10,12,14	2.52	4 (40%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	XYP	G	3	2	9,9,10	2.26	3 (33%)	10,12,14	2.11	3 (30%)
2	AHR	G	4	2	9,9,10	2.18	2 (22%)	11,12,14	1.44	2 (18%)
2	XYP	G	5	2	9,9,10	1.85	2 (22%)	10,12,14	3.82	5 (50%)
2	XYP	H	1	2	10,10,10	1.49	3 (30%)	14,14,14	1.26	2 (14%)
2	XYP	H	2	2	9,9,10	1.54	2 (22%)	10,12,14	2.43	3 (30%)
2	XYP	H	3	2	9,9,10	1.97	3 (33%)	10,12,14	2.23	3 (30%)
2	AHR	H	4	2	9,9,10	2.03	3 (33%)	11,12,14	1.38	1 (9%)
2	XYP	H	5	2	9,9,10	1.78	3 (33%)	10,12,14	3.93	5 (50%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	XYP	E	1	2	-	-	0/1/1/1
2	XYP	E	2	2	-	-	0/1/1/1
2	XYP	E	3	2	-	-	0/1/1/1
2	AHR	E	4	2	-	0/2/15/18	0/1/1/1
2	XYP	E	5	2	-	-	0/1/1/1
2	XYP	F	1	2	-	-	0/1/1/1
2	XYP	F	2	2	-	-	0/1/1/1
2	XYP	F	3	2	-	-	0/1/1/1
2	AHR	F	4	2	-	0/2/15/18	0/1/1/1
2	XYP	F	5	2	-	-	0/1/1/1
2	XYP	G	1	2	-	-	0/1/1/1
2	XYP	G	2	2	-	-	0/1/1/1
2	XYP	G	3	2	-	-	0/1/1/1
2	AHR	G	4	2	-	0/2/15/18	0/1/1/1
2	XYP	G	5	2	-	-	0/1/1/1
2	XYP	H	1	2	-	-	0/1/1/1
2	XYP	H	2	2	-	-	0/1/1/1
2	XYP	H	3	2	-	-	0/1/1/1
2	AHR	H	4	2	-	0/2/15/18	0/1/1/1
2	XYP	H	5	2	-	-	0/1/1/1

The worst 5 of 52 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	G	3	XYP	C2-C3	4.98	1.60	1.52

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	E	5	XYP	O5-C5	-4.73	1.34	1.43
2	G	4	AHR	C1-C2	4.44	1.59	1.51
2	H	4	AHR	C1-C2	4.43	1.59	1.51
2	G	5	XYP	O5-C5	-4.01	1.35	1.43

The worst 5 of 64 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	H	5	XYP	C5-O5-C1	-8.49	97.71	111.42
2	G	5	XYP	C5-O5-C1	-8.47	97.74	111.42
2	E	5	XYP	C5-O5-C1	-7.80	98.82	111.42
2	F	5	XYP	C5-O5-C1	-7.23	99.75	111.42
2	E	5	XYP	C1-C2-C3	-6.96	99.51	109.64


There are no chirality outliers.



There are no torsion outliers.


There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.







5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	381/385 (98%)	-0.70	0 100 100	21, 35, 52, 62	0
1	B	380/385 (98%)	-0.62	0 100 100	22, 35, 54, 61	1 (0%)
1	C	385/385 (100%)	-0.65	0 100 100	19, 37, 51, 59	2 (0%)
1	D	380/385 (98%)	-0.62	0 100 100	21, 35, 52, 62	0
All	All	1526/1540 (99%)	-0.65	0 100 100	19, 35, 52, 62	3 (0%)

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	XYP	G	5	9/10	0.94	0.09	21,21,22,22	0
2	XYP	F	3	9/10	0.95	0.05	21,21,22,22	0
2	AHR	F	4	9/10	0.95	0.09	20,21,22,22	0
2	AHR	E	4	9/10	0.95	0.07	20,21,22,22	0
2	XYP	H	3	9/10	0.95	0.06	21,21,22,22	0
2	XYP	E	3	9/10	0.96	0.05	20,21,21,22	0
2	XYP	H	2	9/10	0.96	0.06	21,21,21,21	0
2	XYP	F	1	10/10	0.96	0.05	21,21,21,22	0
2	AHR	H	4	9/10	0.96	0.06	20,21,21,22	0

Continued on next page...

Continued from previous page...




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	XYP	G	2	9/10	0.97	0.05	20,21,21,22	0
2	XYP	G	3	9/10	0.97	0.05	21,21,21,22	0
2	AHR	G	4	9/10	0.97	0.05	20,21,21,22	0
2	XYP	F	2	9/10	0.97	0.05	20,20,21,21	0
2	XYP	H	1	10/10	0.97	0.05	20,21,21,21	0
2	XYP	E	5	9/10	0.97	0.05	20,21,21,21	0
2	XYP	E	2	9/10	0.97	0.05	20,20,21,21	0
2	XYP	F	5	9/10	0.97	0.05	21,21,22,22	0
2	XYP	H	5	9/10	0.97	0.06	21,22,22,22	0
2	XYP	G	1	10/10	0.98	0.04	21,21,21,22	0
2	XYP	E	1	10/10	0.98	0.04	20,21,21,21	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.







Electron density around Chain F:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around Chain G:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.