wwPDB X-ray Structure Validation Summary Report (i)

Nov 10, 2026 – 03:24 pm GMT

PDB ID	:	9TRU / pdb_00009tru
Title	:	Zebrafish dUTPase in complex with staphylococcal Stl repressor
Authors	:	Perey-Simon, V.; Toth, Z.S.; Dombovari, D.; Leveles, I.; Vertessy, G.B.; Nyiri, K.; Nagy, G.N.
Deposited on	:	2025-12-27
Resolution	:	2.26 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:


MolProbity	:	4-5-2 with Phenix2.2
Xtriage (Phenix)	:	2.2
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.015 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.51

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.26 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1898 (2.26-2.26)
Clashscore	190562	2005 (2.26-2.26)
Ramachandran outliers	187476	1965 (2.26-2.26)
Sidechain outliers	187428	1966 (2.26-2.26)
RSRZ outliers	180081	1898 (2.26-2.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	186	3% 67% 5% 27%
1	B	186	3% 66% 5% 28%
1	C	186	2% 67% 30%
1	D	186	3% 61% 10% 29%
1	E	186	63% 7% 28%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	186	2% 66% 6% 27%
2	G	156	1% 81% 13% 6%
2	H	156	1% 78% 15% 6%
2	I	156	3% 87% 6% 7%
2	J	156	12% 77% 15% 8%
2	K	156	4% 82% 11% 7%
2	L	156	22% 73% 19% 7%

2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 13671 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Deoxyuridine 5'-triphosphate nucleotidohydrolase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	136	Total	C	N	O	S	0	0	0
			1031	659	177	192	3			
1	B	133	Total	C	N	O	S	0	0	0
			1018	651	173	191	3			
1	C	131	Total	C	N	O	S	0	0	0
			1004	642	170	189	3			
1	D	132	Total	C	N	O	S	0	0	0
			1018	651	176	188	3			
1	E	133	Total	C	N	O	S	0	0	0
			1031	657	177	194	3			
1	F	135	Total	C	N	O	S	0	0	0
			1039	662	179	195	3			

There are 132 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-20	MET	-	initiating methionine	UNP Q5XJ23
A	-19	GLY	-	expression tag	UNP Q5XJ23
A	-18	SER	-	expression tag	UNP Q5XJ23
A	-17	SER	-	expression tag	UNP Q5XJ23
A	-16	HIS	-	expression tag	UNP Q5XJ23
A	-15	HIS	-	expression tag	UNP Q5XJ23
A	-14	HIS	-	expression tag	UNP Q5XJ23
A	-13	HIS	-	expression tag	UNP Q5XJ23
A	-12	HIS	-	expression tag	UNP Q5XJ23
A	-11	HIS	-	expression tag	UNP Q5XJ23
A	-10	SER	-	expression tag	UNP Q5XJ23
A	-9	SER	-	expression tag	UNP Q5XJ23
A	-8	GLY	-	expression tag	UNP Q5XJ23
A	-7	LEU	-	expression tag	UNP Q5XJ23
A	-6	VAL	-	expression tag	UNP Q5XJ23
A	-5	PRO	-	expression tag	UNP Q5XJ23
A	-4	ARG	-	expression tag	UNP Q5XJ23

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	-3	GLY	-	expression tag	UNP Q5XJ23
A	-2	SER	-	expression tag	UNP Q5XJ23
A	-1	HIS	-	expression tag	UNP Q5XJ23
A	0	MET	-	expression tag	UNP Q5XJ23
A	1	SER	-	expression tag	UNP Q5XJ23
B	-20	MET	-	initiating methionine	UNP Q5XJ23
B	-19	GLY	-	expression tag	UNP Q5XJ23
B	-18	SER	-	expression tag	UNP Q5XJ23
B	-17	SER	-	expression tag	UNP Q5XJ23
B	-16	HIS	-	expression tag	UNP Q5XJ23
B	-15	HIS	-	expression tag	UNP Q5XJ23
B	-14	HIS	-	expression tag	UNP Q5XJ23
B	-13	HIS	-	expression tag	UNP Q5XJ23
B	-12	HIS	-	expression tag	UNP Q5XJ23
B	-11	HIS	-	expression tag	UNP Q5XJ23
B	-10	SER	-	expression tag	UNP Q5XJ23
B	-9	SER	-	expression tag	UNP Q5XJ23
B	-8	GLY	-	expression tag	UNP Q5XJ23
B	-7	LEU	-	expression tag	UNP Q5XJ23
B	-6	VAL	-	expression tag	UNP Q5XJ23
B	-5	PRO	-	expression tag	UNP Q5XJ23
B	-4	ARG	-	expression tag	UNP Q5XJ23
B	-3	GLY	-	expression tag	UNP Q5XJ23
B	-2	SER	-	expression tag	UNP Q5XJ23
B	-1	HIS	-	expression tag	UNP Q5XJ23
B	0	MET	-	expression tag	UNP Q5XJ23
B	1	SER	-	expression tag	UNP Q5XJ23
C	-20	MET	-	initiating methionine	UNP Q5XJ23
C	-19	GLY	-	expression tag	UNP Q5XJ23
C	-18	SER	-	expression tag	UNP Q5XJ23
C	-17	SER	-	expression tag	UNP Q5XJ23
C	-16	HIS	-	expression tag	UNP Q5XJ23
C	-15	HIS	-	expression tag	UNP Q5XJ23
C	-14	HIS	-	expression tag	UNP Q5XJ23
C	-13	HIS	-	expression tag	UNP Q5XJ23
C	-12	HIS	-	expression tag	UNP Q5XJ23
C	-11	HIS	-	expression tag	UNP Q5XJ23
C	-10	SER	-	expression tag	UNP Q5XJ23
C	-9	SER	-	expression tag	UNP Q5XJ23
C	-8	GLY	-	expression tag	UNP Q5XJ23
C	-7	LEU	-	expression tag	UNP Q5XJ23
C	-6	VAL	-	expression tag	UNP Q5XJ23

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	-5	PRO	-	expression tag	UNP Q5XJ23
C	-4	ARG	-	expression tag	UNP Q5XJ23
C	-3	GLY	-	expression tag	UNP Q5XJ23
C	-2	SER	-	expression tag	UNP Q5XJ23
C	-1	HIS	-	expression tag	UNP Q5XJ23
C	0	MET	-	expression tag	UNP Q5XJ23
C	1	SER	-	expression tag	UNP Q5XJ23
D	-20	MET	-	initiating methionine	UNP Q5XJ23
D	-19	GLY	-	expression tag	UNP Q5XJ23
D	-18	SER	-	expression tag	UNP Q5XJ23
D	-17	SER	-	expression tag	UNP Q5XJ23
D	-16	HIS	-	expression tag	UNP Q5XJ23
D	-15	HIS	-	expression tag	UNP Q5XJ23
D	-14	HIS	-	expression tag	UNP Q5XJ23
D	-13	HIS	-	expression tag	UNP Q5XJ23
D	-12	HIS	-	expression tag	UNP Q5XJ23
D	-11	HIS	-	expression tag	UNP Q5XJ23
D	-10	SER	-	expression tag	UNP Q5XJ23
D	-9	SER	-	expression tag	UNP Q5XJ23
D	-8	GLY	-	expression tag	UNP Q5XJ23
D	-7	LEU	-	expression tag	UNP Q5XJ23
D	-6	VAL	-	expression tag	UNP Q5XJ23
D	-5	PRO	-	expression tag	UNP Q5XJ23
D	-4	ARG	-	expression tag	UNP Q5XJ23
D	-3	GLY	-	expression tag	UNP Q5XJ23
D	-2	SER	-	expression tag	UNP Q5XJ23
D	-1	HIS	-	expression tag	UNP Q5XJ23
D	0	MET	-	expression tag	UNP Q5XJ23
D	1	SER	-	expression tag	UNP Q5XJ23
E	-20	MET	-	initiating methionine	UNP Q5XJ23
E	-19	GLY	-	expression tag	UNP Q5XJ23
E	-18	SER	-	expression tag	UNP Q5XJ23
E	-17	SER	-	expression tag	UNP Q5XJ23
E	-16	HIS	-	expression tag	UNP Q5XJ23
E	-15	HIS	-	expression tag	UNP Q5XJ23
E	-14	HIS	-	expression tag	UNP Q5XJ23
E	-13	HIS	-	expression tag	UNP Q5XJ23
E	-12	HIS	-	expression tag	UNP Q5XJ23
E	-11	HIS	-	expression tag	UNP Q5XJ23
E	-10	SER	-	expression tag	UNP Q5XJ23
E	-9	SER	-	expression tag	UNP Q5XJ23
E	-8	GLY	-	expression tag	UNP Q5XJ23

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
E	-7	LEU	-	expression tag	UNP Q5XJ23
E	-6	VAL	-	expression tag	UNP Q5XJ23
E	-5	PRO	-	expression tag	UNP Q5XJ23
E	-4	ARG	-	expression tag	UNP Q5XJ23
E	-3	GLY	-	expression tag	UNP Q5XJ23
E	-2	SER	-	expression tag	UNP Q5XJ23
E	-1	HIS	-	expression tag	UNP Q5XJ23
E	0	MET	-	expression tag	UNP Q5XJ23
E	1	SER	-	expression tag	UNP Q5XJ23
F	-20	MET	-	initiating methionine	UNP Q5XJ23
F	-19	GLY	-	expression tag	UNP Q5XJ23
F	-18	SER	-	expression tag	UNP Q5XJ23
F	-17	SER	-	expression tag	UNP Q5XJ23
F	-16	HIS	-	expression tag	UNP Q5XJ23
F	-15	HIS	-	expression tag	UNP Q5XJ23
F	-14	HIS	-	expression tag	UNP Q5XJ23
F	-13	HIS	-	expression tag	UNP Q5XJ23
F	-12	HIS	-	expression tag	UNP Q5XJ23
F	-11	HIS	-	expression tag	UNP Q5XJ23
F	-10	SER	-	expression tag	UNP Q5XJ23
F	-9	SER	-	expression tag	UNP Q5XJ23
F	-8	GLY	-	expression tag	UNP Q5XJ23
F	-7	LEU	-	expression tag	UNP Q5XJ23
F	-6	VAL	-	expression tag	UNP Q5XJ23
F	-5	PRO	-	expression tag	UNP Q5XJ23
F	-4	ARG	-	expression tag	UNP Q5XJ23
F	-3	GLY	-	expression tag	UNP Q5XJ23
F	-2	SER	-	expression tag	UNP Q5XJ23
F	-1	HIS	-	expression tag	UNP Q5XJ23
F	0	MET	-	expression tag	UNP Q5XJ23
F	1	SER	-	expression tag	UNP Q5XJ23

- Molecule 2 is a protein called Orf20.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	G	146	Total	C	N	O	S	0	0	0
			1202	772	192	236	2			
2	I	145	Total	C	N	O	S	0	0	0
			1194	768	192	232	2			
2	J	144	Total	C	N	O	S	0	0	0
			1162	747	182	231	2			
2	H	146	Total	C	N	O	S	0	0	0
			1199	771	190	236	2			

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	L	145	Total	C	N	O	S	0	0	0
			1163	750	183	228	2			
2	K	145	Total	C	N	O	S	0	0	0
			1175	756	185	232	2			

- Molecule 3 is MAGNESIUM ION (CCD ID: MG) (formula: Mg) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Mg	0	0
			1	1		
3	D	1	Total	Mg	0	0
			1	1		

- Molecule 4 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	40	Total	O	0	0
			40	40		
4	B	57	Total	O	0	0
			57	57		
4	C	32	Total	O	0	0
			32	32		
4	D	41	Total	O	0	0
			41	41		
4	E	51	Total	O	0	0
			51	51		
4	F	38	Total	O	0	0
			38	38		
4	G	53	Total	O	0	0
			53	53		
4	I	32	Total	O	0	0
			32	32		
4	J	14	Total	O	0	0
			14	14		
4	H	47	Total	O	0	0
			47	47		
4	L	10	Total	O	0	0
			10	10		
4	K	18	Total	O	0	0
			18	18		

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Deoxyuridine 5'-triphosphate nucleotidohydrolase

Chain A:




• Molecule 1: Deoxyuridine 5'-triphosphate nucleotidohydrolase

Chain B:



• Molecule 1: Deoxyuridine 5'-triphosphate nucleotidohydrolase


Chain C:



• Molecule 1: Deoxyuridine 5'-triphosphate nucleotidohydrolase


Chain D:






• Molecule 1: Deoxyuridine 5'-triphosphate nucleotidohydrolase







• Molecule 1: Deoxyuridine 5'-triphosphate nucleotidohydrolase







• Molecule 2: Orf20







• Molecule 2: Orf20






• Molecule 2: Orf20








• Molecule 2: Orf20

Chain H:







• Molecule 2: Orf20

Chain L:








• Molecule 2: Orf20

Chain K:





4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α, β, γ	59.51Å 128.01Å 127.07Å 90.00° 93.72° 90.00°	Depositor
Resolution (Å)	59.38 – 2.26 59.38 – 2.26	Depositor EDS
% Data completeness (in resolution range)	99.0 (59.38-2.26) 99.1 (59.38-2.26)	Depositor EDS
R_{merge}	0.14	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	1.18 (at 2.27Å)	Xtriage
Refinement program	PHENIX 1.20.1_4487	Depositor
R, R_{free}	0.224 , 0.265 0.227 , 0.264	Depositor DCC
R_{free} test set	1927 reflections (2.17%)	wwPDB-VP
Wilson B-factor (Å^2)	45.4	Xtriage
Anisotropy	0.201	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.35 , 52.9	EDS
L-test for twinning[2]	$< \|L\| > = 0.48$, $< L^2 > = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o,F_c correlation	0.94	EDS
Total number of atoms	13671	wwPDB-VP
Average B, all atoms (Å^2)	57.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 14.72% of the height of the origin peak. No significant pseudotranslation is detected.*

[1] Intensities estimated from amplitudes.

[2] Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#$\|Z\|$ >5	RMSZ	#$\|Z\|$ >5
1	A	0.20	0/1053	0.41	0/1426
1	B	0.22	0/1039	0.43	0/1405
1	C	0.20	0/1025	0.42	0/1387
1	D	0.20	0/1039	0.41	0/1404
1	E	0.23	0/1052	0.45	0/1421
1	F	0.21	0/1060	0.45	0/1432
2	G	0.21	0/1227	0.40	0/1655
2	H	0.21	0/1224	0.40	0/1650
2	I	0.18	0/1219	0.36	0/1643
2	J	0.15	0/1187	0.33	0/1605
2	K	0.16	0/1200	0.34	0/1622
2	L	0.14	0/1188	0.37	0/1606
All	All	0.19	0/13513	0.40	0/18256

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1031	0	1010	6	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	1018	0	1000	7	0
1	C	1004	0	979	5	0
1	D	1018	0	1008	14	0
1	E	1031	0	1019	9	0
1	F	1039	0	1021	11	0
2	G	1202	0	1175	10	0
2	H	1199	0	1176	20	0
2	I	1194	0	1179	6	0
2	J	1162	0	1114	16	0
2	K	1175	0	1130	10	0
2	L	1163	0	1115	19	0
3	A	1	0	0	0	0
3	D	1	0	0	0	0
4	A	40	0	0	1	0
4	B	57	0	0	1	0
4	C	32	0	0	0	0
4	D	41	0	0	0	0
4	E	51	0	0	1	0
4	F	38	0	0	2	0
4	G	53	0	0	0	0
4	H	47	0	0	1	0
4	I	32	0	0	0	0
4	J	14	0	0	0	0
4	K	18	0	0	1	0
4	L	10	0	0	0	0
All	All	13671	0	12926	119	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 119 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:F:24:ARG:HD2	1:F:25:ALA:H	1.33	0.92
2:H:96:LYS:NZ	2:H:137:ASN:OD1	2.14	0.79
1:D:92:LYS:HE3	2:I:114:SER:HB3	1.67	0.76
2:H:96:LYS:HE2	2:H:136:VAL:HG22	1.71	0.73
2:H:8:ALA:HB1	2:H:55:VAL:HG22	1.72	0.72

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	134/186 (72%)	134 (100%)	0	0	100	100
1	B	129/186 (69%)	129 (100%)	0	0	100	100
1	C	127/186 (68%)	125 (98%)	2 (2%)	0	100	100
1	D	128/186 (69%)	127 (99%)	1 (1%)	0	100	100
1	E	129/186 (69%)	128 (99%)	1 (1%)	0	100	100
1	F	131/186 (70%)	129 (98%)	2 (2%)	0	100	100
2	G	144/156 (92%)	143 (99%)	1 (1%)	0	100	100
2	H	144/156 (92%)	142 (99%)	2 (1%)	0	100	100
2	I	143/156 (92%)	142 (99%)	1 (1%)	0	100	100
2	J	142/156 (91%)	142 (100%)	0	0	100	100
2	K	143/156 (92%)	143 (100%)	0	0	100	100
2	L	143/156 (92%)	141 (99%)	2 (1%)	0	100	100
All	All	1637/2052 (80%)	1625 (99%)	12 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	104/149 (70%)	100 (96%)	4 (4%)	29	36
1	B	105/149 (70%)	104 (99%)	1 (1%)	68	77

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	C	103/149 (69%)	102 (99%)	1 (1%)	68 77
1	D	105/149 (70%)	105 (100%)	0	100 100
1	E	108/149 (72%)	103 (95%)	5 (5%)	24 28
1	F	107/149 (72%)	103 (96%)	4 (4%)	30 37
2	G	133/141 (94%)	130 (98%)	3 (2%)	44 55
2	H	133/141 (94%)	131 (98%)	2 (2%)	57 68
2	I	133/141 (94%)	132 (99%)	1 (1%)	73 80
2	J	127/141 (90%)	125 (98%)	2 (2%)	55 66
2	K	128/141 (91%)	127 (99%)	1 (1%)	73 80
2	L	125/141 (89%)	122 (98%)	3 (2%)	43 54
All	All	1411/1740 (81%)	1384 (98%)	27 (2%)	50 61

5 of 27 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	F	152	THR
2	G	154	THR
2	L	25	MET
2	G	151	ILE
2	I	55	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 10 such sidechains are listed below:

Mol	Chain	Res	Type
2	J	50	ASN
2	J	107	ASN
2	L	48	ASN
2	G	102	ASN
2	G	137	ASN

5.3.3 RNA

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

5.6 Ligand geometry (i)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	136/186 (73%)	0.53	5 (3%)	45	45	38, 51, 72, 107	0
1	B	133/186 (71%)	0.32	6 (4%)	38	37	37, 43, 69, 101	0
1	C	131/186 (70%)	0.39	4 (3%)	51	51	37, 50, 74, 94	0
1	D	132/186 (70%)	0.39	5 (3%)	44	44	36, 46, 68, 78	0
1	E	133/186 (71%)	0.31	1 (0%)	82	84	33, 42, 70, 96	0
1	F	135/186 (72%)	0.40	3 (2%)	62	63	37, 52, 74, 83	0
2	G	146/156 (93%)	0.32	2 (1%)	73	74	33, 45, 63, 99	0
2	H	146/156 (93%)	0.43	1 (0%)	84	85	38, 48, 80, 97	0
2	I	145/156 (92%)	0.40	5 (3%)	48	48	38, 52, 82, 105	0
2	J	144/156 (92%)	0.99	18 (12%)	8	7	39, 76, 121, 149	0
2	K	145/156 (92%)	0.76	6 (4%)	41	41	49, 63, 84, 108	0
2	L	145/156 (92%)	1.37	34 (23%)	2	1	48, 97, 131, 147	0
All	All	1671/2052 (81%)	0.56	90 (5%)	31	29	33, 52, 103, 149	0

The worst 5 of 90 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	152	THR	5.0
2	J	32	LEU	4.1
2	G	154	THR	3.7
2	I	136	VAL	3.7
2	J	83	GLY	3.7

6.2 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no oligosaccharides in this entry.

6.4 Ligands (i)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
3	MG	D	201	1/1	0.65	0.39	65,65,65,65	0
3	MG	A	201	1/1	0.66	0.33	67,67,67,67	0


The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around MG D 201:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)








Electron density around MG A 201:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)



6.5 Other polymers (i)

There are no such residues in this entry.