



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 09:13 AM UTC

PDB ID : 4TS0 / pdb_00004ts0
Title : Crystal structure of the Spinach RNA aptamer in complex with DFHBI, barium ions
Authors : Warner, K.D.; Chen, M.C.; Song, W.; Strack, R.L.; Thorn, A.; Jaffrey, S.R.; Ferre-D'Amare, A.R.
Deposited on : 2014-06-18
Resolution : 2.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3866 (2.80-2.80)
Clashscore	190562	4276 (2.80-2.80)
RSRZ outliers	180081	3869 (2.80-2.80)
RNA backbone	3983	1114 (3.00-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	X	44	 57% 32% 5% 7%
2	Y	47	 57% 30% 13%
3	A	2	 100%
3	B	2	 100%
3	C	2	 100%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	D	2	50% 50%

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 2068 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a RNA chain called Spinach RNA aptamer, bimolecular construct.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	X	44	Total	C	N	O	P	0	0	0
			935	415	176	300	44			

- Molecule 2 is a RNA chain called Spinach RNA aptamer, bimolecular construct.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	Y	47	Total	C	N	O	P	0	0	0
			997	443	175	332	47			

- Molecule 3 is an oligosaccharide called beta-D-fructofuranose-(2-1)-alpha-D-glucopyranose.



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
3	A	2	Total	C	O	0	0	0
			23	12	11			
3	B	2	Total	C	O	0	0	0
			23	12	11			
3	C	2	Total	C	O	0	0	0
			23	12	11			
3	D	2	Total	C	O	0	0	0
			23	12	11			


- Molecule 4 is POTASSIUM ION (CCD ID: K) (formula: K).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	X	5	Total	K	0	0
			5	5		
4	Y	5	Total	K	0	0
			5	5		

- Molecule 5 is BARIUM ION (CCD ID: BA) (formula: Ba).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	X	1	Total	Ba	0	0
			1	1		

- Molecule 6 is (5Z)-5-(3,5-difluoro-4-hydroxybenzylidene)-2,3-dimethyl-3,5-dihydro-4H-imidazol-4-one (CCD ID: 38E) (formula: C₁₂H₁₀F₂N₂O₂).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
6	Y	1	Total	C	F	N	O	0	0
			18	12	2	2	2		

- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	X	7	Total	O	0	0
			7	7		
7	Y	8	Total	O	0	0
			8	8		

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Spinach RNA aptamer, bimolecular construct

Chain X: 



- Molecule 2: Spinach RNA aptamer, bimolecular construct

Chain Y: 



- Molecule 3: beta-D-fructofuranose-(2-1)-alpha-D-glucopyranose

Chain A: 




- Molecule 3: beta-D-fructofuranose-(2-1)-alpha-D-glucopyranose

Chain B: 



- Molecule 3: beta-D-fructofuranose-(2-1)-alpha-D-glucopyranose

Chain C: 



- Molecule 3: beta-D-fructofuranose-(2-1)-alpha-D-glucopyranose

Chain D: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	39.76Å 49.50Å 186.26Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 2.80 50.00 – 2.80	Depositor EDS
% Data completeness (in resolution range)	99.9 (50.00-2.80) 99.9 (50.00-2.80)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.03 (at 2.51Å)	Xtriage
Refinement program	REFMAC 5.8.0069	Depositor
R, R_{free}	0.206 , 0.239 0.209 , 0.247	Depositor DCC
R_{free} test set	654 reflections (4.87%)	wwPDB-VP
Wilson B-factor (Å ²)	83.6	Xtriage
Anisotropy	0.595	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 63.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	2068	wwPDB-VP
Average B, all atoms (Å ²)	91.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.45% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: 38E, FRU, GLC, CCC, BA, K

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	X	0.53	0/1022	1.08	6/1595 (0.4%)
2	Y	0.56	0/1088	1.02	4/1696 (0.2%)
All	All	0.54	0/2110	1.05	10/3291 (0.3%)

There are no bond length outliers.

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	X	22	G	C1'-C2'-O2'	7.99	120.38	108.40
2	Y	63	G	C2'-C3'-O3'	-5.98	104.73	113.70
1	X	27	A	P-O5'-C5'	-5.94	112.00	120.90
1	X	27	A	C1'-C2'-O2'	5.88	120.62	111.80
1	X	30	G	C3'-C2'-O2'	5.70	119.25	110.70
1	X	19	G	O4'-C1'-C2'	-5.58	102.02	107.60
1	X	12	A	C3'-C2'-O2'	-5.41	106.48	114.60
2	Y	63	G	C3'-C2'-O2'	-5.32	102.72	110.70
2	Y	73	U	C3'-C2'-O2'	5.30	118.65	110.70
2	Y	68	G	C4'-C3'-O3'	-5.10	101.75	109.40

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	X	935	0	469	20	0
2	Y	997	0	499	15	0
3	A	23	0	21	1	0
3	B	23	0	21	6	0
3	C	23	0	21	0	0
3	D	23	0	21	2	0
4	X	5	0	0	0	0
4	Y	5	0	0	0	0
5	X	1	0	0	0	0
6	Y	18	0	9	1	0
7	X	7	0	0	0	0
7	Y	8	0	0	0	0
All	All	2068	0	1061	37	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

All (37) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:X:19:G:N3	3:B:1:GLC:O6	2.12	0.82
2:Y:49:G:H5''	2:Y:49:G:H8	1.50	0.77
1:X:19:G:N2	3:B:1:GLC:O6	2.31	0.63
1:X:33:C:O2'	3:D:1:GLC:H2	1.99	0.62
2:Y:94:U:H6	2:Y:94:U:C5'	2.13	0.62
1:X:42:U:C4	1:X:43:G:N7	2.70	0.60
2:Y:81:G:O2'	3:B:2:FRU:H4	2.02	0.58
2:Y:93:G:C2'	2:Y:94:U:H5'	2.34	0.57
1:X:19:G:H21	3:B:1:GLC:C6	2.18	0.57
1:X:19:G:C2	3:B:1:GLC:O6	2.55	0.54
1:X:9:C:O2'	1:X:12:A:OP1	2.20	0.54
1:X:43:G:H5''	1:X:44:CCC:P	2.48	0.54
1:X:19:G:H8	1:X:19:G:H5''	1.73	0.54
1:X:43:G:H5''	1:X:44:CCC:OP2	2.07	0.54
2:Y:62:U:H2'	2:Y:63:G:OP1	2.09	0.53
2:Y:49:G:H5''	2:Y:49:G:C8	2.37	0.53
2:Y:65:G:H3'	2:Y:65:G:N3	2.25	0.51
6:Y:206:38E:H2	6:Y:206:38E:N	2.27	0.50
2:Y:93:G:H2'	2:Y:94:U:H5'	1.95	0.48
1:X:1:G:H2'	1:X:2:A:C8	2.49	0.48
1:X:42:U:N3	1:X:43:G:C8	2.84	0.46
1:X:19:G:H5''	1:X:19:G:C8	2.50	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:X:36:G:H2'	1:X:37:C:O4'	2.16	0.46
2:Y:78:U:H2'	2:Y:79:C:C6	2.50	0.46
1:X:25:G:H1'	1:X:26:G:C8	2.52	0.45
2:Y:94:U:C6	2:Y:94:U:C3'	2.99	0.45
3:A:1:GLC:O5	3:A:2:FRU:O6	2.29	0.45
2:Y:62:U:C2'	2:Y:63:G:OP1	2.67	0.43
2:Y:94:U:C6	2:Y:94:U:H3'	2.53	0.43
2:Y:82:U:H5''	2:Y:83:A:OP1	2.18	0.42
1:X:43:G:O3'	1:X:44:CCC:O4'	2.38	0.41
2:Y:94:U:H6	2:Y:94:U:O5'	2.01	0.41
1:X:33:C:O2'	3:D:1:GLC:C2	2.67	0.41
1:X:19:G:N3	3:B:1:GLC:C6	2.84	0.41
1:X:42:U:C2	1:X:43:G:C8	3.07	0.41
2:Y:94:U:C6	2:Y:94:U:C4'	3.04	0.41
1:X:10:G:H5''	1:X:12:A:OP1	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

There are no protein molecules in this entry.

5.3.2 Protein sidechains [i](#)

There are no protein molecules in this entry.

5.3.3 RNA [i](#)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
1	X	43/44 (97%)	7 (16%)	1 (2%)
2	Y	46/47 (97%)	10 (21%)	2 (4%)
All	All	89/91 (97%)	17 (19%)	3 (3%)

All (17) RNA backbone outliers are listed below:

Mol	Chain	Res	Type
1	X	12	A

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	X	13	U
1	X	14	G
1	X	19	G
1	X	27	A
1	X	43	G
1	X	44	CCC
2	Y	58	U
2	Y	59	U
2	Y	61	U
2	Y	62	U
2	Y	66	U
2	Y	67	A
2	Y	72	G
2	Y	73	U
2	Y	83	A
2	Y	94	U

All (3) RNA pucker outliers are listed below:

Mol	Chain	Res	Type
1	X	12	A
2	Y	61	U
2	Y	68	G

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	CCC	X	44	1,2	20,25,26	0.92	1 (5%)	27,38,41	1.76	6 (22%)
2	CCC	Y	101	1,2	20,25,26	0.95	1 (5%)	27,38,41	1.90	7 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CCC	X	44	1,2	-	3/7/35/36	0/3/3/3
2	CCC	Y	101	1,2	-	2/7/35/36	0/3/3/3

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	Y	101	CCC	C6-C5	2.12	1.40	1.35
1	X	44	CCC	C6-C5	2.07	1.39	1.35

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	Y	101	CCC	C2'-C1'-N1	-5.70	103.42	114.24
2	Y	101	CCC	O2C-PC-O1C	4.10	123.14	109.89
1	X	44	CCC	O2'-PC-O1C	-3.97	105.29	115.76
1	X	44	CCC	O2C-PC-O1C	3.82	122.23	109.89
1	X	44	CCC	C2'-C1'-N1	-3.54	107.52	114.24
2	Y	101	CCC	O4'-C1'-N1	2.94	115.03	108.36
1	X	44	CCC	O4'-C1'-N1	2.92	114.97	108.36
2	Y	101	CCC	O2'-PC-O1C	-2.91	108.07	115.76
1	X	44	CCC	O3'-PC-O1C	-2.41	109.40	115.76
2	Y	101	CCC	O2-C2-N3	-2.19	118.88	122.33
2	Y	101	CCC	C5'-C4'-C3'	-2.18	107.11	114.38
2	Y	101	CCC	O4'-C4'-C3'	2.08	109.31	104.92
1	X	44	CCC	O2-C2-N3	-2.07	119.06	122.33

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	Y	101	CCC	O4'-C4'-C5'-O5'
1	X	44	CCC	C3'-C4'-C5'-O5'
2	Y	101	CCC	C3'-C4'-C5'-O5'
1	X	44	CCC	C4'-C5'-O5'-P
1	X	44	CCC	O4'-C4'-C5'-O5'

There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	X	44	CCC	3	0

5.5 Carbohydrates [i](#)

8 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	GLC	A	1	3	11,11,12	0.64	0	15,15,17	1.37	2 (13%)
3	FRU	A	2	3	11,12,12	0.72	0	10,18,18	1.13	1 (10%)
3	GLC	B	1	3	11,11,12	0.65	0	15,15,17	1.49	3 (20%)
3	FRU	B	2	3	11,12,12	1.02	1 (9%)	10,18,18	1.40	3 (30%)
3	GLC	C	1	3	11,11,12	0.50	0	15,15,17	1.06	1 (6%)
3	FRU	C	2	3	11,12,12	0.87	0	10,18,18	1.18	1 (10%)
3	GLC	D	1	3	11,11,12	0.98	1 (9%)	15,15,17	2.60	5 (33%)
3	FRU	D	2	4,3	11,12,12	0.73	0	10,18,18	1.92	2 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GLC	A	1	3	-	1/2/19/22	0/1/1/1
3	FRU	A	2	3	-	5/5/24/24	0/1/1/1
3	GLC	B	1	3	-	2/2/19/22	0/1/1/1
3	FRU	B	2	3	-	3/5/24/24	0/1/1/1
3	GLC	C	1	3	-	2/2/19/22	0/1/1/1
3	FRU	C	2	3	-	3/5/24/24	0/1/1/1
3	GLC	D	1	3	-	2/2/19/22	0/1/1/1
3	FRU	D	2	4,3	-	3/5/24/24	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	2	FRU	O2-C2	2.61	1.45	1.40
3	D	1	GLC	O5-C5	2.03	1.47	1.43

All (18) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	D	1	GLC	O5-C5-C6	5.84	119.03	107.66
3	D	1	GLC	C1-O5-C5	-5.43	104.91	112.19
3	D	2	FRU	O1-C1-C2	-4.59	101.53	111.67
3	D	1	GLC	C3-C4-C5	-3.88	103.20	110.23
3	B	1	GLC	C3-C4-C5	-3.04	104.73	110.23
3	D	2	FRU	O3-C3-C4	-2.96	102.79	113.25
3	A	1	GLC	C1-O5-C5	2.94	116.13	112.19
3	D	1	GLC	O3-C3-C4	2.55	116.39	110.38
3	D	1	GLC	C2-C3-C4	-2.46	106.53	110.86
3	A	2	FRU	O2-C2-O5	2.44	114.02	109.33
3	C	1	GLC	O5-C5-C6	2.39	112.32	107.66
3	B	1	GLC	O4-C4-C3	2.20	115.56	110.38
3	B	1	GLC	C2-C3-C4	-2.19	107.01	110.86
3	C	2	FRU	O2-C2-O5	-2.19	105.13	109.33
3	B	2	FRU	O5-C5-C4	-2.17	99.95	105.42
3	B	2	FRU	O2-C2-O5	2.13	113.42	109.33
3	A	1	GLC	C3-C4-C5	2.11	114.06	110.23
3	B	2	FRU	O5-C5-C6	2.04	114.45	108.89

There are no chirality outliers.

All (21) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	2	FRU	O1-C1-C2-O2
3	B	2	FRU	O1-C1-C2-C3
3	B	2	FRU	O1-C1-C2-O2
3	B	2	FRU	O1-C1-C2-O5
3	C	2	FRU	O1-C1-C2-O2
3	D	2	FRU	O1-C1-C2-C3
3	D	2	FRU	O1-C1-C2-O2
3	D	2	FRU	O1-C1-C2-O5
3	A	2	FRU	C4-C5-C6-O6
3	A	2	FRU	O5-C5-C6-O6
3	D	1	GLC	O5-C5-C6-O6
3	C	1	GLC	C4-C5-C6-O6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
3	A	2	FRU	O1-C1-C2-O5
3	D	1	GLC	C4-C5-C6-O6
3	C	1	GLC	O5-C5-C6-O6
3	B	1	GLC	C4-C5-C6-O6
3	B	1	GLC	O5-C5-C6-O6
3	A	1	GLC	O5-C5-C6-O6
3	C	2	FRU	C4-C5-C6-O6
3	C	2	FRU	O1-C1-C2-O5
3	A	2	FRU	O1-C1-C2-C3


There are no ring outliers.

5 monomers are involved in 9 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	2	FRU	1	0
3	B	2	FRU	1	0
3	A	1	GLC	1	0
3	D	1	GLC	2	0
3	B	1	GLC	5	0



The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.

Oligosaccharide Chain A



Oligosaccharide Chain B





5.6 Ligand geometry

Of 12 ligands modelled in this entry, 11 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
6	38E	Y	206	-	19,19,19	4.29	8 (42%)	28,28,28	3.98	12 (42%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	38E	Y	206	-	-	0/4/20/20	0/2/2/2

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	Y	206	38E	C6-C7	13.43	1.48	1.35
6	Y	206	38E	C1-C	7.62	1.48	1.39
6	Y	206	38E	C1-C2	6.28	1.46	1.39
6	Y	206	38E	C8-N	4.76	1.38	1.32
6	Y	206	38E	O-C9	4.07	1.31	1.23
6	Y	206	38E	C9-N1	-3.08	1.32	1.39
6	Y	206	38E	C4-C6	2.51	1.51	1.46
6	Y	206	38E	O1-C1	-2.15	1.32	1.36

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	Y	206	38E	C7-C9-N1	9.57	107.85	103.36
6	Y	206	38E	C5-C-C1	-9.03	116.72	123.79
6	Y	206	38E	F-C-C1	8.01	122.91	117.13
6	Y	206	38E	C3-C2-C1	-7.96	117.56	123.79
6	Y	206	38E	O-C9-C7	-6.36	126.96	131.02
6	Y	206	38E	C4-C6-C7	-4.22	124.85	129.87

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	Y	206	38E	C6-C7-C9	4.05	127.27	122.36
6	Y	206	38E	C9-C7-N	-3.53	106.42	108.95
6	Y	206	38E	C4-C5-C	3.42	121.93	119.27
6	Y	206	38E	C-C1-C2	3.20	121.27	116.82
6	Y	206	38E	F1-C2-C3	2.62	123.88	118.64
6	Y	206	38E	C5-C4-C3	2.11	121.99	119.11

There are no chirality outliers.


There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	Y	206	38E	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	X	43/44 (97%)	-0.27	0 100 100	56, 81, 133, 150	0
2	Y	46/47 (97%)	-0.29	0 100 100	52, 78, 140, 191	0
All	All	89/91 (97%)	-0.28	0 100 100	52, 79, 140, 191	0

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	CCC	X	44	23/24	0.83	0.12	112,143,160,165	0
2	CCC	Y	101	23/24	0.85	0.09	103,120,147,156	0

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	GLC	A	1	11/12	-	-	100,120,142,142	0
3	FRU	A	2	12/12	-	-	97,117,128,137	0
3	GLC	B	1	11/12	-	-	87,113,132,136	0
3	FRU	B	2	12/12	-	-	91,139,150,160	0
3	GLC	C	1	11/12	0.71	0.09	91,99,104,106	0

Continued on next page...

Continued from previous page...




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	FRU	D	2	12/12	0.75	0.14	72,110,128,144	0
3	FRU	C	2	12/12	0.81	0.14	81,93,98,99	0
3	GLC	D	1	11/12	0.85	0.15	76,89,121,149	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around Chain B:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around Chain C:

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)






6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	K	Y	205	1/1	0.74	0.18	122,122,122,122	0
4	K	X	102	1/1	0.85	0.13	104,104,104,104	0
4	K	X	104	1/1	0.90	0.08	91,91,91,91	0
4	K	Y	203	1/1	0.92	0.14	84,84,84,84	0
4	K	X	103	1/1	0.93	0.09	97,97,97,97	0
4	K	X	105	1/1	0.95	0.09	89,89,89,89	0
4	K	Y	204	1/1	0.96	0.08	71,71,71,71	0
4	K	Y	202	1/1	0.97	0.05	88,88,88,88	0
4	K	Y	201	1/1	0.97	0.06	65,65,65,65	0
5	BA	X	106	1/1	0.97	0.06	140,140,140,140	0
4	K	X	101	1/1	0.98	0.05	66,66,66,66	0
6	38E	Y	206	18/18	0.98	0.07	51,57,68,74	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.