



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 05:52 PM UTC

PDB ID : 8TSC / pdb_00008tsc
Title : Human PI3K p85alpha/p110alpha H1047R bound to compound 3
Authors : Holliday, M.; Tang, Y.; Bulku, A.; Wilbur, J.; Fraser, J.
Deposited on : 2023-08-11
Resolution : 3.62 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.62 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1062 (3.72-3.52)
Clashscore	190562	1092 (3.72-3.52)
Ramachandran outliers	187476	1057 (3.72-3.52)
Sidechain outliers	187428	1055 (3.72-3.52)
RSRZ outliers	180081	1060 (3.72-3.52)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	1060	
2	B	300	

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 21128 atoms, of which 10567 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Phosphatidylinositol 4,5-bisphosphate 3-kinase catalytic subunit alpha isoform.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	1008	Total	C	H	N	O	S	0	0	0
			16502	5260	8275	1411	1488	68			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-9	GLY	-	expression tag	UNP P42336
A	-8	SER	-	expression tag	UNP P42336
A	-7	PRO	-	expression tag	UNP P42336
A	-6	GLY	-	expression tag	UNP P42336
A	-5	ILE	-	expression tag	UNP P42336
A	-4	SER	-	expression tag	UNP P42336
A	-3	GLY	-	expression tag	UNP P42336
A	-2	GLY	-	expression tag	UNP P42336
A	-1	GLY	-	expression tag	UNP P42336
A	0	GLY	-	expression tag	UNP P42336
A	1	GLY	-	expression tag	UNP P42336
A	1047	ARG	HIS	engineered mutation	UNP P42336


- Molecule 2 is a protein called Phosphatidylinositol 3-kinase regulatory subunit alpha.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	B	270	Total	C	H	N	O	S	0	0	0
			4572	1441	2273	409	442	7			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	316	GLY	-	expression tag	UNP P27986
B	317	PRO	-	expression tag	UNP P27986

- Molecule 3 is (1S)-7-[3-fluoro-5-(trifluoromethyl)benzamido]-N-methyl-1-(2-methylphenyl)-3-oxo-2,3-dihydro-1H-isindole-5-carboxamide (CCD ID: UJC) (formula: $C_{25}H_{19}F_4N_3O_3$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	F	H	N	O	
3	A	1	54	25	4	19	3	3	0

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	85.56Å 120.28Å 190.44Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	69.72 – 3.62 69.72 – 3.62	Depositor EDS
% Data completeness (in resolution range)	99.7 (69.72-3.62) 99.7 (69.72-3.62)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.73 (at 3.58Å)	Xtriage
Refinement program	PHENIX 1.20.1_4487	Depositor
R, R_{free}	0.216 , 0.255 0.218 , 0.254	Depositor DCC
R_{free} test set	1122 reflections (4.86%)	wwPDB-VP
Wilson B-factor (Å ²)	131.9	Xtriage
Anisotropy	0.167	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 81.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	21128	wwPDB-VP
Average B, all atoms (Å ²)	160.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.55% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: UJC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.11	0/8408	0.30	0/11358
2	B	0.12	0/2338	0.34	0/3130
All	All	0.11	0/10746	0.31	0/14488

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	8227	8275	8269	29	0
2	B	2299	2273	2271	10	0
3	A	35	19	0	0	0
All	All	10561	10567	10540	37	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (37) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:992:ARG:NH1	1:A:1027:ALA:O	2.28	0.66
1:A:531:LEU:HA	1:A:534:ILE:HD12	1.82	0.61
1:A:675:MET:HE1	1:A:685:GLY:N	2.17	0.60
2:B:356:LEU:HD23	2:B:357:VAL:N	2.18	0.59
1:A:528:LYS:HA	1:A:531:LEU:HD12	1.85	0.57
1:A:913:ILE:HG22	1:A:913:ILE:O	2.04	0.56
1:A:530:GLN:HA	1:A:533:ALA:HB3	1.88	0.55
1:A:709:MET:HE2	1:A:841:ILE:HD13	1.90	0.53
1:A:208:THR:C	1:A:209:LEU:HD12	2.36	0.50
2:B:501:GLN:O	2:B:505:SER:OG	2.29	0.49
1:A:41:THR:HG23	1:A:44:THR:H	1.79	0.48
1:A:709:MET:HE1	1:A:847:LEU:HD21	1.96	0.48
1:A:480:PHE:O	1:A:482:SER:N	2.48	0.47
1:A:342:THR:O	1:A:471:PRO:HA	2.15	0.47
2:B:562:ARG:O	2:B:566:ILE:HD13	2.15	0.47
2:B:335:TRP:CD1	2:B:356:LEU:HD21	2.50	0.46
1:A:37:LEU:HD22	1:A:37:LEU:N	2.30	0.46
1:A:873:ASN:O	1:A:876:THR:HG22	2.17	0.45
2:B:355:PHE:CZ	2:B:427:PRO:HB3	2.52	0.43
1:A:820:MET:HE1	1:A:907:ALA:HB1	2.00	0.43
2:B:532:LYS:HA	2:B:535:ILE:HG22	2.00	0.43
1:A:974:THR:HG23	1:A:977:PHE:H	1.84	0.43
1:A:177:LEU:H	1:A:177:LEU:HD23	1.84	0.42
1:A:440:LYS:HA	1:A:476:GLU:HA	2.01	0.42
1:A:412:ARG:NE	1:A:412:ARG:HA	2.34	0.42
2:B:464:ASP:O	2:B:468:GLU:HG2	2.20	0.42
1:A:711:LYS:O	1:A:715:LEU:HG	2.20	0.42
1:A:525:GLU:O	1:A:526:ASN:C	2.62	0.42
1:A:807:LEU:HD12	1:A:846:GLY:HA3	2.01	0.42
1:A:300:ASP:HB3	1:A:697:MET:HG2	2.02	0.41
1:A:237:GLU:O	1:A:241:LEU:HD13	2.19	0.41
1:A:543:ILE:O	1:A:543:ILE:HG23	2.20	0.41
1:A:199:SER:N	1:A:200:PRO:CD	2.84	0.41
2:B:389:LYS:C	2:B:396:LEU:HD21	2.46	0.40
1:A:603:ASP:OD1	1:A:603:ASP:C	2.63	0.40
1:A:29:GLY:HA3	2:B:501:GLN:CA	2.52	0.40
1:A:29:GLY:HA3	2:B:501:GLN:HA	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	1000/1060 (94%)	953 (95%)	47 (5%)	0	100	100
2	B	266/300 (89%)	258 (97%)	8 (3%)	0	100	100
All	All	1266/1360 (93%)	1211 (96%)	55 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	923/960 (96%)	907 (98%)	16 (2%)	53	65
2	B	251/277 (91%)	246 (98%)	5 (2%)	48	63
All	All	1174/1237 (95%)	1153 (98%)	21 (2%)	51	65

All (21) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	36	CYS
1	A	197	ILE
1	A	329	VAL
1	A	342	THR
1	A	392	TYR
1	A	461	VAL
1	A	491	VAL
1	A	523	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	524	ARG
1	A	528	LYS
1	A	529	GLU
1	A	602	LEU
1	A	654	LEU
1	A	720	LYS
1	A	982	GLU
1	A	1036	LEU
2	B	369	THR
2	B	420	LEU
2	B	428	VAL
2	B	429	SER
2	B	434	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	362	HIS
1	A	384	ASN
1	A	526	ASN
1	A	554	HIS
1	A	855	HIS
1	A	1048	HIS
2	B	385	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	UJC	A	1101	-	37,38,38	1.70	6 (16%)	52,57,57	1.38	5 (9%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	UJC	A	1101	-	-	3/24/36/36	0/4/4/4

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	1101	UJC	C08-N10	7.46	1.41	1.34
3	A	1101	UJC	C03-N02	3.41	1.37	1.33
3	A	1101	UJC	C21-C19	2.09	1.41	1.37
3	A	1101	UJC	C28-C13	2.08	1.42	1.39
3	A	1101	UJC	C18-C19	2.06	1.41	1.37
3	A	1101	UJC	C07-C12	2.02	1.41	1.39

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	1101	UJC	C11-N10-C08	-4.51	111.47	114.03
3	A	1101	UJC	C34-C29-C11	4.25	123.68	120.26
3	A	1101	UJC	C28-C13-C12	-2.58	117.95	121.07
3	A	1101	UJC	C18-C19-C21	-2.44	120.53	123.50
3	A	1101	UJC	C07-C08-N10	2.31	107.51	106.51

There are no chirality outliers.


All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	1101	UJC	N10-C11-C29-C34
3	A	1101	UJC	N10-C11-C29-C30
3	A	1101	UJC	C12-C13-N14-C15

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	1008/1060 (95%)	-0.45	1 (0%) 92 86	102, 148, 202, 261	0
2	B	270/300 (90%)	-0.39	0 100 100	121, 185, 251, 291	0
All	All	1278/1360 (93%)	-0.44	1 (0%) 92 86	102, 154, 223, 291	0

All (1) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1050	GLY	2.6

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.


6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	UJC	A	1101	35/35	0.93	0.10	94,126,164,230	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers

as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.