



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 7, 2026 – 03:32 AM UTC

PDB ID : 4TTI / pdb_00004tti
Title : Crystal structure of double mutant E. Coli purine nucleoside phosphorylase with 4 FMC molecules
Authors : Stefanic, Z.; Bzowska, A.
Deposited on : 2014-06-21
Resolution : 1.89 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtrriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.89 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7789 (1.90-1.90)
Clashscore	190562	8410 (1.90-1.90)
Ramachandran outliers	187476	8333 (1.90-1.90)
Sidechain outliers	187428	8333 (1.90-1.90)
RSRZ outliers	180081	7790 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	237	95% 5%
1	B	237	89% 9% .
1	C	237	2% 84% 9% . 5%
1	D	237	2% 96% . .
1	E	237	3% 89% 11%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	237	 95% 5%

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 12099 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Purine nucleoside phosphorylase DeoD-type.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	237	1809	1145	311	337	16	0	3	0
1	B	237	1805	1144	308	337	16	0	3	0
1	C	224	1701	1082	286	317	16	0	3	0
1	D	237	1793	1134	307	337	15	0	1	0
1	E	237	1793	1135	305	337	16	0	1	0
1	F	237	1820	1151	317	337	15	0	4	0

There are 12 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	204	ALA	ASP	engineered mutation	UNP P0ABP8
A	217	ALA	ARG	engineered mutation	UNP P0ABP8
B	204	ALA	ASP	engineered mutation	UNP P0ABP8
B	217	ALA	ARG	engineered mutation	UNP P0ABP8
C	204	ALA	ASP	engineered mutation	UNP P0ABP8
C	217	ALA	ARG	engineered mutation	UNP P0ABP8
D	204	ALA	ASP	engineered mutation	UNP P0ABP8
D	217	ALA	ARG	engineered mutation	UNP P0ABP8
E	204	ALA	ASP	engineered mutation	UNP P0ABP8
E	217	ALA	ARG	engineered mutation	UNP P0ABP8
F	204	ALA	ASP	engineered mutation	UNP P0ABP8
F	217	ALA	ARG	engineered mutation	UNP P0ABP8

- Molecule 2 is (1S)-1-(7-amino-1H-pyrazolo[4,3-d]pyrimidin-3-yl)-1,4-anhydro-D-ribitol (CCD ID: FMC) (formula: C₁₀H₁₃N₅O₄).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
2	A	1	19	10	5	4	0	0
2	C	1	19	10	5	4	0	0
2	D	1	19	10	5	4	0	0
2	F	1	19	10	5	4	0	0

- Molecule 3 is PHOSPHATE ION (CCD ID: PO4) (formula: O₄P).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total O P 5 4 1	0	0
3	B	1	Total O P 5 4 1	0	0
3	C	1	Total O P 5 4 1	0	0
3	D	1	Total O P 5 4 1	0	0
3	E	1	Total O P 5 4 1	0	0
3	F	1	Total O P 5 4 1	0	0

- Molecule 4 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	C	1	Total O S 5 4 1	0	0
4	C	1	Total O S 5 4 1	0	0
4	D	1	Total O S 5 4 1	0	0
4	E	1	Total O S 5 4 1	0	0
4	F	1	Total O S 5 4 1	0	0

- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	237	Total O 237 237	0	0
5	B	183	Total O 183 183	9	0
5	C	206	Total O 206 206	0	0
5	D	192	Total O 192 192	0	0
5	E	193	Total O 193 193	6	0
5	F	236	Total O 236 236	0	0

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Purine nucleoside phosphorylase DeoD-type

Chain A: 





- Molecule 1: Purine nucleoside phosphorylase DeoD-type

Chain B: 




- Molecule 1: Purine nucleoside phosphorylase DeoD-type

Chain C: 





- Molecule 1: Purine nucleoside phosphorylase DeoD-type

Chain D: 



- Molecule 1: Purine nucleoside phosphorylase DeoD-type

Chain E: 



- Molecule 1: Purine nucleoside phosphorylase DeoD-type

Chain F: 95% 5%



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	61.93Å 124.17Å 189.44Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.70 – 1.89 47.70 – 1.89	Depositor EDS
% Data completeness (in resolution range)	98.8 (47.70-1.89) 98.7 (47.70-1.89)	Depositor EDS
R_{merge}	0.17	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.44 (at 1.90Å)	Xtrriage
Refinement program	PHENIX (phenix.refine: 1.8.2_1309)	Depositor
R, R_{free}	0.167 , 0.210 0.166 , 0.210	Depositor DCC
R_{free} test set	5807 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	17.7	Xtrriage
Anisotropy	0.400	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 47.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	12099	wwPDB-VP
Average B, all atoms (Å ²)	21.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.17% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FMC, PO4, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.45	0/1847	0.78	2/2488 (0.1%)
1	B	0.47	1/1843 (0.1%)	0.79	0/2484
1	C	0.43	0/1737	0.78	0/2339
1	D	0.44	0/1825	0.76	0/2461
1	E	0.44	0/1825	0.76	0/2460
1	F	0.45	0/1861	0.76	0/2506
All	All	0.45	1/10938 (0.0%)	0.77	2/14738 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	C	0	1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	66	ILE	CA-CB	6.17	1.57	1.54

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	95	LEU	CA-C-N	5.53	125.62	119.32
1	A	95	LEU	C-N-CA	5.53	125.62	119.32

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	C	219	THR	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1809	0	1830	6	0
1	B	1805	0	1826	15	0
1	C	1701	0	1721	15	0
1	D	1793	0	1799	5	0
1	E	1793	0	1804	15	0
1	F	1820	0	1847	10	0
2	A	19	0	13	1	0
2	C	19	0	13	2	0
2	D	19	0	13	1	0
2	F	19	0	13	2	0
3	A	5	0	0	0	0
3	B	5	0	0	0	0
3	C	5	0	0	1	0
3	D	5	0	0	0	0
3	E	5	0	0	0	0
3	F	5	0	0	0	0
4	C	10	0	0	1	0
4	D	5	0	0	0	0
4	E	5	0	0	1	0
4	F	5	0	0	0	0
5	A	237	0	0	1	0
5	B	183	0	0	2	0
5	C	206	0	0	1	0
5	D	192	0	0	0	0
5	E	193	0	0	3	0
5	F	236	0	0	5	0
All	All	12099	0	10879	65	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (65) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:219:THR:HB	1:C:221:PHE:H	1.39	0.87
1:C:219:THR:HB	1:C:221:PHE:N	2.05	0.70
1:C:222:ASN:ND2	4:C:303:SO4:O2	2.23	0.68
1:C:219:THR:HA	1:C:220:THR:HB	1.79	0.64
1:C:237:LYS:O	5:C:557:HOH:O	2.15	0.63
1:E:95:LEU:HD23	1:E:171:GLU:HG3	1.81	0.62
1:E:35:ASP:O	5:E:549:HOH:O	2.17	0.59
1:F:30:GLU:OE1	5:F:401:HOH:O	2.17	0.59
2:D:302:FMC:N3	2:D:302:FMC:H2'	2.18	0.58
1:B:101:ARG:HB3	1:B:220:THR:HG21	1.85	0.58
1:B:117:ARG:NH2	1:D:122:ASP:O	2.35	0.57
1:B:30:GLU:OE1	5:B:401:HOH:O	2.17	0.56
1:F:37[B]:ARG:NH1	5:F:612:HOH:O	2.38	0.56
1:E:83:LYS:NZ	5:E:560:HOH:O	2.38	0.56
1:F:37[B]:ARG:NH1	5:F:616:HOH:O	2.40	0.55
2:A:300:FMC:N3	2:A:300:FMC:H2'	2.22	0.54
1:E:220:THR:HG21	4:E:302:SO4:O1	2.08	0.53
1:A:57:ILE:HD11	1:A:232:VAL:HG11	1.91	0.53
1:E:15:VAL:HG22	1:E:84:LYS:HB2	1.90	0.53
1:F:205:HIS:CE1	1:F:207:ARG:HB2	2.43	0.53
1:B:100:LEU:HA	1:B:202:VAL:HG22	1.90	0.53
1:C:97:HIS:O	1:C:149:ARG:NH1	2.41	0.53
2:C:301:FMC:H2'	2:C:301:FMC:N3	2.23	0.53
1:C:24:ARG:HD3	1:C:221:PHE:CE1	2.45	0.52
1:E:220:THR:HG22	1:E:223:ASP:CG	2.34	0.51
1:B:101:ARG:HE	1:B:220:THR:CG2	2.23	0.51
1:E:220:THR:HG23	1:E:223:ASP:H	1.75	0.51
1:B:114:LYS:HD3	1:E:114:LYS:HG3	1.93	0.51
1:E:13:ALA:HB2	1:E:56:LYS:HG2	1.92	0.50
1:A:236:ASP:O	1:A:237:LYS:HB2	2.11	0.49
1:A:13:ALA:HB2	1:A:56:LYS:HG2	1.95	0.49
2:F:301:FMC:H2'	2:F:301:FMC:N3	2.28	0.49
1:B:212:THR:OG1	1:B:216:GLU:HG3	2.13	0.48
1:F:149[A]:ARG:NE	5:F:566:HOH:O	2.47	0.48
1:C:62:HIS:CE1	1:C:181:GLU:HG2	2.48	0.48
1:B:74:LYS:HD3	1:B:74:LYS:C	2.39	0.47
5:B:405:HOH:O	1:D:172:LYS:HG3	2.13	0.47
1:C:2:THR:OG1	1:C:75:GLU:OE1	2.31	0.47
1:E:220:THR:HG23	1:E:222:ASN:N	2.30	0.46
1:D:62:HIS:CE1	1:D:181:GLU:HG2	2.50	0.46
1:B:73:THR:HG22	1:B:85:ILE:HD13	1.97	0.46
1:C:29:ALA:HA	1:C:33:LEU:HD22	1.98	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:43:ARG:HA	1:E:21:ASP:OD1	2.16	0.45
1:E:220:THR:CG2	1:E:223:ASP:H	2.29	0.45
2:C:301:FMC:H1'	3:C:302:PO4:O2	2.17	0.45
2:F:301:FMC:N7	5:F:599:HOH:O	2.33	0.44
1:C:73:THR:HG22	1:C:85:ILE:HD13	1.99	0.44
1:D:172:LYS:HG2	1:D:173:TYR:CD1	2.53	0.44
1:E:76:LEU:HB3	1:E:82:VAL:HG21	1.99	0.43
1:C:173:TYR:CZ	1:E:119:ARG:HB3	2.52	0.43
1:B:212:THR:HG23	1:B:217:ALA:HB2	2.00	0.43
1:B:107[A]:MET:HE2	1:B:107[A]:MET:HB2	1.93	0.43
1:D:107:MET:HE3	1:D:107:MET:HB3	1.87	0.42
1:C:43:ARG:HA	1:F:21:ASP:OD1	2.20	0.42
1:A:41:ASN:OD1	5:A:583:HOH:O	2.22	0.42
1:C:90:SER:HB2	1:C:203:SER:HB3	2.02	0.42
1:B:101:ARG:HE	1:B:220:THR:HG21	1.84	0.42
1:C:142:LYS:HE3	1:C:142:LYS:HB2	1.85	0.41
1:B:105:ILE:HG23	1:B:197:LEU:HD11	2.02	0.41
1:F:74:LYS:HD3	1:F:74:LYS:C	2.45	0.41
1:E:226:LYS:NZ	5:E:586:HOH:O	2.45	0.41
1:B:95:LEU:HD23	1:B:171:GLU:HG3	2.03	0.41
1:F:62:HIS:CE1	1:F:181:GLU:HG2	2.55	0.41
1:A:135[A]:ARG:HH22	1:F:139:ASP:CG	2.29	0.40
1:A:128:ILE:HD12	1:F:107:MET:HE3	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	238/237 (100%)	232 (98%)	6 (2%)	0	100 100
1	B	238/237 (100%)	232 (98%)	6 (2%)	0	100 100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	C	223/237 (94%)	216 (97%)	6 (3%)	1 (0%)	30	22
1	D	236/237 (100%)	225 (95%)	9 (4%)	2 (1%)	16	8
1	E	236/237 (100%)	225 (95%)	9 (4%)	2 (1%)	16	8
1	F	239/237 (101%)	232 (97%)	7 (3%)	0	100	100
All	All	1410/1422 (99%)	1362 (97%)	43 (3%)	5 (0%)	30	22

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	209	HIS
1	D	215	ALA
1	E	212	THR
1	E	215	ALA
1	C	220	THR

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	189/186 (102%)	184 (97%)	5 (3%)	40	35
1	B	189/186 (102%)	179 (95%)	10 (5%)	20	12
1	C	179/186 (96%)	173 (97%)	6 (3%)	32	25
1	D	186/186 (100%)	182 (98%)	4 (2%)	45	42
1	E	187/186 (100%)	179 (96%)	8 (4%)	26	18
1	F	190/186 (102%)	186 (98%)	4 (2%)	47	44
All	All	1120/1116 (100%)	1083 (97%)	37 (3%)	36	26

All (37) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	37	ARG
1	A	107[A]	MET

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	107[B]	MET
1	A	179	GLU
1	A	237	LYS
1	B	15[A]	VAL
1	B	15[B]	VAL
1	B	107[A]	MET
1	B	107[B]	MET
1	B	179	GLU
1	B	202	VAL
1	B	207	ARG
1	B	208	THR
1	B	212	THR
1	B	237	LYS
1	C	33	LEU
1	C	101	ARG
1	C	104	VAL
1	C	149	ARG
1	C	165	GLU
1	C	179	GLU
1	D	107	MET
1	D	179	GLU
1	D	207	ARG
1	D	209	HIS
1	E	23	LEU
1	E	34	GLU
1	E	37	ARG
1	E	100	LEU
1	E	120	PHE
1	E	165	GLU
1	E	179	GLU
1	E	210	GLU
1	F	37[A]	ARG
1	F	37[B]	ARG
1	F	175	ILE
1	F	179	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	40	ASN
1	D	218	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

15 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	FMC	F	301	-	19,21,21	1.34	3 (15%)	20,31,31	1.27	3 (15%)
3	PO4	B	301	-	4,4,4	0.88	0	6,6,6	0.64	0
4	SO4	D	303	-	4,4,4	0.22	0	6,6,6	0.20	0
3	PO4	C	302	-	4,4,4	1.00	0	6,6,6	0.54	0
3	PO4	E	301	-	4,4,4	0.93	0	6,6,6	0.60	0
4	SO4	C	304	-	4,4,4	0.24	0	6,6,6	0.11	0
3	PO4	A	301	-	4,4,4	1.26	0	6,6,6	0.85	0
2	FMC	C	301	-	19,21,21	1.28	3 (15%)	20,31,31	1.25	2 (10%)
3	PO4	D	301	-	4,4,4	0.95	0	6,6,6	0.39	0
3	PO4	F	302	-	4,4,4	1.19	0	6,6,6	0.40	0
4	SO4	E	302	-	4,4,4	0.23	0	6,6,6	0.26	0
2	FMC	D	302	-	19,21,21	1.36	3 (15%)	20,31,31	1.23	2 (10%)
4	SO4	F	303	-	4,4,4	0.23	0	6,6,6	0.20	0
4	SO4	C	303	-	4,4,4	0.19	0	6,6,6	0.15	0
2	FMC	A	300	-	19,21,21	1.30	3 (15%)	20,31,31	1.25	2 (10%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FMC	F	301	-	-	5/6/22/22	0/3/3/3
2	FMC	A	300	-	-	2/6/22/22	0/3/3/3
2	FMC	C	301	-	-	2/6/22/22	0/3/3/3
2	FMC	D	302	-	-	4/6/22/22	0/3/3/3

All (12) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	302	FMC	C9-N8	4.33	1.36	1.29
2	F	301	FMC	C9-N8	3.99	1.36	1.29
2	C	301	FMC	C9-N8	3.93	1.35	1.29
2	A	300	FMC	C9-N8	3.78	1.35	1.29
2	A	300	FMC	N7-N8	2.30	1.42	1.36
2	D	302	FMC	C5-C6	2.29	1.48	1.40
2	A	300	FMC	C5-C6	2.24	1.48	1.40
2	D	302	FMC	N7-N8	2.23	1.41	1.36
2	C	301	FMC	C5-C6	2.13	1.48	1.40
2	F	301	FMC	N7-N8	2.10	1.41	1.36
2	C	301	FMC	N7-N8	2.09	1.41	1.36
2	F	301	FMC	C5-C6	2.07	1.48	1.40

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	301	FMC	N3-C2-N1	-3.21	123.72	128.58
2	D	302	FMC	N3-C2-N1	-3.07	123.93	128.58
2	F	301	FMC	N3-C2-N1	-3.02	124.01	128.58
2	A	300	FMC	N3-C2-N1	-2.98	124.07	128.58
2	C	301	FMC	C2-N1-C6	2.28	122.47	118.73
2	F	301	FMC	C2-N1-C6	2.12	122.21	118.73
2	A	300	FMC	C9-N8-N7	2.10	110.59	106.38
2	D	302	FMC	C9-N8-N7	2.04	110.47	106.38
2	F	301	FMC	C9-N8-N7	2.00	110.40	106.38

There are no chirality outliers.

All (13) torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
2	A	300	FMC	C2'-C1'-C9-C4
2	A	300	FMC	C2'-C1'-C9-N8
2	D	302	FMC	C2'-C1'-C9-C4
2	D	302	FMC	C2'-C1'-C9-N8
2	F	301	FMC	C2'-C1'-C9-C4
2	F	301	FMC	C2'-C1'-C9-N8
2	F	301	FMC	O4'-C4'-C5'-O5'
2	C	301	FMC	C2'-C1'-C9-N8
2	F	301	FMC	O4'-C1'-C9-N8
2	F	301	FMC	C3'-C4'-C5'-O5'
2	D	302	FMC	O4'-C4'-C5'-O5'
2	C	301	FMC	C2'-C1'-C9-C4
2	D	302	FMC	C3'-C4'-C5'-O5'


There are no ring outliers.

7 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	F	301	FMC	2	0
3	C	302	PO4	1	0
2	C	301	FMC	2	0
4	E	302	SO4	1	0
2	D	302	FMC	1	0
4	C	303	SO4	1	0
2	A	300	FMC	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	237/237 (100%)	-0.60	1 (0%) 88 90	8, 17, 30, 43	3 (1%)
1	B	237/237 (100%)	-0.42	0 100 100	10, 18, 39, 54	3 (1%)
1	C	224/237 (94%)	-0.44	5 (2%) 62 66	8, 17, 40, 62	3 (1%)
1	D	237/237 (100%)	-0.36	5 (2%) 63 67	11, 19, 40, 81	1 (0%)
1	E	237/237 (100%)	-0.28	6 (2%) 58 62	11, 20, 40, 71	1 (0%)
1	F	237/237 (100%)	-0.63	1 (0%) 88 90	9, 16, 27, 42	4 (1%)
All	All	1409/1422 (99%)	-0.46	18 (1%) 75 78	8, 18, 37, 81	15 (1%)

All (18) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	E	214	ALA	4.9
1	E	212	THR	4.3
1	D	214	ALA	4.2
1	E	215	ALA	4.0
1	E	213	THR	3.9
1	C	219	THR	3.5
1	C	100	LEU	3.4
1	D	208	THR	2.8
1	C	97	HIS	2.5
1	C	204	ALA	2.4
1	D	207	ARG	2.4
1	D	209	HIS	2.4
1	E	211	GLN	2.4
1	A	1	ALA	2.3
1	C	101	ARG	2.2
1	F	237	LYS	2.2
1	E	209	HIS	2.0
1	D	213	THR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	SO4	C	304	5/5	0.78	0.13	38,38,52,53	5
4	SO4	D	303	5/5	0.81	0.14	36,39,42,44	5
2	FMC	C	301	19/19	0.82	0.13	15,15,15,15	19
2	FMC	F	301	19/19	0.83	0.12	15,15,15,15	19
2	FMC	D	302	19/19	0.84	0.11	15,15,15,15	19
2	FMC	A	300	19/19	0.85	0.12	15,15,15,15	19
4	SO4	C	303	5/5	0.88	0.10	36,37,43,47	5
4	SO4	E	302	5/5	0.91	0.10	29,32,34,43	5
3	PO4	B	301	5/5	0.92	0.08	30,31,36,38	0
4	SO4	F	303	5/5	0.95	0.07	16,20,25,29	5
3	PO4	E	301	5/5	0.97	0.07	24,27,33,33	0
3	PO4	D	301	5/5	0.97	0.07	24,24,28,29	0
3	PO4	F	302	5/5	0.99	0.03	10,10,14,15	0
3	PO4	A	301	5/5	0.99	0.04	13,13,16,16	0
3	PO4	C	302	5/5	0.99	0.04	23,23,27,31	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around FMC C 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around FMC F 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around FMC D 302:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.