



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 03:23 AM UTC

PDB ID : 1TV8 / pdb_00001tv8
Title : Structure of MoaA in complex with S-adenosylmethionine
Authors : Haenzelmann, P.; Schindelin, H.
Deposited on : 2004-06-28
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6164 (2.20-2.20)
Clashscore	190562	6851 (2.20-2.20)
Ramachandran outliers	187476	6768 (2.20-2.20)
Sidechain outliers	187428	6769 (2.20-2.20)
RSRZ outliers	180081	6166 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	340	6% 81% 12% ...
1	B	340	10% 70% 20% 5% .

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 5505 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Molybdenum cofactor biosynthesis protein A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	327	Total	C	N	O	S	0	0	0
			2641	1673	455	500	13			
1	B	326	Total	C	N	O	S	0	0	0
			2632	1668	454	497	13			

- Molecule 2 is SULFATE ION (CCD ID: SO4) (formula: O₄S).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 3 is S-ADENOSYLMETHIONINE (CCD ID: SAM) (formula: C₁₅H₂₂N₆O₅S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	S	0	0
			27	15	6	5	1		
3	B	1	Total	C	N	O	S	0	0
			27	15	6	5	1		

- Molecule 4 is IRON/SULFUR CLUSTER (CCD ID: SF4) (formula: Fe₄S₄).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	Fe	S	0	0
			8	4	4		
4	A	1	Total	Fe	S	0	0
			8	4	4		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	Fe	S	0	0
			8	4	4		
4	B	1	Total	Fe	S	0	0
			8	4	4		

- Molecule 5 is (2R,3S)-1,4-DIMERCAPTOBUTANE-2,3-DIOL (CCD ID: DTU) (formula: $C_4H_{10}O_2S_2$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	O	S	0	0
			8	4	2	2		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	96	Total	O	0	0
			96	96		
6	B	32	Total	O	0	0
			32	32		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Molybdenum cofactor biosynthesis protein A



• Molecule 1: Molybdenum cofactor biosynthesis protein A



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	47.98Å 103.31Å 189.93Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	95.35 – 2.20 94.96 – 2.21	Depositor EDS
% Data completeness (in resolution range)	96.7 (95.35-2.20) 96.7 (94.96-2.21)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.75 (at 2.20Å)	Xtriage
Refinement program	REFMAC 5.1.24	Depositor
R, R_{free}	0.213 , 0.241 0.234 , 0.254	Depositor DCC
R_{free} test set	2399 reflections (5.09%)	wwPDB-VP
Wilson B-factor (Å ²)	48.9	Xtriage
Anisotropy	0.535	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 45.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	5505	wwPDB-VP
Average B, all atoms (Å ²)	43.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.51% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SAM, SF4, SO4, DTU

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.11	7/2684 (0.3%)	1.05	6/3613 (0.2%)
1	B	1.18	17/2675 (0.6%)	1.08	8/3601 (0.2%)
All	All	1.15	24/5359 (0.4%)	1.06	14/7214 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	3
1	B	0	1
All	All	0	4

All (24) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	170	LYS	C-O	15.89	1.44	1.24
1	B	170	LYS	CA-C	13.73	1.71	1.52
1	B	131	ASP	C-O	11.88	1.38	1.23
1	B	169	GLN	CA-C	9.74	1.64	1.52
1	B	133	THR	C-O	8.33	1.33	1.24
1	B	134	LEU	N-CA	8.29	1.56	1.46
1	B	131	ASP	C-N	8.26	1.46	1.33
1	B	171	GLY	N-CA	7.30	1.56	1.45
1	A	327	ASN	CG-ND2	6.93	1.47	1.33
1	A	327	ASN	CG-OD1	6.70	1.36	1.23
1	A	79	MET	CG-SD	-6.52	1.64	1.80
1	A	197	MET	SD-CE	-6.44	1.63	1.79
1	B	174	ASP	C-O	6.19	1.31	1.24
1	B	137	SER	CA-CB	6.02	1.63	1.53

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	168	ILE	C-N	5.96	1.41	1.33
1	A	89	LYS	C-O	-5.79	1.17	1.24
1	B	169	GLN	C-O	5.76	1.30	1.24
1	B	137	SER	CB-OG	5.66	1.53	1.42
1	B	144	LYS	CA-C	5.65	1.60	1.52
1	A	39	ASP	C-O	5.61	1.30	1.24
1	B	139	ASN	CA-C	5.23	1.59	1.52
1	B	134	LEU	C-O	5.08	1.30	1.24
1	B	14	ARG	C-O	5.05	1.30	1.23
1	A	118	ALA	CA-CB	-5.01	1.45	1.53

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	202	ASP	N-CA-C	10.25	126.11	113.50
1	B	170	LYS	N-CA-C	8.69	121.91	111.82
1	A	210	VAL	N-CA-C	8.25	126.49	109.34
1	B	170	LYS	CA-C-O	6.60	127.56	119.97
1	A	245	ASP	CA-C-N	6.12	132.72	121.70
1	A	245	ASP	C-N-CA	6.12	132.72	121.70
1	B	81	ARG	N-CA-C	6.09	117.92	111.28
1	B	133	THR	N-CA-C	-6.08	104.34	110.97
1	B	20	VAL	CB-CA-C	5.82	116.35	110.65
1	B	134	LEU	CA-C-O	5.77	126.61	119.97
1	A	327	ASN	OD1-CG-ND2	5.44	128.04	122.60
1	B	73	THR	OG1-CB-CG2	-5.09	99.13	109.30
1	A	41	PHE	N-CA-C	-5.04	102.92	110.23
1	B	35	GLU	N-CA-C	-5.03	102.94	110.23

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	201	ASN	Peptide
1	A	209	LYS	Peptide
1	A	210	VAL	Peptide
1	B	328	ARG	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2641	0	2637	31	0
1	B	2632	0	2631	53	0
2	A	5	0	0	0	0
2	B	5	0	0	0	0
3	A	27	0	21	1	0
3	B	27	0	21	5	0
4	A	16	0	0	0	0
4	B	16	0	0	0	0
5	A	8	0	9	2	0
6	A	96	0	0	7	0
6	B	32	0	0	11	0
All	All	5505	0	5319	84	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (84) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:266:ARG:HB3	6:A:1598:HOH:O	1.49	1.12
1:B:32:MET:HE3	6:B:2529:HOH:O	1.62	0.98
1:B:30:TYR:HE1	3:B:2501:SAM:HN61	1.12	0.98
1:A:245:ASP:HB3	1:A:246:ASN:HB2	1.48	0.96
1:B:136:GLN:CG	1:B:143:ILE:O	2.15	0.95
1:B:136:GLN:HG2	1:B:143:ILE:O	1.71	0.90
1:A:245:ASP:HB2	1:A:246:ASN:HB3	1.54	0.88
1:A:245:ASP:CB	1:A:246:ASN:CB	2.55	0.85
1:A:245:ASP:CB	1:A:246:ASN:HB3	2.07	0.85
1:A:165:ASN:HD21	1:A:192:ARG:HH11	1.25	0.84
1:A:245:ASP:HB3	1:A:246:ASN:CB	2.12	0.78
1:B:136:GLN:HG3	1:B:143:ILE:O	1.80	0.78
1:B:15:ASP:N	6:B:2521:HOH:O	2.15	0.78
1:B:30:TYR:CE1	3:B:2501:SAM:N6	2.56	0.73
1:A:71:ARG:HH11	1:A:102:THR:HG22	1.55	0.71
1:B:34:LYS:NZ	6:B:2510:HOH:O	2.23	0.70

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:14:ARG:CA	6:B:2521:HOH:O	2.44	0.66
1:B:73:THR:HG22	1:B:74:GLY:H	1.61	0.64
1:A:169:GLN:HE21	1:A:209:LYS:NZ	1.96	0.63
1:B:14:ARG:HA	6:B:2521:HOH:O	1.98	0.63
1:B:98:ASP:HB3	6:B:2509:HOH:O	1.98	0.62
1:A:165:ASN:HD21	1:A:192:ARG:NH1	1.96	0.61
1:B:301:LEU:HD22	1:B:305:PHE:CE1	2.36	0.60
1:B:280:PHE:O	1:B:281:ALA:HB2	2.04	0.57
1:B:169:GLN:H	1:B:173:ASN:HB2	1.68	0.57
1:A:169:GLN:HE21	1:A:209:LYS:HZ2	1.52	0.56
1:A:245:ASP:HB2	1:A:246:ASN:CB	2.26	0.55
1:A:297:THR:HG22	1:A:300:GLU:H	1.73	0.54
1:A:6:LYS:NZ	1:B:323:GLN:HE22	2.06	0.54
1:B:26:PHE:CE1	1:B:104:ASN:HB2	2.42	0.54
1:B:87:ILE:HD12	1:B:115:LEU:HD22	1.89	0.53
3:B:2501:SAM:O3'	6:B:2525:HOH:O	2.15	0.53
1:B:13:ILE:HG12	1:B:309:TRP:CD2	2.43	0.53
1:B:177:ILE:HA	1:B:193:PHE:CE2	2.44	0.53
1:B:14:ARG:O	1:B:265:THR:O	2.28	0.52
1:B:279:LEU:HD11	6:B:2533:HOH:O	2.09	0.52
1:B:30:TYR:CD1	1:B:30:TYR:O	2.64	0.51
1:B:205:TRP:CE3	1:B:256:VAL:HG22	2.47	0.50
1:B:30:TYR:O	1:B:30:TYR:CG	2.64	0.49
1:B:14:ARG:O	1:B:15:ASP:HB2	2.12	0.49
1:B:23:ARG:NE	6:B:2520:HOH:O	2.46	0.49
1:B:58:ILE:HD13	1:B:269:LEU:HD21	1.95	0.49
1:A:14:ARG:HD3	6:A:1594:HOH:O	2.13	0.48
1:B:134:LEU:O	1:B:138:ILE:HG13	2.14	0.48
1:B:14:ARG:O	1:B:15:ASP:CB	2.61	0.48
1:B:14:ARG:C	6:B:2521:HOH:O	2.50	0.48
1:A:73:THR:OG1	3:A:1501:SAM:HG2	2.12	0.48
1:B:73:THR:CG2	3:B:2501:SAM:HG2	2.44	0.47
1:A:116:TYR:CD2	1:A:116:TYR:C	2.93	0.46
1:A:165:ASN:ND2	1:A:192:ARG:HH11	2.02	0.46
1:A:260:PHE:CZ	5:A:1502:DTU:O3	2.64	0.46
1:A:202:ASP:N	1:A:203:ASN:HA	2.30	0.46
6:A:1592:HOH:O	1:B:328:ARG:HG3	2.14	0.46
1:A:58:ILE:HD13	1:A:269:LEU:HD11	1.98	0.46
1:B:172:ILE:HG22	1:B:173:ASN:HD22	1.81	0.46
1:B:259:SER:HB3	1:B:318:ASP:OD1	2.15	0.46
1:A:74:GLY:HA3	1:A:75:GLY:C	2.41	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:47:ASN:H	1:A:47:ASN:HD22	1.64	0.45
1:B:24:CYS:O	1:B:27:ARG:NH1	2.47	0.45
1:A:14:ARG:NE	6:A:1594:HOH:O	2.49	0.45
1:A:153:ASP:OD2	1:A:187:LYS:NZ	2.50	0.45
1:B:127:LEU:O	1:B:180:MET:HE1	2.17	0.45
1:A:97:GLU:HG2	6:A:1596:HOH:O	2.17	0.45
1:A:266:ARG:CB	6:A:1598:HOH:O	2.31	0.44
1:B:61:VAL:CG1	1:B:302:LYS:HA	2.48	0.44
1:A:292:ILE:HG22	1:A:301:LEU:HD13	1.98	0.44
1:A:260:PHE:HZ	5:A:1502:DTU:O3	1.99	0.44
1:B:178:ILE:N	1:B:178:ILE:HD12	2.33	0.44
1:B:133:THR:O	1:B:136:GLN:N	2.51	0.44
1:B:27:ARG:CZ	1:B:141:ARG:NH2	2.81	0.44
1:B:69:LYS:NZ	1:B:98:ASP:OD2	2.50	0.43
1:B:178:ILE:HD12	1:B:178:ILE:H	1.84	0.43
1:A:14:ARG:CD	6:A:1594:HOH:O	2.67	0.42
1:B:178:ILE:HB	1:B:179:PRO:HD3	2.01	0.42
1:B:205:TRP:CD2	1:B:256:VAL:HG22	2.54	0.42
1:B:176:GLN:C	1:B:179:PRO:HD2	2.45	0.42
1:B:224:PHE:HB3	1:B:226:ILE:HD12	2.02	0.41
1:B:73:THR:HG22	3:B:2501:SAM:HG2	2.01	0.41
1:B:266:ARG:CD	6:B:2533:HOH:O	2.68	0.41
1:B:280:PHE:O	1:B:281:ALA:CB	2.68	0.41
1:A:58:ILE:HG21	1:A:269:LEU:HD11	2.03	0.40
1:B:111:HIS:O	1:B:115:LEU:HG	2.21	0.40
1:B:79:MET:HE3	1:B:79:MET:HB3	1.96	0.40
1:B:132:ASP:CG	1:B:144:LYS:HB3	2.46	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	325/340 (96%)	309 (95%)	13 (4%)	3 (1%)	14	14
1	B	324/340 (95%)	312 (96%)	6 (2%)	6 (2%)	6	4
All	All	649/680 (95%)	621 (96%)	19 (3%)	9 (1%)	9	7

All (9) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	210	VAL
1	A	246	ASN
1	B	15	ASP
1	B	174	ASP
1	B	36	VAL
1	B	281	ALA
1	B	295	GLY
1	B	315	ARG
1	A	209	LYS

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	291/302 (96%)	271 (93%)	20 (7%)	14	16
1	B	290/302 (96%)	263 (91%)	27 (9%)	8	9
All	All	581/604 (96%)	534 (92%)	47 (8%)	11	12

All (47) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	5	ILE
1	A	17	ARG
1	A	18	LEU
1	A	34	LYS
1	A	46	LYS
1	A	47	ASN
1	A	102	THR

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	110	LYS
1	A	114	LYS
1	A	120	LEU
1	A	134	LEU
1	A	192	ARG
1	A	210	VAL
1	A	226	ILE
1	A	255	SER
1	A	282	THR
1	A	292	ILE
1	A	297	THR
1	A	299	GLU
1	A	315	ARG
1	B	13	ILE
1	B	20	VAL
1	B	34	LYS
1	B	47	ASN
1	B	68	LYS
1	B	71	ARG
1	B	81	ARG
1	B	83	LEU
1	B	101	LEU
1	B	120	LEU
1	B	134	LEU
1	B	148	ILE
1	B	192	ARG
1	B	197	MET
1	B	215	GLU
1	B	226	ILE
1	B	246	ASN
1	B	253	ILE
1	B	256	VAL
1	B	258	GLN
1	B	269	LEU
1	B	282	THR
1	B	292	ILE
1	B	301	LEU
1	B	303	GLU
1	B	315	ARG
1	B	329	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	47	ASN
1	A	91	ASN
1	A	92	GLN
1	A	165	ASN
1	A	169	GLN
1	A	203	ASN
1	B	136	GLN
1	B	173	ASN
1	B	222	GLN
1	B	243	HIS
1	B	323	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

9 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	SAM	A	1501	4	27,29,29	1.11	3 (11%)	34,42,42	2.39	13 (38%)
2	SO4	B	2403	-	4,4,4	0.34	0	6,6,6	0.57	0
5	DTU	A	1502	4	7,7,7	0.90	0	4,8,8	2.43	1 (25%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	SF4	A	1402	1,5	0,12,12	-	-	-		
4	SF4	A	1401	1,3	0,12,12	-	-	-		
2	SO4	A	1403	-	4,4,4	0.35	0	6,6,6	1.07	0
4	SF4	B	2401	1,3	0,12,12	-	-	-		
4	SF4	B	2402	1	0,12,12	-	-	-		
3	SAM	B	2501	4	27,29,29	1.21	3 (11%)	34,42,42	2.22	13 (38%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	SAM	A	1501	4	-	3/17/33/33	0/3/3/3
5	DTU	A	1502	4	-	4/8/8/8	-
4	SF4	A	1402	1,5	-	-	0/6/5/5
4	SF4	A	1401	1,3	-	-	0/6/5/5
4	SF4	B	2401	1,3	-	-	0/6/5/5
4	SF4	B	2402	1	-	-	0/6/5/5
3	SAM	B	2501	4	-	2/17/33/33	0/3/3/3

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	2501	SAM	C2-N3	3.07	1.39	1.33
3	B	2501	SAM	C8-N7	2.81	1.37	1.31
3	A	1501	SAM	C2-N3	2.73	1.38	1.33
3	B	2501	SAM	C2-N1	2.60	1.38	1.33
3	A	1501	SAM	C2-N1	2.50	1.38	1.33
3	A	1501	SAM	C8-N7	2.38	1.36	1.31

All (27) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	2501	SAM	N3-C2-N1	-6.50	118.75	128.58
3	A	1501	SAM	C5-C4-N3	-5.82	118.71	126.72
3	A	1501	SAM	N3-C2-N1	-5.21	120.69	128.58
3	A	1501	SAM	O4'-C1'-N9	4.73	117.18	108.09
5	A	1502	DTU	C3-C4-S4	-4.44	102.04	114.43
3	B	2501	SAM	C5-C4-N3	-4.42	120.63	126.72
3	A	1501	SAM	C2-N3-C4	4.05	121.72	111.83

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	1501	SAM	N3-C4-N9	3.85	133.72	127.17
3	B	2501	SAM	C2-N3-C4	3.71	120.88	111.83
3	A	1501	SAM	C5-N7-C8	3.61	109.13	103.45
3	A	1501	SAM	N9-C8-N7	-3.43	109.08	113.94
3	B	2501	SAM	C5-N7-C8	3.17	108.42	103.45
3	B	2501	SAM	OXT-C-O	-3.10	117.04	124.08
3	B	2501	SAM	N9-C8-N7	-3.06	109.60	113.94
3	A	1501	SAM	CG-SD-C5'	-3.01	96.06	103.43
3	A	1501	SAM	C4-C5-N7	-2.99	107.16	110.58
3	B	2501	SAM	O4'-C1'-N9	2.97	113.79	108.09
3	A	1501	SAM	O4'-C1'-C2'	-2.77	100.69	106.62
3	B	2501	SAM	C4-C5-N7	-2.72	107.47	110.58
3	B	2501	SAM	O4'-C4'-C5'	2.64	115.55	108.88
3	B	2501	SAM	C5-C4-N9	2.51	108.54	105.81
3	A	1501	SAM	C4'-O4'-C1'	-2.49	103.98	109.47
3	B	2501	SAM	C4'-O4'-C1'	-2.36	104.25	109.47
3	A	1501	SAM	O4'-C4'-C5'	2.28	114.65	108.88
3	B	2501	SAM	N3-C4-N9	2.15	130.83	127.17
3	A	1501	SAM	C6-C5-C4	2.12	120.07	117.18
3	B	2501	SAM	O2'-C2'-C3'	-2.01	105.38	111.82

There are no chirality outliers.

All (9) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	1501	SAM	C-CA-CB-CG
5	A	1502	DTU	O2-C2-C3-C4
5	A	1502	DTU	C2-C3-C4-S4
5	A	1502	DTU	O3-C3-C4-S4
5	A	1502	DTU	O2-C2-C3-O3
3	B	2501	SAM	C-CA-CB-CG
3	A	1501	SAM	N-CA-CB-CG
3	A	1501	SAM	CA-CB-CG-SD
3	B	2501	SAM	CA-CB-CG-SD

There are no ring outliers.

3 monomers are involved in 8 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	1501	SAM	1	0
5	A	1502	DTU	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	2501	SAM	5	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	327/340 (96%)	0.68	19 (5%) 29 25	28, 40, 55, 62	0
1	B	326/340 (95%)	0.89	33 (10%) 12 10	29, 42, 54, 61	0
All	All	653/680 (96%)	0.78	52 (7%) 18 16	28, 41, 55, 62	0

All (52) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	30	TYR	5.4
1	B	283	VAL	5.1
1	B	282	THR	4.2
1	B	226	ILE	3.9
1	B	40	ASP	3.8
1	B	35	GLU	3.6
1	B	136	GLN	3.6
1	B	131	ASP	3.5
1	B	203	ASN	3.5
1	A	201	ASN	3.3
1	B	171	GLY	3.2
1	B	193	PHE	3.1
1	B	133	THR	3.1
1	A	95	GLY	3.1
1	B	21	THR	3.0
1	B	143	ILE	2.9
1	B	248	VAL	2.9
1	B	28	CYS	2.9
1	A	98	ASP	2.9
1	A	178	ILE	2.9
1	A	292	ILE	2.8
1	A	303	GLU	2.8
1	B	280	PHE	2.6
1	B	31	CYS	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	39	ASP	2.6
1	B	47	ASN	2.6
1	B	260	PHE	2.6
1	A	202	ASP	2.5
1	A	171	GLY	2.5
1	B	26	PHE	2.4
1	A	36	VAL	2.4
1	A	260	PHE	2.4
1	B	37	PHE	2.4
1	B	104	ASN	2.3
1	A	182	GLU	2.3
1	A	40	ASP	2.3
1	A	280	PHE	2.3
1	B	36	VAL	2.3
1	A	279	LEU	2.3
1	B	173	ASN	2.2
1	B	281	ALA	2.2
1	B	174	ASP	2.2
1	B	325	VAL	2.2
1	B	290	ALA	2.2
1	B	94	ASP	2.1
1	A	297	THR	2.1
1	B	201	ASN	2.1
1	A	15	ASP	2.1
1	A	328	ARG	2.1
1	B	24	CYS	2.1
1	B	130	ILE	2.1
1	A	94	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,




median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	SO4	B	2403	5/5	0.66	0.19	81,83,85,86	0
5	DTU	A	1502	8/8	0.81	0.18	77,87,88,95	0
3	SAM	B	2501	27/27	0.82	0.20	67,78,80,80	0
2	SO4	A	1403	5/5	0.84	0.18	70,72,76,79	0
4	SF4	B	2401	8/8	0.93	0.19	37,42,44,44	0
3	SAM	A	1501	27/27	0.93	0.12	54,71,75,75	0
4	SF4	B	2402	8/8	0.94	0.13	35,37,40,45	0
4	SF4	A	1402	8/8	0.97	0.08	35,38,38,44	0
4	SF4	A	1401	8/8	0.97	0.07	36,36,37,37	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around SAM B 2501:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around SAM A 1501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.