



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 14, 2026 – 05:30 PM UTC

PDB ID : 4TV8 / pdb_00004tv8
Title : Tubulin-Maytansine complex
Authors : Prota, A.E.; Bargsten, K.; Diaz, J.F.; Marsh, M.; Cuevas, C.; Liniger, M.; Neuhaus, C.; Andreu, J.M.; Altmann, K.H.; Steinmetz, M.O.
Deposited on : 2014-06-26
Resolution : 2.10 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6658 (2.10-2.10)
Clashscore	190562	7164 (2.10-2.10)
Ramachandran outliers	187476	7099 (2.10-2.10)
Sidechain outliers	187428	7100 (2.10-2.10)
RSRZ outliers	180081	6662 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	451	 2% 87% 10% .
1	C	451	 2% 88% 9% .
2	B	445	 4% 89% 7% .
2	D	445	 3% 86% 9% 5%
3	E	143	 8% 76% 9% 15%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
4	F	384	8% 79% 8% 13%

2 Entry composition [i](#)

There are 13 unique types of molecules in this entry. The entry contains 18278 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Tubulin alpha-1B chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	436	Total	C	N	O	S	0	7	0
			3433	2175	580	654	24			
1	C	440	Total	C	N	O	S	0	11	0
			3483	2208	585	665	25			

- Molecule 2 is a protein called Tubulin beta-2B chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	427	Total	C	N	O	S	0	4	0
			3365	2116	572	650	27			
2	D	422	Total	C	N	O	S	0	3	0
			3326	2091	563	644	28			

- Molecule 3 is a protein called Stathmin-4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	E	121	Total	C	N	O	S	0	1	0
			1006	621	182	198	5			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
E	3	MET	-	expression tag	UNP P63043
E	4	ALA	-	expression tag	UNP P63043

- Molecule 4 is a protein called Tubulin-Tyrosine Ligase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	F	335	Total	C	N	O	S	0	2	0
			2763	1776	473	500	14			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
F	379	HIS	-	expression tag	UNP E1BQ43
F	380	HIS	-	expression tag	UNP E1BQ43
F	381	HIS	-	expression tag	UNP E1BQ43
F	382	HIS	-	expression tag	UNP E1BQ43
F	383	HIS	-	expression tag	UNP E1BQ43
F	384	HIS	-	expression tag	UNP E1BQ43

- Molecule 5 is GUANOSINE-5'-TRIPHOSPHATE (CCD ID: GTP) (formula: $C_{10}H_{16}N_5O_{14}P_3$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	A	1	Total	C	N	O	P	0	0
			32	10	5	14	3		
5	C	1	Total	C	N	O	P	0	0
			32	10	5	14	3		

- Molecule 6 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	1	Total	Mg	0	0
			1	1		
6	B	1	Total	Mg	0	0
			1	1		
6	C	1	Total	Mg	0	0
			1	1		
6	D	1	Total	Mg	0	0
			1	1		

- Molecule 7 is CALCIUM ION (CCD ID: CA) (formula: Ca).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	1	Total	Ca	0	0
			1	1		
7	B	2	Total	Ca	0	0
			2	2		
7	C	1	Total	Ca	0	0
			1	1		

- Molecule 8 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	A	1	Total	C	O	0	0
			6	3	3		
8	C	1	Total	C	O	0	0
			6	3	3		


- Molecule 9 is GUANOSINE-5'-DIPHOSPHATE (CCD ID: GDP) (formula: C₁₀H₁₅N₅O₁₁P₂).



- MES
-
- The chemical structure of MES (3-(3-sulfamoylpropyl)carbazole) is shown. It consists of a carbazole ring system (a benzene ring fused to a five-membered nitrogen-containing ring) connected via a propyl chain to a sulfamoyl group. The atoms are labeled: O1 (oxygen of the carbazole ring), C2, C3, C5, C6, C7, C8 (carbon atoms), N4 (nitrogen of the carbazole ring), S (sulfur of the sulfamoyl group), O2S, O1S (oxygens of the sulfamoyl group), and O3S (oxygen of the sulfamoyl group). The structure is drawn with green lines for the carbazole ring and blue lines for the propyl chain and sulfamoyl group.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
10	B	1	Total	C	N	O	S	0	0
			12	6	1	4	1		

- Molecule 11 is (3beta,4beta,5beta,10beta,11E,13E)-maytansine (CCD ID: 3GT) (formula: C₃₄H₄₆ClN₃O₁₀).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	Cl	N	O		
11	D	1	48	34	1	3	10	0	0

- Molecule 12 is PHOSPHOMETHYLPHOSPHONIC ACID ADENYLATE ESTER (CCD ID: ACP) (formula: C₁₁H₁₈N₅O₁₂P₃).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
12	F	1	31	11	5	12	3	0	0


- Molecule 13 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
13	A	134	Total 134	O 134	0	0
13	B	113	Total 113	O 113	0	0
13	C	252	Total 252	O 252	0	0
13	D	97	Total 97	O 97	0	0
13	E	30	Total 30	O 30	0	0
13	F	45	Total 45	O 45	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Tubulin alpha-1B chain




- Molecule 1: Tubulin alpha-1B chain




- Molecule 2: Tubulin beta-2B chain





- Molecule 2: Tubulin beta-2B chain







● Molecule 3: Stathmin-4



● Molecule 4: Tubulin-Tyrosine Ligase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	103.86Å 155.57Å 180.62Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	62.42 – 2.10 62.42 – 2.10	Depositor EDS
% Data completeness (in resolution range)	99.8 (62.42-2.10) 99.8 (62.42-2.10)	Depositor EDS
R_{merge}	0.14	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.99 (at 2.10Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.9_1692)	Depositor
R, R_{free}	0.189 , 0.218 0.197 , 0.223	Depositor DCC
R_{free} test set	8416 reflections (4.93%)	wwPDB-VP
Wilson B-factor (Å ²)	50.0	Xtriage
Anisotropy	0.183	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 50.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	18278	wwPDB-VP
Average B, all atoms (Å ²)	67.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.43% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MES, MG, 3GT, CA, ACP, GDP, GTP, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.29	0/3527	0.72	4/4786 (0.1%)
1	C	0.29	0/3594	0.70	0/4882
2	B	0.29	0/3451	0.68	0/4674
2	D	0.28	0/3408	0.70	0/4617
3	E	0.26	0/1017	0.64	0/1349
4	F	0.24	0/2830	0.65	0/3823
All	All	0.28	0/17827	0.69	4/24131 (0.0%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	262	TYR	CA-C-N	6.31	125.81	119.24
1	A	262	TYR	C-N-CA	6.31	125.81	119.24
1	A	174	ALA	CA-C-N	5.31	125.62	119.47
1	A	174	ALA	C-N-CA	5.31	125.62	119.47

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3433	0	3364	22	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	3483	0	3415	24	0
2	B	3365	0	3243	21	0
2	D	3326	0	3213	20	0
3	E	1006	0	1026	7	0
4	F	2763	0	2740	17	0
5	A	32	0	12	1	0
5	C	32	0	12	1	0
6	A	1	0	0	0	0
6	B	1	0	0	0	0
6	C	1	0	0	0	0
6	D	1	0	0	0	0
7	A	1	0	0	0	0
7	B	2	0	0	0	0
7	C	1	0	0	0	0
8	A	6	0	8	0	0
8	C	6	0	8	0	0
9	B	28	0	12	1	0
9	D	28	0	12	1	0
10	B	12	0	13	3	0
11	D	48	0	46	2	0
12	F	31	0	14	3	0
13	A	134	0	0	1	0
13	B	113	0	0	2	0
13	C	252	0	0	2	0
13	D	97	0	0	1	0
13	E	30	0	0	1	0
13	F	45	0	0	0	0
All	All	18278	0	17138	110	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 110 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:71:GLU:HB2	1:C:98:ASP:HB3	1.67	0.76
1:C:381:THR:HG22	1:C:383:ALA:H	1.53	0.74
1:C:270:ALA:HB3	1:C:302:MET:HE2	1.72	0.72
2:B:147[A]:SER:HG	2:B:190:SER:HG	1.41	0.69
1:A:18:ASN:HD21	1:A:78:VAL:HG13	1.60	0.66

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	439/451 (97%)	432 (98%)	7 (2%)	0	100	100
1	C	449/451 (100%)	441 (98%)	8 (2%)	0	100	100
2	B	427/445 (96%)	420 (98%)	7 (2%)	0	100	100
2	D	421/445 (95%)	412 (98%)	9 (2%)	0	100	100
3	E	118/143 (82%)	118 (100%)	0	0	100	100
4	F	325/384 (85%)	315 (97%)	10 (3%)	0	100	100
All	All	2179/2319 (94%)	2138 (98%)	41 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	374/379 (99%)	371 (99%)	3 (1%)	73	81
1	C	382/379 (101%)	380 (100%)	2 (0%)	81	88
2	B	370/383 (97%)	370 (100%)	0	100	100
2	D	368/383 (96%)	366 (100%)	2 (0%)	81	88
3	E	110/127 (87%)	107 (97%)	3 (3%)	39	45
4	F	304/342 (89%)	301 (99%)	3 (1%)	68	76

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
All	All	1908/1993 (96%)	1895 (99%)	13 (1%)	78	83

5 of 13 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	E	22	VAL
3	E	92	ASN
4	F	326	LYS
4	F	196	HIS
4	F	260	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 16 such sidechains are listed below:

Mol	Chain	Res	Type
4	F	260	ASN
4	F	45	ASN
1	C	216	ASN
3	E	92	ASN
2	B	436	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 17 ligands modelled in this entry, 8 are monoatomic - leaving 9 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The

Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
9	GDP	D	501	6	29,30,30	1.18	4 (13%)	45,47,47	1.74	5 (11%)
10	MES	B	505	-	12,12,12	2.28	1 (8%)	15,16,16	1.30	2 (13%)
9	GDP	B	501	6	29,30,30	1.20	3 (10%)	45,47,47	1.72	6 (13%)
11	3GT	D	502	-	49,51,51	1.68	8 (16%)	58,77,77	2.18	15 (25%)
5	GTP	C	501	6	33,34,34	1.22	4 (12%)	50,54,54	1.63	6 (12%)
12	ACP	F	401	-	31,33,33	1.73	9 (29%)	47,52,52	1.86	10 (21%)
8	GOL	C	503	-	5,5,5	0.37	0	5,5,5	0.32	0
8	GOL	A	504	-	5,5,5	0.36	0	5,5,5	0.31	0
5	GTP	A	501	6	33,34,34	1.24	5 (15%)	50,54,54	1.64	6 (12%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
9	GDP	D	501	6	-	3/16/32/32	0/3/3/3
10	MES	B	505	-	-	3/6/14/14	0/1/1/1
9	GDP	B	501	6	-	4/16/32/32	0/3/3/3
11	3GT	D	502	-	-	11/58/84/84	0/2/4/4
5	GTP	C	501	6	-	7/22/38/38	0/3/3/3
12	ACP	F	401	-	-	8/19/38/38	0/3/3/3
8	GOL	C	503	-	-	2/4/4/4	-
8	GOL	A	504	-	-	2/4/4/4	-
5	GTP	A	501	6	-	5/22/38/38	0/3/3/3

The worst 5 of 34 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
10	B	505	MES	C8-S	-7.68	1.66	1.77
11	D	502	3GT	OBB-CBC	6.65	1.45	1.35
11	D	502	3GT	OBB-C	5.30	1.46	1.34
12	F	401	ACP	C5-C4	4.70	1.47	1.39
11	D	502	3GT	CBI-CBH	4.24	1.52	1.47

The worst 5 of 50 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
11	D	502	3GT	OBK-CBH-CBF	-6.56	109.55	117.41
11	D	502	3GT	CBI-OBK-CBH	6.37	64.42	60.81
9	B	501	GDP	C5-C4-N3	-5.98	118.88	128.39
9	D	501	GDP	C5-C4-N3	-5.95	118.92	128.39
12	F	401	ACP	C5-C4-N3	-5.92	118.57	126.72

There are no chirality outliers.

5 of 45 torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
5	A	501	GTP	PB-O3B-PG-O2G
5	A	501	GTP	C5'-O5'-PA-O3A
5	A	501	GTP	C5'-O5'-PA-O1A
5	A	501	GTP	C5'-O5'-PA-O2A
5	C	501	GTP	C5'-O5'-PA-O3A

There are no ring outliers.


7 monomers are involved in 12 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
9	D	501	GDP	1	0
10	B	505	MES	3	0
9	B	501	GDP	1	0
11	D	502	3GT	2	0
5	C	501	GTP	1	0
12	F	401	ACP	3	0
5	A	501	GTP	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.




Ligand 3GT D 502




Bond lengths





Bond angles




Torsions



Rings





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	436/451 (96%)	0.23	8 (1%) 67 70	32, 60, 95, 133	7 (1%)
1	C	440/451 (97%)	0.00	7 (1%) 70 73	26, 48, 76, 111	11 (2%)
2	B	427/445 (95%)	0.27	19 (4%) 39 41	27, 56, 96, 139	6 (1%)
2	D	422/445 (94%)	0.42	13 (3%) 51 54	34, 67, 100, 139	6 (1%)
3	E	121/143 (84%)	0.65	11 (9%) 15 15	32, 73, 111, 128	1 (0%)
4	F	335/384 (87%)	0.73	32 (9%) 13 14	34, 85, 161, 181	2 (0%)
All	All	2181/2319 (94%)	0.33	90 (4%) 41 43	26, 62, 113, 181	33 (1%)

The worst 5 of 90 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	278	ARG	4.6
4	F	161	LEU	4.6
4	F	149	ALA	4.6
2	B	248	LEU	4.5
1	A	281	ALA	4.5

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands ⓘ




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
8	GOL	A	504	6/6	0.68	0.20	86,87,90,92	0
7	CA	B	504	1/1	0.79	0.22	115,115,115,115	0
12	ACP	F	401	31/31	0.82	0.14	87,97,167,172	0
11	3GT	D	502	48/48	0.85	0.16	80,89,105,188	0
9	GDP	D	501	28/28	0.91	0.11	56,64,74,85	0
10	MES	B	505	12/12	0.91	0.11	57,65,77,82	0
8	GOL	C	503	6/6	0.92	0.18	77,79,82,82	0
6	MG	B	502	1/1	0.93	0.12	39,39,39,39	0
7	CA	A	503	1/1	0.94	0.07	85,85,85,85	0
7	CA	B	503	1/1	0.95	0.06	101,101,101,101	0
6	MG	D	503	1/1	0.97	0.08	62,62,62,62	0
9	GDP	B	501	28/28	0.97	0.06	32,43,47,50	0
5	GTP	A	501	32/32	0.98	0.06	37,44,49,74	0
5	GTP	C	501	32/32	0.98	0.05	30,37,41,46	0
7	CA	C	504	1/1	0.99	0.03	67,67,67,67	0
6	MG	A	502	1/1	0.99	0.02	44,44,44,44	0
6	MG	C	502	1/1	1.00	0.05	41,41,41,41	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around ACP F 401:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around 3GT D 502:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around GDP D 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around GDP B 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around GTP A 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around GTP C 501:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.