



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 1, 2026 – 06:57 PM UTC

PDB ID : 4TVT / pdb_00004tv
Title : New ligand for thaumatin discovered using acoustic high throughput screening
Authors : Teplitsky, E.; Joshi, K.; Ericson, D.L.; Scalia, A.; Mullen, J.D.; Sweet, R.M.;
Soares, A.S.
Deposited on : 2014-06-28
Resolution : 1.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1216 (1.20-1.20)
Clashscore	190562	1265 (1.20-1.20)
Ramachandran outliers	187476	1226 (1.20-1.20)
Sidechain outliers	187428	1226 (1.20-1.20)
RSRZ outliers	180081	1214 (1.20-1.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	207	5% 74% 21% .

2 Entry composition [i](#)


There are 5 unique types of molecules in this entry. The entry contains 2124 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Thaumatin-1.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	207	1604	995	284	306	19	0	6	0

- Molecule 2 is ASCORBIC ACID (CCD ID: ASC) (formula: C₆H₈O₆).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			12	6	6		
2	A	1	Total	C	O	0	0
			12	6	6		
2	A	1	Total	C	O	0	0
			12	6	6		
2	A	1	Total	C	O	0	0
			12	6	6		
2	A	1	Total	C	O	0	0
			12	6	6		

- Molecule 3 is L(+)-TARTARIC ACID (CCD ID: TLA) (formula: $C_4H_6O_6$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			10	4	6		

- Molecule 4 is SODIUM ION (CCD ID: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	8	Total	Na	0	0
			8	8		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	442	Total	O	0	0
			442	442		

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Thaumatin-1



4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, α , β , γ	57.84Å 57.84Å 149.58Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	53.95 – 1.20 53.95 – 1.20	Depositor EDS
% Data completeness (in resolution range)	89.8 (53.95-1.20) 89.9 (53.95-1.20)	Depositor EDS
R_{merge}	0.03	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.02 (at 1.20Å)	Xtriage
Refinement program	REFMAC 5.8.0069	Depositor
R, R_{free}	0.145 , 0.163 0.144 , 0.162	Depositor DCC
R_{free} test set	3518 reflections (4.39%)	wwPDB-VP
Wilson B-factor (Å ²)	12.6	Xtriage
Anisotropy	0.371	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.40 , 57.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.98	EDS
Total number of atoms	2124	wwPDB-VP
Average B, all atoms (Å ²)	19.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.42% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NA, ASC, TLA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.96	45/1640 (2.7%)	1.49	14/2218 (0.6%)

All (45) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	127	ALA	C-N	-8.25	1.22	1.33
1	A	95	TYR	CA-C	-7.46	1.43	1.53
1	A	37	TRP	NE1-CE2	-7.30	1.29	1.37
1	A	32	ASN	CG-ND2	-7.27	1.18	1.33
1	A	67	LYS	CA-C	-7.08	1.43	1.52
1	A	200	ARG	NE-CZ	-6.91	1.25	1.33
1	A	143	GLY	C-O	6.90	1.33	1.23
1	A	109	ASN	CG-OD1	6.89	1.36	1.23
1	A	185	LEU	CA-C	-6.47	1.43	1.52
1	A	10	SER	CA-C	-6.40	1.44	1.52
1	A	82	ARG	CZ-NH2	-6.39	1.25	1.33
1	A	68	THR	N-CA	-6.36	1.38	1.46
1	A	161	THR	N-CA	6.33	1.54	1.46
1	A	4	GLU	CD-OE1	-6.32	1.13	1.25
1	A	64	GLY	N-CA	-6.31	1.36	1.45
1	A	37	TRP	CA-CB	-6.27	1.44	1.53
1	A	63	SER	CA-C	-6.18	1.45	1.52
1	A	179	ASP	CG-OD1	6.12	1.36	1.25
1	A	23	ALA	CA-C	-6.02	1.45	1.52
1	A	97	LYS	N-CA	5.96	1.53	1.45
1	A	109	ASN	CG-ND2	-5.91	1.20	1.33
1	A	67	LYS	CA-CB	-5.84	1.43	1.53
1	A	204	CYS	CA-C	-5.73	1.45	1.52
1	A	57	TYR	CB-CG	-5.71	1.39	1.51
1	A	160	THR	CB-OG1	-5.70	1.34	1.43
1	A	78	LYS	CA-C	-5.62	1.45	1.52

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	38	THR	CA-CB	-5.62	1.45	1.53
1	A	69	GLY	N-CA	-5.60	1.37	1.45
1	A	133	GLN	CA-CB	-5.60	1.43	1.53
1	A	125	ARG	N-CA	-5.58	1.39	1.46
1	A	187	LYS	CA-CB	-5.53	1.43	1.53
1	A	186	ASP	N-CA	-5.38	1.39	1.46
1	A	202	THR	C-O	5.35	1.30	1.24
1	A	124	VAL	CB-CG2	-5.33	1.34	1.52
1	A	184	VAL	CA-C	-5.32	1.46	1.52
1	A	184	VAL	C-O	5.29	1.30	1.24
1	A	120	GLY	N-CA	-5.29	1.38	1.45
1	A	171	ARG	CZ-NH1	-5.27	1.25	1.32
1	A	153	GLN	CG-CD	-5.24	1.39	1.52
1	A	141	PRO	C-O	5.14	1.30	1.24
1	A	155	SER	N-CA	-5.11	1.40	1.46
1	A	200	ARG	CZ-NH2	5.08	1.40	1.33
1	A	135	PRO	CA-C	-5.03	1.46	1.52
1	A	125	ARG	NE-CZ	-5.01	1.27	1.33
1	A	14	TRP	C-O	5.01	1.29	1.23

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	200	ARG	NE-CZ-NH2	-8.85	111.23	119.20
1	A	175	ARG	NE-CZ-NH2	6.93	125.44	119.20
1	A	67	LYS	O-C-N	-6.33	114.48	122.27
1	A	200	ARG	NE-CZ-NH1	6.22	127.72	121.50
1	A	153	GLN	OE1-CD-NE2	-6.12	116.48	122.60
1	A	184	VAL	CA-C-O	-5.93	114.78	120.95
1	A	185	LEU	CA-C-O	5.58	126.42	119.78
1	A	141	PRO	O-C-N	5.54	129.36	122.22
1	A	160	THR	N-CA-CB	-5.30	102.09	110.25
1	A	32	ASN	OD1-CG-ND2	-5.25	117.35	122.60
1	A	23	ALA	O-C-N	-5.11	116.53	122.81
1	A	42	GLU	CB-CG-CD	5.07	121.23	112.60
1	A	141	PRO	CA-C-N	5.04	126.44	120.14
1	A	141	PRO	C-N-CA	5.04	126.44	120.14

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1604	0	1531	21	0
2	A	60	0	35	3	0
3	A	10	0	4	0	0
4	A	8	0	0	0	0
5	A	442	0	0	16	1
All	All	2124	0	1570	24	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (24) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:304:ASC:H61	5:A:403:HOH:O	1.25	1.33
1:A:159[B]:CYS:SG	5:A:798:HOH:O	2.17	1.00
2:A:304:ASC:C6	5:A:403:HOH:O	1.90	0.99
1:A:179:ASP:HB3	5:A:801:HOH:O	1.71	0.91
1:A:179:ASP:CB	5:A:801:HOH:O	2.17	0.90
1:A:160:THR:HG21	5:A:468:HOH:O	1.73	0.88
1:A:179:ASP:CG	5:A:801:HOH:O	2.21	0.82
1:A:122[A]:ARG:NH1	5:A:629:HOH:O	2.11	0.80
1:A:160:THR:CG2	5:A:468:HOH:O	2.28	0.78
1:A:44:GLY:H	1:A:93:ASN:HD22	1.33	0.77
1:A:30:GLN:HE21	1:A:32:ASN:HD21	1.35	0.74
1:A:55:ASP:OD1	5:A:404:HOH:O	2.16	0.61
1:A:30:GLN:HE21	1:A:32:ASN:ND2	1.99	0.60
1:A:122[A]:ARG:NH2	1:A:124:VAL:HG12	2.19	0.57
1:A:163:LYS:HE2	5:A:466:HOH:O	2.08	0.53
2:A:303:ASC:H4	5:A:804:HOH:O	2.04	0.52
1:A:163:LYS:CE	5:A:466:HOH:O	2.60	0.50
1:A:109:ASN:HD22	1:A:109:ASN:H	1.65	0.45
1:A:160:THR:CB	5:A:468:HOH:O	2.66	0.42
1:A:122[C]:ARG:NH2	5:A:653:HOH:O	2.19	0.42
1:A:31[A]:LEU:HD21	1:A:37:TRP:HB2	2.02	0.41
1:A:119:ARG:NH1	5:A:409:HOH:O	2.52	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:86:THR:H	1:A:109:ASN:ND2	2.19	0.41
1:A:78:LYS:HB2	1:A:78:LYS:HE2	1.89	0.41

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:A:555:HOH:O	5:A:555:HOH:O[7_645]	2.15	0.05

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	211/207 (102%)	209 (99%)	2 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	173/167 (104%)	164 (95%)	9 (5%)	21	2

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	76[A]	ARG
1	A	76[B]	ARG
1	A	109	ASN
1	A	119	ARG
1	A	122[A]	ARG
1	A	122[C]	ARG
1	A	159[A]	CYS
1	A	159[B]	CYS
1	A	176	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	30	GLN
1	A	32	ASN
1	A	93	ASN
1	A	104	ASN
1	A	109	ASN
1	A	146	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 14 ligands modelled in this entry, 8 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	ASC	A	302	-	12,12,12	1.74	4 (33%)	17,17,17	2.40	6 (35%)
3	TLA	A	306	-	9,9,9	2.05	3 (33%)	12,12,12	1.86	3 (25%)
2	ASC	A	303	-	12,12,12	1.21	1 (8%)	17,17,17	1.99	5 (29%)
2	ASC	A	305	-	12,12,12	1.11	1 (8%)	17,17,17	1.62	4 (23%)
2	ASC	A	304	-	12,12,12	2.16	5 (41%)	17,17,17	1.96	6 (35%)
2	ASC	A	301	-	12,12,12	1.61	2 (16%)	17,17,17	1.57	4 (23%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ASC	A	302	-	-	2/6/22/22	0/1/1/1
3	TLA	A	306	-	-	0/12/12/12	-
2	ASC	A	303	-	-	0/6/22/22	0/1/1/1
2	ASC	A	305	-	-	0/6/22/22	0/1/1/1
2	ASC	A	304	-	-	0/6/22/22	0/1/1/1
2	ASC	A	301	-	-	0/6/22/22	0/1/1/1

All (16) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	304	ASC	C5-C4	-4.05	1.45	1.53
2	A	301	ASC	O1-C1	3.88	1.29	1.21
3	A	306	TLA	O3-C3	3.69	1.49	1.42
3	A	306	TLA	O4-C4	3.18	1.31	1.22
2	A	302	ASC	C5-C4	3.15	1.58	1.53
2	A	304	ASC	C2-C3	-3.07	1.27	1.35
3	A	306	TLA	C3-C4	-3.03	1.48	1.52
2	A	304	ASC	O4-C4	-2.97	1.40	1.45
2	A	304	ASC	O1-C1	2.93	1.27	1.21
2	A	302	ASC	O4-C4	-2.91	1.40	1.45
2	A	303	ASC	O5-C5	-2.48	1.38	1.43
2	A	302	ASC	O4-C1	2.42	1.39	1.36
2	A	301	ASC	O2-C2	2.28	1.41	1.33
2	A	305	ASC	C6-C5	2.26	1.57	1.52

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	302	ASC	C6-C5	2.14	1.57	1.52
2	A	304	ASC	O5-C5	2.12	1.47	1.43

All (28) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	302	ASC	C1-C2-C3	-5.98	100.44	107.77
2	A	304	ASC	O1-C1-C2	-4.41	122.72	129.33
2	A	303	ASC	O4-C1-C2	-4.10	106.23	109.86
3	A	306	TLA	O4-C4-C3	-3.99	111.00	121.62
2	A	304	ASC	C4-O4-C1	-3.97	104.92	109.27
2	A	302	ASC	C4-C3-C2	3.94	115.56	109.55
2	A	302	ASC	O1-C1-C2	-3.70	123.79	129.33
2	A	301	ASC	O4-C1-O1	-3.63	116.49	121.27
2	A	303	ASC	C4-O4-C1	3.53	113.14	109.27
2	A	305	ASC	O4-C1-O1	-3.38	116.81	121.27
2	A	303	ASC	O6-C6-C5	-3.29	104.25	111.16
3	A	306	TLA	O41-C4-C3	3.24	122.30	113.31
2	A	304	ASC	O4-C1-O1	3.06	125.31	121.27
2	A	305	ASC	O5-C5-C4	2.97	116.36	110.74
2	A	301	ASC	O1-C1-C2	2.80	133.52	129.33
3	A	306	TLA	O11-C1-O1	2.69	130.19	124.08
2	A	302	ASC	O4-C1-O1	2.67	124.80	121.27
2	A	303	ASC	O4-C4-C3	-2.52	101.23	103.76
2	A	304	ASC	O4-C4-C3	2.49	106.25	103.76
2	A	301	ASC	O6-C6-C5	-2.44	106.02	111.16
2	A	305	ASC	O3-C3-C2	2.35	138.41	132.40
2	A	304	ASC	O4-C1-C2	2.33	111.92	109.86
2	A	302	ASC	O4-C4-C3	-2.24	101.51	103.76
2	A	305	ASC	O3-C3-C4	-2.17	112.45	117.99
2	A	303	ASC	O5-C5-C4	2.17	114.85	110.74
2	A	302	ASC	O5-C5-C4	-2.16	106.64	110.74
2	A	301	ASC	C4-C3-C2	2.14	112.81	109.55
2	A	304	ASC	C5-C4-C3	-2.07	110.91	114.71

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	302	ASC	C4-C5-C6-O6
2	A	302	ASC	O5-C5-C6-O6

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	303	ASC	1	0
2	A	304	ASC	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	207/207 (100%)	-0.05	10 (4%)	35 37	5, 12, 27, 61	6 (2%)

All (10) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	207	ALA	4.6
1	A	60	ASP	2.9
1	A	142	GLY	2.9
1	A	119	ARG	2.3
1	A	206	THR	2.3
1	A	57	TYR	2.3
1	A	143	GLY	2.1
1	A	160	THR	2.1
1	A	161	THR	2.1
1	A	163	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	ASC	A	305	12/12	0.89	0.12	12,14,17,18	12
2	ASC	A	304	12/12	0.90	0.12	12,15,21,21	12
4	NA	A	310	1/1	0.90	0.11	23,23,23,23	0
4	NA	A	308	1/1	0.93	0.10	34,34,34,34	0
2	ASC	A	302	12/12	0.93	0.10	14,18,27,34	12
4	NA	A	307	1/1	0.94	0.09	35,35,35,35	0
4	NA	A	312	1/1	0.95	0.08	23,23,23,23	0
4	NA	A	311	1/1	0.96	0.07	18,18,18,18	0
2	ASC	A	303	12/12	0.96	0.08	12,18,21,22	12
2	ASC	A	301	12/12	0.97	0.06	11,15,17,17	0
4	NA	A	314	1/1	0.97	0.08	20,20,20,20	0
3	TLA	A	306	10/10	0.98	0.05	11,13,17,18	0
4	NA	A	313	1/1	0.99	0.04	14,14,14,14	0
4	NA	A	309	1/1	0.99	0.06	19,19,19,19	0

6.5 Other polymers [i](#)

There are no such residues in this entry.