



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 8, 2026 – 05:25 PM UTC

PDB ID : 5TW1 / pdb_00005tw1
Title : Crystal structure of a Mycobacterium smegmatis transcription initiation complex with RbpA
Authors : Hubin, E.A.; Darst, S.A.; Campbell, E.A.
Deposited on : 2016-11-10
Resolution : 2.76 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION




The reported resolution of this entry is 2.76 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1009 (2.76-2.76)
Clashscore	190562	1044 (2.76-2.76)
Ramachandran outliers	187476	1024 (2.76-2.76)
Sidechain outliers	187428	1024 (2.76-2.76)
RSRZ outliers	180081	1009 (2.76-2.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	J	114	
2	A	350	
2	B	350	
2	T	350	
3	C	1169	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
4	D	1317	
5	E	107	
6	F	466	
7	O	31	
8	P	26	
9	G	17	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
10	SO4	C	1203	-	-	X	-
10	SO4	D	2005	-	-	X	-

2 Entry composition

There are 14 unique types of molecules in this entry. The entry contains 26644 atoms, of which 36 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called RNA polymerase-binding protein RbpA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	J	83	Total	C	N	O	S	0	0	0
			667	419	118	128	2			

- Molecule 2 is a protein called DNA-directed RNA polymerase subunit alpha.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	A	218	Total	C	N	O	S	0	0	0
			1617	1020	276	318	3			
2	B	233	Total	C	N	O	S	0	0	0
			1667	1054	289	322	2			
2	T	53	Total	C	N	O	S	0	0	0
			374	236	65	72	1			

- Molecule 3 is a protein called DNA-directed RNA polymerase subunit beta.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	1099	Total	C	N	O	S	0	0	0
			8250	5164	1448	1603	35			

- Molecule 4 is a protein called DNA-directed RNA polymerase subunit beta'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	D	1248	Total	C	N	O	S	0	0	0
			9588	6016	1727	1805	40			

- Molecule 5 is a protein called DNA-directed RNA polymerase subunit omega.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
5	E	76	Total	C	N	O		0	0	0
			592	378	100	114				

- Molecule 6 is a protein called RNA polymerase sigma factor SigA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
6	F	302	Total	C	N	O	S	0	0	0
			2396	1502	433	454	7			

- Molecule 7 is a DNA chain called DNA (31-MER).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
7	O	31	Total	C	N	O	P	0	0	0
			634	305	114	185	30			


- Molecule 8 is a DNA chain called DNA (26-MER).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
8	P	26	Total	C	N	O	P	0	0	0
			526	254	94	153	25			

- Molecule 9 is a protein called Unknown peptide.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
9	G	17	Total	C	N	O	0	0	0
			85	51	17	17			

- Molecule 10 is SULFATE ION (CCD ID: SO4) (formula: O₄S).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
10	C	1	Total	O	S	0	0
			5	4	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
10	C	1	Total	O	S	0	0
			5	4	1		
10	C	1	Total	O	S	0	0
			5	4	1		
10	D	1	Total	O	S	0	0
			5	4	1		
10	D	1	Total	O	S	0	0
			5	4	1		
10	D	1	Total	O	S	0	0
			5	4	1		
10	D	1	Total	O	S	0	0
			5	4	1		
10	F	1	Total	O	S	0	0
			5	4	1		
10	F	1	Total	O	S	0	0
			5	4	1		
10	F	1	Total	O	S	0	0
			5	4	1		
10	F	1	Total	O	S	0	0
			5	4	1		

- Molecule 11 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
11	C	1	Total C H O 10 2 6 2	0	0
11	C	1	Total C H O 10 2 6 2	0	0
11	D	1	Total C H O 10 2 6 2	0	0
11	D	1	Total C H O 10 2 6 2	0	0
11	D	1	Total C H O 10 2 6 2	0	0
11	F	1	Total C H O 10 2 6 2	0	0

- Molecule 12 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
12	D	2	Total Zn 2 2	0	0

- Molecule 13 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
13	D	1	Total Mg 1 1	0	0


- Molecule 14 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
14	J	2	Total O 2 2	0	0
14	A	2	Total O 2 2	0	0
14	B	2	Total O 2 2	0	0
14	C	30	Total O 30 30	0	0
14	D	57	Total O 57 57	0	0
14	E	3	Total O 3 3	0	0
14	F	16	Total O 16 16	0	0

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
14	O	10	Total	O	0	0
			10	10		
14	P	3	Total	O	0	0
			3	3		










- Molecule 8: DNA (26-MER)



- Molecule 9: Unknown peptide



There are no outlier residues recorded for this chain.

4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	133.01Å 161.63Å 139.21Å 90.00° 107.72° 90.00°	Depositor
Resolution (Å)	54.91 – 2.76 54.91 – 2.76	Depositor EDS
% Data completeness (in resolution range)	99.5 (54.91-2.76) 99.5 (54.91-2.76)	Depositor EDS
R_{merge}	0.23	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.16 (at 2.77Å)	Xtriage
Refinement program	PHENIX (1.10.1_2155)	Depositor
R, R_{free}	0.239 , 0.280 (Not available) , 0.249	Depositor DCC
R_{free} test set	1989 reflections (1.37%)	wwPDB-VP
Wilson B-factor (Å ²)	78.4	Xtriage
Anisotropy	0.150	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 47.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.35$	Xtriage
Estimated twinning fraction	0.003 for l,-k,h	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	26644	wwPDB-VP
Average B, all atoms (Å ²)	85.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.46% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, SO4, MG, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	J	0.19	0/681	0.44	0/923
2	A	0.19	0/1641	0.43	0/2236
2	B	0.18	0/1693	0.47	0/2316
2	T	0.12	0/376	0.31	0/511
3	C	0.20	0/8394	0.46	2/11410 (0.0%)
4	D	0.22	0/9742	0.45	1/13189 (0.0%)
5	E	0.21	0/604	0.48	0/822
6	F	0.17	0/2426	0.37	0/3273
7	O	0.28	0/710	0.48	0/1095
8	P	0.29	0/589	0.49	0/906
All	All	0.21	0/26856	0.45	3/36681 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	J	0	1
3	C	0	5
All	All	0	6

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	D	901	ALA	CB-CA-C	-9.54	104.44	115.79
3	C	949	LYS	N-CA-C	7.11	122.13	113.17
3	C	1132	ASP	N-CA-C	-6.81	99.42	108.34

There are no chirality outliers.

5 of 6 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
3	C	368	ARG	Peptide
3	C	433	GLN	Peptide
3	C	982	SER	Peptide
3	C	985	PRO	Peptide
1	J	70	PRO	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	J	667	0	649	55	0
2	A	1617	0	1636	182	0
2	B	1667	0	1636	194	0
2	T	374	0	344	29	0
3	C	8250	0	7989	712	0
4	D	9588	0	9552	651	0
5	E	592	0	583	53	0
6	F	2396	0	2422	146	0
7	O	634	0	350	33	0
8	P	526	0	296	16	0
9	G	85	0	19	0	0
10	C	15	0	0	2	0
10	D	20	0	0	2	0
10	F	25	0	0	1	0
11	C	8	12	12	1	0
11	D	12	18	18	3	0
11	F	4	6	6	1	0
12	D	2	0	0	0	0
13	D	1	0	0	0	0
14	A	2	0	0	0	0
14	B	2	0	0	0	0
14	C	30	0	0	16	0
14	D	57	0	0	17	0
14	E	3	0	0	0	0
14	F	16	0	0	3	0
14	J	2	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
14	O	10	0	0	3	0
14	P	3	0	0	1	0
All	All	26608	36	25512	1898	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 36.

The worst 5 of 1898 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
6:F:253:MET:HE3	6:F:297:MET:HA	1.24	1.15
2:B:184:GLU:HA	2:B:187:THR:HG22	1.28	1.15
6:F:253:MET:HE2	6:F:300:GLN:HB2	1.26	1.15
3:C:940:ALA:HB1	3:C:941:ALA:HA	1.28	1.12
4:D:527:LEU:HD21	4:D:581:MET:HE1	1.31	1.12

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	J	81/114 (71%)	75 (93%)	6 (7%)	0	100	100
2	A	214/350 (61%)	202 (94%)	12 (6%)	0	100	100
2	B	231/350 (66%)	216 (94%)	14 (6%)	1 (0%)	30	50
2	T	33/350 (9%)	33 (100%)	0	0	100	100
3	C	1093/1169 (94%)	1038 (95%)	51 (5%)	4 (0%)	30	50
4	D	1238/1317 (94%)	1189 (96%)	46 (4%)	3 (0%)	43	64
5	E	72/107 (67%)	65 (90%)	5 (7%)	2 (3%)	4	6
6	F	298/466 (64%)	294 (99%)	4 (1%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
All	All	3260/4223 (77%)	3112 (96%)	138 (4%)	10 (0%)	36	56

5 of 10 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
5	E	76	VAL
3	C	982	SER
3	C	1134	ALA
4	D	902	GLU
4	D	1194	VAL

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	J	71/98 (72%)	62 (87%)	9 (13%)	4	8
2	A	178/297 (60%)	159 (89%)	19 (11%)	6	12
2	B	171/297 (58%)	150 (88%)	21 (12%)	4	8
2	T	35/297 (12%)	29 (83%)	6 (17%)	2	3
3	C	857/984 (87%)	762 (89%)	95 (11%)	6	11
4	D	994/1095 (91%)	894 (90%)	100 (10%)	7	14
5	E	62/86 (72%)	51 (82%)	11 (18%)	2	3
6	F	252/379 (66%)	227 (90%)	25 (10%)	7	15
All	All	2620/3533 (74%)	2334 (89%)	286 (11%)	6	12

5 of 286 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
4	D	1086	ARG
4	D	1215	SER
6	F	291	GLN
3	C	703	GLU
3	C	636	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 49 such sidechains are listed below:

Mol	Chain	Res	Type
4	D	564	ASN
4	D	778	GLN
4	D	600	GLN
4	D	748	HIS
4	D	853	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 21 ligands modelled in this entry, 3 are monoatomic - leaving 18 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
10	SO4	F	505	-	4,4,4	0.24	0	6,6,6	0.15	0
11	EDO	F	506	-	3,3,3	0.45	0	2,2,2	0.29	0
10	SO4	F	501	-	4,4,4	0.25	0	6,6,6	0.09	0
10	SO4	D	2007	-	4,4,4	0.23	0	6,6,6	0.14	0
11	EDO	D	2009	-	3,3,3	0.42	0	2,2,2	0.35	0
10	SO4	C	1202	-	4,4,4	0.23	0	6,6,6	0.16	0
10	SO4	F	504	-	4,4,4	0.25	0	6,6,6	0.13	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
11	EDO	C	1205	-	3,3,3	0.48	0	2,2,2	0.27	0
11	EDO	C	1204	-	3,3,3	0.40	0	2,2,2	0.45	0
11	EDO	D	2008	-	3,3,3	0.45	0	2,2,2	0.29	0
11	EDO	D	2010	-	3,3,3	0.47	0	2,2,2	0.18	0
10	SO4	D	2004	-	4,4,4	0.24	0	6,6,6	0.30	0
10	SO4	F	502	-	4,4,4	0.23	0	6,6,6	0.25	0
10	SO4	F	503	-	4,4,4	0.24	0	6,6,6	0.16	0
10	SO4	C	1203	-	4,4,4	0.22	0	6,6,6	0.17	0
10	SO4	C	1201	-	4,4,4	0.23	0	6,6,6	0.13	0
10	SO4	D	2005	-	4,4,4	0.20	0	6,6,6	0.15	0
10	SO4	D	2006	-	4,4,4	0.23	0	6,6,6	0.25	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
11	EDO	F	506	-	-	1/1/1/1	-
11	EDO	D	2009	-	-	0/1/1/1	-
11	EDO	C	1205	-	-	1/1/1/1	-
11	EDO	C	1204	-	-	1/1/1/1	-
11	EDO	D	2008	-	-	1/1/1/1	-
11	EDO	D	2010	-	-	0/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
11	C	1204	EDO	O1-C1-C2-O2
11	D	2008	EDO	O1-C1-C2-O2
11	C	1205	EDO	O1-C1-C2-O2
11	F	506	EDO	O1-C1-C2-O2

There are no ring outliers.

6 monomers are involved in 10 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
11	F	506	EDO	1	0
11	C	1205	EDO	1	0
11	D	2008	EDO	3	0
10	F	502	SO4	1	0
10	C	1203	SO4	2	0
10	D	2005	SO4	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	J	83/114 (72%)	0.48	3 (3%) 46 44	64, 97, 141, 158	0
2	A	218/350 (62%)	0.54	12 (5%) 30 30	59, 86, 118, 137	0
2	B	233/350 (66%)	1.00	31 (13%) 7 6	80, 110, 134, 148	0
2	T	53/350 (15%)	1.35	15 (28%) 1 1	113, 149, 165, 169	0
3	C	1099/1169 (94%)	0.47	53 (4%) 35 35	42, 85, 147, 163	0
4	D	1248/1317 (94%)	0.20	30 (2%) 59 57	39, 73, 125, 156	0
5	E	76/107 (71%)	0.28	2 (2%) 57 55	51, 78, 113, 117	0
6	F	302/466 (64%)	0.00	0 100 100	38, 72, 122, 137	0
7	O	31/31 (100%)	-0.50	0 100 100	51, 64, 83, 89	0
8	P	26/26 (100%)	-0.44	0 100 100	59, 68, 83, 93	0
9	G	0/17	-	-	-	-
All	All	3369/4297 (78%)	0.36	146 (4%) 40 38	38, 82, 141, 169	0

The worst 5 of 146 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	155	SER	5.5
2	T	275	LEU	5.4
3	C	228	LEU	4.8
3	C	229	LEU	4.4
2	A	2	LEU	4.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
10	SO4	D	2007	5/5	0.68	0.09	93,98,113,141	0
10	SO4	F	504	5/5	0.73	0.07	112,114,117,128	0
10	SO4	C	1203	5/5	0.76	0.08	103,113,139,197	0
13	MG	D	2003	1/1	0.76	0.28	82,82,82,82	0
10	SO4	C	1202	5/5	0.77	0.08	105,111,115,178	0
10	SO4	C	1201	5/5	0.79	0.07	111,114,124,138	0
11	EDO	D	2009	4/4	0.80	0.12	71,86,103,103	0
11	EDO	C	1205	4/4	0.81	0.15	74,93,101,112	0
11	EDO	F	506	4/4	0.82	0.12	73,88,95,95	0
11	EDO	C	1204	4/4	0.84	0.11	53,63,75,77	0
11	EDO	D	2008	4/4	0.86	0.10	74,89,105,105	0
10	SO4	F	501	5/5	0.88	0.09	97,98,116,122	0
10	SO4	D	2006	5/5	0.88	0.09	80,103,121,126	0
10	SO4	F	505	5/5	0.88	0.11	80,84,117,167	0
10	SO4	D	2004	5/5	0.89	0.11	69,72,83,87	0
10	SO4	F	502	5/5	0.89	0.09	84,86,102,108	0
10	SO4	F	503	5/5	0.90	0.10	85,92,98,113	0
11	EDO	D	2010	4/4	0.92	0.20	62,75,77,89	0
10	SO4	D	2005	5/5	0.93	0.14	80,85,97,125	0
12	ZN	D	2002	1/1	0.97	0.12	222,222,222,222	0
12	ZN	D	2001	1/1	0.99	0.04	66,66,66,66	0

6.5 Other polymers [i](#)

There are no such residues in this entry.