



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 03:19 AM UTC

PDB ID : 6TWU / pdb_00006twu
Title : MAGI1_2 complexed with a phosphomimetic 16E6 peptide
Authors : Gogl, G.; Cousido-Siah, A.; Trave, G.
Deposited on : 2020-01-13
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4912 (2.40-2.40)
Clashscore	190562	5391 (2.40-2.40)
Ramachandran outliers	187476	5320 (2.40-2.40)
Sidechain outliers	187428	5321 (2.40-2.40)
RSRZ outliers	180081	4916 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	427	5% 83% 15% ..
1	B	427	2% 86% 11% ..
2	C	10	30% 70%

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 6821 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Membrane-associated guanylate kinase, WW and PDZ domain-containing protein 1,Annexin A2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	421	Total	C	N	O	S	0	4	0
			3338	2097	571	656	14			
1	B	421	Total	C	N	O	S	0	1	0
			3326	2091	571	650	14			

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	452	GLY	-	expression tag	UNP Q96QZ7
A	453	SER	-	expression tag	UNP Q96QZ7
A	454	MET	-	expression tag	UNP Q96QZ7
A	455	GLY	-	expression tag	UNP Q96QZ7
A	456	LYS	-	expression tag	UNP Q96QZ7
A	457	PRO	-	expression tag	UNP Q96QZ7
A	559	GLY	-	linker	UNP Q96QZ7
A	560	SER	-	linker	UNP Q96QZ7
A	605	GLU	ALA	conflict	UNP P07355
B	452	GLY	-	expression tag	UNP Q96QZ7
B	453	SER	-	expression tag	UNP Q96QZ7
B	454	MET	-	expression tag	UNP Q96QZ7
B	455	GLY	-	expression tag	UNP Q96QZ7
B	456	LYS	-	expression tag	UNP Q96QZ7
B	457	PRO	-	expression tag	UNP Q96QZ7
B	559	GLY	-	linker	UNP Q96QZ7
B	560	SER	-	linker	UNP Q96QZ7
B	605	GLU	ALA	conflict	UNP P07355


- Molecule 2 is a protein called Protein E6.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	C	3	Total	C	N	O	0	0	0
			27	16	4	7			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	156	GLU	THR	conflict	UNP P03126

- Molecule 3 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is CITRIC ACID (CCD ID: CIT) (formula: $C_6H_8O_7$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			13	6	7		
4	B	1	Total	C	O	0	0
			13	6	7		
4	B	1	Total	C	O	0	0
			13	6	7		

- Molecule 5 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	5	Total	Ca	0	0
			5	5		
5	B	6	Total	Ca	0	0
			6	6		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	30	Total	O	0	0
			30	30		
6	B	32	Total	O	0	0
			32	32		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Membrane-associated guanylate kinase, WW and PDZ domain-containing protein 1, Annexin A2



- Molecule 1: Membrane-associated guanylate kinase, WW and PDZ domain-containing protein 1, Annexin A2



- Molecule 2: Protein E6



4 Data and refinement statistics

Property	Value	Source
Space group	P 2 21 21	Depositor
Cell constants a, b, c, α , β , γ	61.16Å 96.44Å 197.71Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.22 – 2.40 48.22 – 2.40	Depositor EDS
% Data completeness (in resolution range)	99.9 (48.22-2.40) 99.9 (48.22-2.40)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.36 (at 2.39Å)	Xtriage
Refinement program	PHENIX 1.14_3260, PHENIX 1.14_3260	Depositor
R, R_{free}	0.216 , 0.255 0.222 , 0.261	Depositor DCC
R_{free} test set	2307 reflections (4.94%)	wwPDB-VP
Wilson B-factor (Å ²)	65.4	Xtriage
Anisotropy	0.084	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 49.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	6821	wwPDB-VP
Average B, all atoms (Å ²)	87.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 19.16% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, CIT, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.20	0/3388	0.35	0/4562
1	B	0.18	0/3376	0.36	0/4542
2	C	0.61	0/26	0.83	0/32
All	All	0.20	0/6790	0.36	0/9136

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3338	0	3306	31	0
1	B	3326	0	3323	33	0
2	C	27	0	24	5	0
3	A	12	0	16	0	0
3	B	6	0	8	1	0
4	A	13	0	5	2	0
4	B	26	0	10	2	0
5	A	5	0	0	0	0
5	B	6	0	0	0	0
6	A	30	0	0	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	B	32	0	0	4	0
All	All	6821	0	6692	68	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (68) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:520:VAL:HB	1:B:551:GLU:HG2	1.58	0.85
1:B:485:THR:HG22	2:C:156:GLU:O	1.76	0.82
1:A:721[A]:ASP:OD1	1:A:721[A]:ASP:N	2.16	0.79
1:B:657:MET:HE1	1:B:698:ILE:HG12	1.69	0.74
1:A:665[B]:ASP:OD1	1:A:812:ARG:NH2	2.24	0.70
1:A:633:HIS:NE2	1:A:824:ASP:OD2	2.25	0.69
1:A:803:GLN:NE2	6:A:1002:HOH:O	2.26	0.69
1:B:520:VAL:HG13	1:B:528:LEU:HD11	1.76	0.66
1:A:822:THR:H	4:A:902:CIT:H42	1.60	0.64
1:B:795:PHE:O	1:B:799:VAL:HG12	1.98	0.63
1:B:820:LYS:O	1:B:823:ARG:NH2	2.32	0.62
1:B:485:THR:HA	2:C:156:GLU:O	2.00	0.62
1:B:482:PHE:N	2:C:158:LEU:O	2.33	0.61
1:B:783:ILE:HD13	1:B:795:PHE:HB3	1.83	0.61
1:A:471:ILE:HD12	1:A:512:LYS:HE2	1.85	0.59
1:A:626:LEU:HD12	1:A:642:LEU:HD11	1.86	0.57
1:B:498:ILE:HD11	1:B:519:ILE:HD11	1.86	0.56
4:B:903:CIT:O2	4:B:903:CIT:O7	2.23	0.55
4:A:902:CIT:O1	4:A:902:CIT:O7	2.21	0.55
1:B:485:THR:CG2	2:C:156:GLU:O	2.50	0.54
1:B:520:VAL:HG13	1:B:528:LEU:CD1	2.37	0.54
1:A:666:SER:OG	1:A:829:ARG:NH2	2.39	0.53
1:B:669:GLU:OE2	1:B:812:ARG:NH1	2.42	0.53
1:B:501:LEU:HD21	1:B:513:MET:HG2	1.90	0.53
1:B:699:ILE:HG23	1:B:707:ARG:HH11	1.75	0.51
1:B:489:GLY:H	1:B:494:GLU:HG3	1.76	0.51
1:B:812:ARG:NH2	6:B:1012:HOH:O	2.37	0.51
1:B:662:THR:HG21	1:B:701:ASP:HB3	1.92	0.51
1:B:523:ASN:O	1:B:524:ASP:HB2	2.11	0.50
1:A:740:ALA:HB1	1:A:748:ASP:HB3	1.93	0.50
1:B:487:VAL:HB	1:B:499:LYS:HD2	1.94	0.49
1:A:808:TYR:CZ	1:A:812:ARG:HD2	2.48	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:461:ARG:HA	1:A:528:LEU:HB3	1.95	0.48
1:B:463:PRO:HG3	1:B:528:LEU:HD12	1.95	0.48
1:B:592:GLU:H	1:B:592:GLU:HG2	1.37	0.48
4:B:903:CIT:H42	6:B:1016:HOH:O	2.15	0.47
1:B:498:ILE:HG13	1:B:513:MET:SD	2.55	0.47
1:B:485:THR:CB	2:C:156:GLU:O	2.62	0.47
1:A:721[A]:ASP:HB3	1:A:762:PRO:HG2	1.97	0.46
1:A:665[B]:ASP:CG	1:A:812:ARG:HH22	2.23	0.46
1:A:721[B]:ASP:CG	1:A:762:PRO:HG2	2.41	0.46
1:A:520:VAL:HG13	1:A:528:LEU:HD11	1.97	0.45
1:A:583:THR:O	1:A:587:THR:OG1	2.31	0.45
1:A:662:THR:HG21	1:A:701:ASP:HB3	1.98	0.45
1:A:763:HIS:O	1:A:767:VAL:HG13	2.15	0.45
1:B:735:ARG:HG3	1:B:774:TYR:CE1	2.52	0.45
1:A:795:PHE:O	1:A:799:VAL:HG13	2.17	0.44
1:A:820:LYS:O	1:A:823:ARG:NH2	2.41	0.44
1:A:759:ARG:HB2	1:A:764:LEU:HG	2.00	0.44
1:A:466:LEU:HD11	1:A:528:LEU:HD11	2.00	0.44
1:A:604:ASN:O	1:A:608:GLN:HG2	2.17	0.44
1:B:512:LYS:O	1:B:554:ARG:NH1	2.51	0.44
1:A:705:ASP:HB2	1:A:791:LEU:HD22	2.00	0.44
1:A:827:LEU:O	1:A:831:MET:HG2	2.18	0.43
1:B:705:ASP:HB2	1:B:791:LEU:HD22	2.00	0.43
1:B:477:LYS:HD2	1:B:542:ILE:HB	2.00	0.43
1:B:731:ASP:HB3	6:B:1008:HOH:O	2.18	0.43
1:A:676:ASN:ND2	1:A:718:ARG:O	2.48	0.42
1:A:477:LYS:HB2	1:A:482:PHE:CZ	2.54	0.42
1:B:626:LEU:HD12	1:B:642:LEU:HD11	2.02	0.42
1:A:598:ILE:O	1:A:602:ARG:HG2	2.20	0.42
1:A:780:LEU:HD23	1:A:780:LEU:HA	1.89	0.42
1:A:856:TYR:O	1:A:860:GLN:HG2	2.20	0.41
1:B:657:MET:HE3	1:B:667:LEU:HD13	2.01	0.41
1:B:491:GLU:O	1:B:492:PRO:C	2.63	0.41
1:A:539:PHE:CE1	1:A:550:LEU:HD21	2.56	0.41
3:B:902:GOL:O3	6:B:1001:HOH:O	2.22	0.40
1:B:699:ILE:O	1:B:707:ARG:NH1	2.55	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	423/427 (99%)	409 (97%)	12 (3%)	2 (0%)	24	37
1	B	420/427 (98%)	406 (97%)	12 (3%)	2 (0%)	24	37
2	C	1/10 (10%)	1 (100%)	0	0	100	100
All	All	844/864 (98%)	816 (97%)	24 (3%)	4 (0%)	24	37

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	557	PRO
1	A	590	VAL
1	B	492	PRO
1	B	590	VAL

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	361/370 (98%)	340 (94%)	21 (6%)	18	32
1	B	362/370 (98%)	347 (96%)	15 (4%)	27	46
2	C	3/10 (30%)	2 (67%)	1 (33%)	0	0
All	All	726/750 (97%)	689 (95%)	37 (5%)	22	37

All (37) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	473	THR
1	A	490	ASP
1	A	491	GLU
1	A	509	LEU
1	A	515	THR
1	A	548	VAL
1	A	592	GLU
1	A	618	THR
1	A	627	LYS
1	A	662	THR
1	A	664	GLU
1	A	680	GLN
1	A	696	LYS
1	A	721[A]	ASP
1	A	721[B]	ASP
1	A	745	LYS
1	A	767	VAL
1	A	787	VAL
1	A	788	LYS
1	A	799	VAL
1	A	861	ASP
1	B	513	MET
1	B	523	ASN
1	B	535	VAL
1	B	551	GLU
1	B	576	ARG
1	B	592	GLU
1	B	616	ARG
1	B	627	LYS
1	B	664	GLU
1	B	680	GLN
1	B	701	ASP
1	B	745	LYS
1	B	787	VAL
1	B	799	VAL
1	B	861	ASP
2	C	157	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	523	ASN
1	B	497	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	571	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 17 ligands modelled in this entry, 11 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	CIT	B	901	-	12,12,12	1.02	0	17,17,17	1.44	2 (11%)
4	CIT	A	902	-	12,12,12	1.23	3 (25%)	17,17,17	1.35	2 (11%)
3	GOL	B	902	-	5,5,5	0.87	0	5,5,5	1.05	0
4	CIT	B	903	-	12,12,12	1.08	0	17,17,17	1.50	2 (11%)
3	GOL	A	901	-	5,5,5	0.85	0	5,5,5	1.10	0
3	GOL	A	903	-	5,5,5	0.91	0	5,5,5	1.09	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	CIT	B	901	-	-	9/16/16/16	-
4	CIT	A	902	-	-	10/16/16/16	-
3	GOL	B	902	-	-	2/4/4/4	-
4	CIT	B	903	-	-	9/16/16/16	-
3	GOL	A	901	-	-	2/4/4/4	-
3	GOL	A	903	-	-	0/4/4/4	-

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	902	CIT	O2-C1	-2.18	1.23	1.30
4	A	902	CIT	O4-C5	-2.08	1.23	1.30
4	A	902	CIT	C3-C6	2.05	1.55	1.53

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	903	CIT	O6-C6-C3	3.64	120.12	113.14
4	B	901	CIT	O6-C6-C3	3.50	119.85	113.14
4	A	902	CIT	O6-C6-C3	3.01	118.91	113.14
4	A	902	CIT	O5-C6-C3	-2.82	116.64	122.09
4	B	903	CIT	O2-C1-C2	2.19	121.28	114.35
4	B	901	CIT	O4-C5-C4	2.16	121.19	114.35

There are no chirality outliers.

All (32) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	902	CIT	C2-C3-C6-O5
4	A	902	CIT	C2-C3-C6-O6
4	A	902	CIT	O7-C3-C6-O5
4	A	902	CIT	O7-C3-C6-O6
4	B	901	CIT	O7-C3-C6-O5
4	B	901	CIT	O7-C3-C6-O6
4	B	901	CIT	C4-C3-C6-O5
4	B	901	CIT	C4-C3-C6-O6
4	B	903	CIT	C1-C2-C3-C4
4	B	903	CIT	C1-C2-C3-C6
4	B	903	CIT	C2-C3-C6-O5
4	B	903	CIT	C2-C3-C6-O6
4	B	903	CIT	O7-C3-C6-O5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
4	B	903	CIT	O7-C3-C6-O6
4	A	902	CIT	C2-C3-C4-C5
4	A	902	CIT	O7-C3-C4-C5
4	A	902	CIT	C6-C3-C4-C5
3	B	902	GOL	O1-C1-C2-O2
4	A	902	CIT	C1-C2-C3-C6
4	B	901	CIT	C2-C3-C4-C5
4	B	903	CIT	C1-C2-C3-O7
3	A	901	GOL	C1-C2-C3-O3
3	B	902	GOL	O1-C1-C2-C3
4	B	901	CIT	C2-C3-C6-O5
4	A	902	CIT	C1-C2-C3-O7
4	B	903	CIT	O1-C1-C2-C3
4	B	903	CIT	O2-C1-C2-C3
4	A	902	CIT	C1-C2-C3-C4
4	B	901	CIT	C6-C3-C4-C5
4	B	901	CIT	C2-C3-C6-O6
3	A	901	GOL	O2-C2-C3-O3
4	B	901	CIT	O7-C3-C4-C5

There are no ring outliers.

3 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	902	CIT	2	0
3	B	902	GOL	1	0
4	B	903	CIT	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	421/427 (98%)	0.17	21 (4%) 34 30	29, 77, 154, 164	4 (0%)
1	B	421/427 (98%)	0.22	10 (2%) 59 55	47, 83, 117, 135	1 (0%)
2	C	3/10 (30%)	0.84	0 100 100	118, 118, 128, 135	0
All	All	845/864 (97%)	0.20	31 (3%) 45 41	29, 80, 152, 164	5 (0%)

All (31) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	519	ILE	4.2
1	A	553	CYS	3.3
1	A	721[A]	ASP	3.3
1	B	500	SER	3.2
1	B	566	VAL	3.1
1	A	544	ILE	3.0
1	B	458	PHE	2.9
1	A	503	LEU	2.9
1	A	520	VAL	2.7
1	B	487	VAL	2.7
1	B	503	LEU	2.7
1	A	502	VAL	2.7
1	B	528	LEU	2.6
1	A	522	VAL	2.6
1	A	524	ASP	2.6
1	A	547	SER	2.6
1	A	458	PHE	2.5
1	B	547	SER	2.5
1	A	538	ILE	2.5
1	A	487	VAL	2.4
1	B	576	ARG	2.4
1	A	501	LEU	2.4
1	B	721[A]	ASP	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	535	VAL	2.3
1	A	525	THR	2.2
1	A	533	ALA	2.2
1	A	471	ILE	2.2
1	A	527	VAL	2.1
1	A	560	SER	2.1
1	B	557	PRO	2.1
1	A	548	VAL	2.1

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	CA	B	907	1/1	0.14	0.12	186,186,186,186	0
5	CA	B	905	1/1	0.56	0.11	131,131,131,131	0
4	CIT	A	902	13/13	0.66	0.12	96,101,112,117	0
5	CA	A	904	1/1	0.70	0.15	142,142,142,142	0
4	CIT	B	903	13/13	0.72	0.14	116,128,133,134	0
5	CA	B	908	1/1	0.72	0.17	115,115,115,115	0
4	CIT	B	901	13/13	0.74	0.13	105,139,150,150	0
5	CA	A	906	1/1	0.75	0.11	139,139,139,139	0
5	CA	A	907	1/1	0.86	0.17	101,101,101,101	0
5	CA	B	904	1/1	0.89	0.18	99,99,99,99	0
3	GOL	A	903	6/6	0.89	0.18	80,95,103,104	0
5	CA	B	906	1/1	0.90	0.08	140,140,140,140	0
5	CA	A	905	1/1	0.93	0.06	163,163,163,163	0
3	GOL	B	902	6/6	0.94	0.10	52,64,70,77	0
3	GOL	A	901	6/6	0.95	0.12	56,63,72,81	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
5	CA	B	909	1/1	0.95	0.07	94,94,94,94	0
5	CA	A	908	1/1	0.98	0.11	85,85,85,85	0

6.5 Other polymers [i](#)

There are no such residues in this entry.