



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 12, 2026 – 05:44 PM UTC

PDB ID : 8TW3 / pdb_00008tw3
Title : hMPV fusion protein complexed with single domain antibodies sdHMPV16 and sdHMPV12
Authors : Rush, S.A.; McLellan, J.S.
Deposited on : 2023-08-19
Resolution : 2.90 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2481 (2.90-2.90)
Clashscore	190562	2690 (2.90-2.90)
Ramachandran outliers	187476	2623 (2.90-2.90)
Sidechain outliers	187428	2625 (2.90-2.90)
RSRZ outliers	180081	2481 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	193	0% 61% 36%
1	Y	193	62% 36%
2	B	192	2% 56% 7% 36%
2	Z	192	9% 54% 8% 38%
3	F	542	0% 68% 28%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	X	542	 A horizontal bar chart showing the quality of chain 3. The bar is divided into four segments: a small red segment at the beginning labeled '2%', followed by a large green segment labeled '61%', a small yellow segment labeled '8%', and a grey segment at the end labeled '31%'. The segments are separated by thin white lines.

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 18851 atoms, of which 9276 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Single domain antibody sdHMPV12.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	123	Total	C	H	N	O	S	0	0	0
			1813	568	894	164	182	5			
1	Y	123	Total	C	H	N	O	S	0	0	0
			1813	568	894	164	182	5			

- Molecule 2 is a protein called Single domain antibody sdHMPV16.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	B	123	Total	C	H	N	O	S	0	0	0
			1826	580	883	170	188	5			
2	Z	119	Total	C	H	N	O	S	0	0	0
			1774	570	851	166	182	5			

- Molecule 3 is a protein called Fusion glycoprotein.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
3	F	388	Total	C	H	N	O	S	0	0	0
			5918	1844	2966	510	577	21			
3	X	376	Total	C	H	N	O	S	0	0	0
			5599	1792	2736	494	556	21			

There are 106 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
F	185	PRO	ALA	conflict	UNP H6X1Z0
F	491	GLY	-	expression tag	UNP H6X1Z0
F	492	GLY	-	expression tag	UNP H6X1Z0
F	493	GLY	-	expression tag	UNP H6X1Z0
F	494	SER	-	expression tag	UNP H6X1Z0
F	495	LEU	-	expression tag	UNP H6X1Z0
F	496	GLU	-	expression tag	UNP H6X1Z0
F	497	VAL	-	expression tag	UNP H6X1Z0

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
F	498	LEU	-	expression tag	UNP H6X1Z0
F	499	PHE	-	expression tag	UNP H6X1Z0
F	500	GLN	-	expression tag	UNP H6X1Z0
F	501	GLY	-	expression tag	UNP H6X1Z0
F	502	PRO	-	expression tag	UNP H6X1Z0
F	503	GLY	-	expression tag	UNP H6X1Z0
F	504	HIS	-	expression tag	UNP H6X1Z0
F	505	HIS	-	expression tag	UNP H6X1Z0
F	506	HIS	-	expression tag	UNP H6X1Z0
F	507	HIS	-	expression tag	UNP H6X1Z0
F	508	HIS	-	expression tag	UNP H6X1Z0
F	509	HIS	-	expression tag	UNP H6X1Z0
F	510	HIS	-	expression tag	UNP H6X1Z0
F	511	HIS	-	expression tag	UNP H6X1Z0
F	512	SER	-	expression tag	UNP H6X1Z0
F	513	ALA	-	expression tag	UNP H6X1Z0
F	514	TRP	-	expression tag	UNP H6X1Z0
F	515	SER	-	expression tag	UNP H6X1Z0
F	516	HIS	-	expression tag	UNP H6X1Z0
F	517	PRO	-	expression tag	UNP H6X1Z0
F	518	GLN	-	expression tag	UNP H6X1Z0
F	519	PHE	-	expression tag	UNP H6X1Z0
F	520	GLU	-	expression tag	UNP H6X1Z0
F	521	LYS	-	expression tag	UNP H6X1Z0
F	522	GLY	-	expression tag	UNP H6X1Z0
F	523	GLY	-	expression tag	UNP H6X1Z0
F	524	GLY	-	expression tag	UNP H6X1Z0
F	525	SER	-	expression tag	UNP H6X1Z0
F	526	GLY	-	expression tag	UNP H6X1Z0
F	527	GLY	-	expression tag	UNP H6X1Z0
F	528	GLY	-	expression tag	UNP H6X1Z0
F	529	GLY	-	expression tag	UNP H6X1Z0
F	530	SER	-	expression tag	UNP H6X1Z0
F	531	GLY	-	expression tag	UNP H6X1Z0
F	532	GLY	-	expression tag	UNP H6X1Z0
F	533	SER	-	expression tag	UNP H6X1Z0
F	534	ALA	-	expression tag	UNP H6X1Z0
F	535	TRP	-	expression tag	UNP H6X1Z0
F	536	SER	-	expression tag	UNP H6X1Z0
F	537	HIS	-	expression tag	UNP H6X1Z0
F	538	PRO	-	expression tag	UNP H6X1Z0
F	539	GLN	-	expression tag	UNP H6X1Z0

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
F	540	PHE	-	expression tag	UNP H6X1Z0
F	541	GLU	-	expression tag	UNP H6X1Z0
F	542	LYS	-	expression tag	UNP H6X1Z0
X	185	PRO	ALA	conflict	UNP H6X1Z0
X	491	GLY	-	expression tag	UNP H6X1Z0
X	492	GLY	-	expression tag	UNP H6X1Z0
X	493	GLY	-	expression tag	UNP H6X1Z0
X	494	SER	-	expression tag	UNP H6X1Z0
X	495	LEU	-	expression tag	UNP H6X1Z0
X	496	GLU	-	expression tag	UNP H6X1Z0
X	497	VAL	-	expression tag	UNP H6X1Z0
X	498	LEU	-	expression tag	UNP H6X1Z0
X	499	PHE	-	expression tag	UNP H6X1Z0
X	500	GLN	-	expression tag	UNP H6X1Z0
X	501	GLY	-	expression tag	UNP H6X1Z0
X	502	PRO	-	expression tag	UNP H6X1Z0
X	503	GLY	-	expression tag	UNP H6X1Z0
X	504	HIS	-	expression tag	UNP H6X1Z0
X	505	HIS	-	expression tag	UNP H6X1Z0
X	506	HIS	-	expression tag	UNP H6X1Z0
X	507	HIS	-	expression tag	UNP H6X1Z0
X	508	HIS	-	expression tag	UNP H6X1Z0
X	509	HIS	-	expression tag	UNP H6X1Z0
X	510	HIS	-	expression tag	UNP H6X1Z0
X	511	HIS	-	expression tag	UNP H6X1Z0
X	512	SER	-	expression tag	UNP H6X1Z0
X	513	ALA	-	expression tag	UNP H6X1Z0
X	514	TRP	-	expression tag	UNP H6X1Z0
X	515	SER	-	expression tag	UNP H6X1Z0
X	516	HIS	-	expression tag	UNP H6X1Z0
X	517	PRO	-	expression tag	UNP H6X1Z0
X	518	GLN	-	expression tag	UNP H6X1Z0
X	519	PHE	-	expression tag	UNP H6X1Z0
X	520	GLU	-	expression tag	UNP H6X1Z0
X	521	LYS	-	expression tag	UNP H6X1Z0
X	522	GLY	-	expression tag	UNP H6X1Z0
X	523	GLY	-	expression tag	UNP H6X1Z0
X	524	GLY	-	expression tag	UNP H6X1Z0
X	525	SER	-	expression tag	UNP H6X1Z0
X	526	GLY	-	expression tag	UNP H6X1Z0
X	527	GLY	-	expression tag	UNP H6X1Z0
X	528	GLY	-	expression tag	UNP H6X1Z0

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
X	529	GLY	-	expression tag	UNP H6X1Z0
X	530	SER	-	expression tag	UNP H6X1Z0
X	531	GLY	-	expression tag	UNP H6X1Z0
X	532	GLY	-	expression tag	UNP H6X1Z0
X	533	SER	-	expression tag	UNP H6X1Z0
X	534	ALA	-	expression tag	UNP H6X1Z0
X	535	TRP	-	expression tag	UNP H6X1Z0
X	536	SER	-	expression tag	UNP H6X1Z0
X	537	HIS	-	expression tag	UNP H6X1Z0
X	538	PRO	-	expression tag	UNP H6X1Z0
X	539	GLN	-	expression tag	UNP H6X1Z0
X	540	PHE	-	expression tag	UNP H6X1Z0
X	541	GLU	-	expression tag	UNP H6X1Z0
X	542	LYS	-	expression tag	UNP H6X1Z0

- Molecule 4 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: $C_8H_{15}NO_6$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	F	1	Total	C	H	N	O	0	0
			27	8	13	1	5		
4	F	1	Total	C	H	N	O	0	0
			27	8	13	1	5		
4	X	1	Total	C	H	N	O	0	0
			27	8	13	1	5		
4	X	1	Total	C	H	N	O	0	0
			27	8	13	1	5		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Single domain antibody sdHMPV12




- Molecule 1: Single domain antibody sdHMPV12




- Molecule 2: Single domain antibody sdHMPV16




- Molecule 2: Single domain antibody sdHMPV16



- Molecule 3: Fusion glycoprotein



- Molecule 3: Fusion glycoprotein



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	80.26Å 115.61Å 158.03Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	44.97 – 2.90 44.97 – 2.90	Depositor EDS
% Data completeness (in resolution range)	97.9 (44.97-2.90) 97.9 (44.97-2.90)	Depositor EDS
R_{merge}	0.21	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.10 (at 2.91Å)	Xtriage
Refinement program	PHENIX (1.20.1_4487: ???)	Depositor
R, R_{free}	0.191 , 0.232 0.191 , 0.232	Depositor DCC
R_{free} test set	1543 reflections (4.63%)	wwPDB-VP
Wilson B-factor (Å ²)	41.0	Xtriage
Anisotropy	0.193	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 35.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	18851	wwPDB-VP
Average B, all atoms (Å ²)	48.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.24% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.15	0/936	0.32	0/1271
1	Y	0.16	0/936	0.33	0/1271
2	B	0.16	0/962	0.34	0/1302
2	Z	0.14	0/941	0.28	0/1273
3	F	0.14	0/2991	0.28	0/4052
3	X	0.14	0/2902	0.29	0/3931
All	All	0.15	0/9668	0.30	0/13100

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	919	894	896	3	0
1	Y	919	894	896	3	0
2	B	943	883	890	12	0
2	Z	923	851	873	9	0
3	F	2952	2966	2973	13	0
3	X	2863	2736	2889	29	0
4	F	28	26	26	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	X	28	26	26	2	0
All	All	9575	9276	9469	67	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 67 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:X:287:LYS:NZ	3:X:323:GLU:OE2	2.00	0.94
3:X:184:ILE:HG22	3:X:186:ASP:H	1.47	0.80
2:Z:97:GLU:N	2:Z:97:GLU:OE1	2.18	0.76
3:X:234:MET:HE1	3:X:244:MET:SD	2.26	0.76
3:F:234:MET:HE1	3:F:244:MET:SD	2.29	0.72

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	121/193 (63%)	117 (97%)	4 (3%)	0	100	100
1	Y	121/193 (63%)	117 (97%)	4 (3%)	0	100	100
2	B	121/192 (63%)	118 (98%)	3 (2%)	0	100	100
2	Z	115/192 (60%)	110 (96%)	5 (4%)	0	100	100
3	F	384/542 (71%)	374 (97%)	10 (3%)	0	100	100
3	X	372/542 (69%)	357 (96%)	15 (4%)	0	100	100
All	All	1234/1854 (67%)	1193 (97%)	41 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	99/151 (66%)	99 (100%)	0	100	100
1	Y	99/151 (66%)	99 (100%)	0	100	100
2	B	98/149 (66%)	96 (98%)	2 (2%)	48	78
2	Z	96/149 (64%)	96 (100%)	0	100	100
3	F	329/453 (73%)	325 (99%)	4 (1%)	63	86
3	X	320/453 (71%)	317 (99%)	3 (1%)	70	90
All	All	1041/1506 (69%)	1032 (99%)	9 (1%)	70	90

5 of 9 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	X	174	THR
3	X	187	LEU
3	F	171	LYS
3	F	272	GLN
3	F	285	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 10 such sidechains are listed below:

Mol	Chain	Res	Type
3	X	353	ASN
3	X	358	ASN
2	Z	114	ASN
3	F	332	HIS
3	X	178	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	NAG	X	601	-	14,14,15	0.19	0	17,19,21	0.55	0
4	NAG	F	601	-	14,14,15	0.22	0	17,19,21	0.48	0
4	NAG	X	602	3	14,14,15	0.32	0	17,19,21	0.48	0
4	NAG	F	602	3	14,14,15	0.21	0	17,19,21	0.51	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	X	601	-	-	0/6/23/26	0/1/1/1
4	NAG	F	601	-	-	0/6/23/26	0/1/1/1
4	NAG	X	602	3	-	0/6/23/26	0/1/1/1
4	NAG	F	602	3	-	2/6/23/26	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	F	602	NAG	O5-C5-C6-O6
4	F	602	NAG	C4-C5-C6-O6

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	X	601	NAG	2	0
4	F	602	NAG	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	123/193 (63%)	-0.22	1 (0%) 82 77	22, 34, 51, 74	0
1	Y	123/193 (63%)	-0.22	0 100 100	20, 35, 56, 73	0
2	B	123/192 (64%)	0.57	3 (2%) 59 50	24, 54, 86, 115	0
2	Z	119/192 (61%)	0.89	17 (14%) 6 5	32, 64, 93, 103	0
3	F	388/542 (71%)	0.04	6 (1%) 72 64	20, 44, 81, 119	0
3	X	376/542 (69%)	0.14	11 (2%) 53 45	18, 45, 86, 101	0
All	All	1252/1854 (67%)	0.15	38 (3%) 52 43	18, 44, 85, 119	0

The worst 5 of 38 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	F	93	GLU	3.8
3	X	64	PRO	3.4
2	B	18	SER	3.1
2	Z	88	TYR	2.9
3	X	185	PRO	2.9

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	NAG	X	601	14/15	0.69	0.19	53,88,110,113	0
4	NAG	F	601	14/15	0.78	0.14	60,78,101,115	0
4	NAG	F	602	14/15	0.86	0.12	53,64,73,78	0
4	NAG	X	602	14/15	0.86	0.13	54,64,76,77	0

6.5 Other polymers [i](#)

There are no such residues in this entry.