



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 2, 2026 – 01:11 AM UTC

PDB ID : 3TXV / pdb_00003txv
Title : Crystal structure of a probable tagatose 6 phosphate kinase from *Sinorhizobium meliloti* 1021
Authors : Agarwal, R.; Chamala, S.; Evans, B.; Foti, R.; Gizzi, A.; Hillerich, B.; Kar, A.; LaFleur, J.; Seidel, R.; Villigas, G.; Zencheck, W.; Almo, S.C.; Swaminathan, S.; New York Structural Genomics Research Consortium (NYSGRG)
Deposited on : 2011-09-23
Resolution : 2.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3866 (2.80-2.80)
Clashscore	190562	4276 (2.80-2.80)
Ramachandran outliers	187476	4196 (2.80-2.80)
Sidechain outliers	187428	4198 (2.80-2.80)
RSRZ outliers	180081	3869 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	450	3% 63% 19% • 14%

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 2878 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Probable tagatose 6-phosphate kinase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
1	A	387	2847	1811	486	538	4	8	0	0	0

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MSE	-	expression tag	UNP Q92UY3
A	2	VAL	-	expression tag	UNP Q92UY3
A	429	ALA	-	expression tag	UNP Q92UY3
A	430	GLU	-	expression tag	UNP Q92UY3
A	431	ASN	-	expression tag	UNP Q92UY3
A	432	LEU	-	expression tag	UNP Q92UY3
A	433	TYR	-	expression tag	UNP Q92UY3
A	434	PHE	-	expression tag	UNP Q92UY3
A	435	GLN	-	expression tag	UNP Q92UY3
A	436	SER	-	expression tag	UNP Q92UY3
A	437	HIS	-	expression tag	UNP Q92UY3
A	438	HIS	-	expression tag	UNP Q92UY3
A	439	HIS	-	expression tag	UNP Q92UY3
A	440	HIS	-	expression tag	UNP Q92UY3
A	441	HIS	-	expression tag	UNP Q92UY3
A	442	HIS	-	expression tag	UNP Q92UY3
A	443	TRP	-	expression tag	UNP Q92UY3
A	444	SER	-	expression tag	UNP Q92UY3
A	445	HIS	-	expression tag	UNP Q92UY3
A	446	PRO	-	expression tag	UNP Q92UY3
A	447	GLN	-	expression tag	UNP Q92UY3
A	448	PHE	-	expression tag	UNP Q92UY3
A	449	GLU	-	expression tag	UNP Q92UY3
A	450	LYS	-	expression tag	UNP Q92UY3


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	31	Total 31	O 31	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Probable tagatose 6-phosphate kinase



4 Data and refinement statistics

Property	Value	Source
Space group	P 63 2 2	Depositor
Cell constants a, b, c, α , β , γ	140.00Å 140.00Å 97.36Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	48.68 – 2.80 48.68 – 2.80	Depositor EDS
% Data completeness (in resolution range)	99.2 (48.68-2.80) 99.1 (48.68-2.80)	Depositor EDS
R_{merge}	0.17	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.43 (at 2.81Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.240 , 0.293 0.243 , 0.294	Depositor DCC
R_{free} test set	716 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å ²)	54.7	Xtriage
Anisotropy	0.107	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 37.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.89	EDS
Total number of atoms	2878	wwPDB-VP
Average B, all atoms (Å ²)	45.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.63% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.15	6/2903 (0.2%)	1.21	19/3946 (0.5%)

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	327	MSE	SE-CE	-7.09	1.74	1.95
1	A	142	ALA	CA-CB	-6.61	1.44	1.53
1	A	326	VAL	CA-CB	-6.16	1.46	1.54
1	A	288	VAL	C-O	-6.07	1.18	1.24
1	A	139	GLU	CA-C	-5.54	1.47	1.52
1	A	322	VAL	CA-CB	-5.23	1.48	1.54

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	44	GLU	N-CA-C	9.80	125.31	113.16
1	A	89	ILE	CB-CA-C	-7.36	101.78	110.91
1	A	330	GLU	CA-C-N	7.29	126.93	119.56
1	A	330	GLU	C-N-CA	7.29	126.93	119.56
1	A	407	ALA	N-CA-C	7.27	119.21	111.28
1	A	213	GLU	N-CA-C	-6.90	103.75	111.28
1	A	158	VAL	N-CA-C	-6.00	103.89	112.35
1	A	193	GLU	N-CA-C	5.83	118.68	110.23
1	A	213	GLU	CA-C-N	5.79	128.04	120.28
1	A	213	GLU	C-N-CA	5.79	128.04	120.28
1	A	44	GLU	CA-C-N	5.56	131.71	121.70
1	A	44	GLU	C-N-CA	5.56	131.71	121.70
1	A	136	CYS	N-CA-C	-5.43	100.06	108.90
1	A	315	ARG	N-CA-C	5.41	117.16	107.80
1	A	289	GLY	CA-C-N	5.33	124.97	119.05
1	A	289	GLY	C-N-CA	5.33	124.97	119.05
1	A	405	PRO	CB-CA-C	-5.16	106.13	112.89
1	A	137	ALA	N-CA-C	5.07	117.58	110.23

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	326	VAL	N-CA-C	5.06	115.78	110.62

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2847	0	2716	56	0
2	A	31	0	0	4	0
All	All	2878	0	2716	56	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (56) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:318:THR:HG22	1:A:321:GLU:H	1.12	1.11
1:A:174:ILE:HD11	1:A:227:VAL:HG22	1.43	1.00
1:A:46:ALA:HB1	1:A:47:PRO:CD	1.94	0.97
1:A:46:ALA:HB1	1:A:47:PRO:HD2	1.44	0.96
1:A:202:GLU:HG2	2:A:474:HOH:O	1.69	0.90
1:A:204:VAL:HG11	1:A:257:LEU:HD21	1.52	0.90
1:A:318:THR:HG22	1:A:321:GLU:N	1.91	0.84
1:A:116:MSE:HG3	1:A:117:ILE:N	1.97	0.80
1:A:136:CYS:O	1:A:139:GLU:HG3	1.82	0.78
1:A:348:GLN:HG3	1:A:352:SER:HB2	1.71	0.72
1:A:40:ARG:NH2	1:A:420:GLN:HG3	2.05	0.71
1:A:394:SER:O	1:A:398:VAL:HG23	1.93	0.68
1:A:204:VAL:HG11	1:A:257:LEU:CD2	2.25	0.64
1:A:116:MSE:HG3	1:A:117:ILE:H	1.64	0.60
1:A:204:VAL:CG1	1:A:257:LEU:HD21	2.29	0.60
1:A:113:ALA:O	1:A:116:MSE:HG3	2.02	0.59
1:A:137:ALA:HB2	2:A:459:HOH:O	2.03	0.58

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:37:ALA:HA	1:A:419:LEU:CD1	2.34	0.57
1:A:134:MSE:H	1:A:134:MSE:SE	2.38	0.56
1:A:135:GLY:HA2	1:A:143:LEU:HD11	1.88	0.55
1:A:230:PRO:HB3	1:A:283:PHE:CE1	2.42	0.55
1:A:173:TYR:HB2	1:A:223:VAL:HG22	1.90	0.54
1:A:207:HIS:O	1:A:211:PHE:HD1	1.91	0.54
1:A:265:HIS:NE2	1:A:287:LYS:HD2	2.23	0.54
1:A:348:GLN:HG3	1:A:352:SER:CB	2.36	0.53
1:A:408:LYS:HB3	1:A:409:PRO:HD3	1.89	0.53
1:A:290:PRO:HD3	2:A:468:HOH:O	2.10	0.50
1:A:25:SER:HB3	1:A:287:LYS:HA	1.93	0.50
1:A:116:MSE:CG	1:A:117:ILE:N	2.73	0.50
1:A:147:THR:O	1:A:151:ARG:HG3	2.12	0.50
1:A:346:ARG:O	1:A:347:LEU:C	2.55	0.49
1:A:324:GLU:OE1	1:A:346:ARG:HG3	2.12	0.49
1:A:174:ILE:HD11	1:A:227:VAL:CG2	2.29	0.49
1:A:160:GLU:CD	1:A:222:ARG:HH21	2.21	0.49
1:A:265:HIS:CD2	1:A:287:LYS:HD2	2.47	0.49
1:A:37:ALA:HA	1:A:419:LEU:HD11	1.94	0.48
1:A:46:ALA:CB	1:A:47:PRO:CD	2.74	0.48
1:A:109:ALA:CB	1:A:136:CYS:SG	3.03	0.47
1:A:135:GLY:HA2	1:A:143:LEU:CD1	2.45	0.47
1:A:244:ARG:HH11	1:A:244:ARG:CG	2.27	0.47
1:A:80:ASP:O	1:A:82:ILE:N	2.48	0.46
1:A:228:VAL:O	1:A:230:PRO:HD3	2.16	0.46
1:A:318:THR:O	1:A:322:VAL:HG23	2.15	0.46
1:A:98:PRO:HG3	1:A:110:MSE:SE	2.67	0.45
1:A:103:HIS:H	1:A:103:HIS:CD2	2.33	0.45
1:A:160:GLU:OE1	1:A:222:ARG:NH2	2.52	0.43
1:A:267:THR:HG21	1:A:286:LEU:CD2	2.48	0.42
1:A:306:ILE:HG23	1:A:310:LEU:HD22	2.01	0.42
1:A:346:ARG:HH11	1:A:346:ARG:HB3	1.84	0.42
1:A:242:TYR:CD2	1:A:278:LEU:HG	2.56	0.41
1:A:301:TYR:C	1:A:305:GLN:HE21	2.26	0.41
1:A:54:CYS:HA	1:A:116:MSE:HE1	2.03	0.41
1:A:30:HIS:HA	1:A:31:PRO:HD2	1.80	0.40
1:A:235:GLY:O	1:A:353:TYR:HB2	2.21	0.40
1:A:54:CYS:HB3	1:A:95:HIS:O	2.22	0.40
1:A:426:CYS:C	2:A:477:HOH:O	2.64	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	381/450 (85%)	362 (95%)	16 (4%)	3 (1%)	16	44

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	81	ARG
1	A	289	GLY
1	A	272	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	270/332 (81%)	235 (87%)	35 (13%)	4	14

All (35) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	27	CYS
1	A	28	SER
1	A	57	VAL
1	A	72	THR
1	A	83	GLU
1	A	90	LEU
1	A	96	LEU
1	A	117	ILE

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	118	THR
1	A	128	LEU
1	A	132	THR
1	A	174	ILE
1	A	192	LEU
1	A	202	GLU
1	A	204	VAL
1	A	227	VAL
1	A	238	ASN
1	A	244	ARG
1	A	250	LEU
1	A	256	GLN
1	A	266	SER
1	A	273	ASP
1	A	275	LEU
1	A	287	LYS
1	A	310	LEU
1	A	315	ARG
1	A	318	THR
1	A	323	THR
1	A	336	LYS
1	A	347	LEU
1	A	348	GLN
1	A	352	SER
1	A	408	LYS
1	A	412	LEU
1	A	420	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	56	GLN
1	A	103	HIS
1	A	420	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å ²)	Q < 0.9
1	A	379/450 (84%)	0.27	13 (3%)	48 39	24, 45, 62, 73	0

All (13) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	192	LEU	5.6
1	A	216	ALA	4.0
1	A	193	GLU	3.9
1	A	238	ASN	3.6
1	A	154	ARG	3.4
1	A	170	LEU	3.1
1	A	21	ARG	3.1
1	A	257	LEU	2.9
1	A	231	GLY	2.5
1	A	137	ALA	2.4
1	A	246	ARG	2.4
1	A	54	CYS	2.2
1	A	194	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.