



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 05:02 PM UTC

PDB ID : 1TZB / pdb_00001tzb
Title : Crystal structure of native phosphoglucose/phosphomannose isomerase from *Pyrobaculum aerophilum*
Authors : Swan, M.K.; Hansen, T.; Schoenheit, P.; Davies, C.
Deposited on : 2004-07-09
Resolution : 1.16 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.16 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1256 (1.18-1.14)
Clashscore	190562	1272 (1.18-1.14)
Ramachandran outliers	187476	1246 (1.18-1.14)
Sidechain outliers	187428	1246 (1.18-1.14)
RSRZ outliers	180081	1255 (1.18-1.14)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	302	20% 90% 9% .
1	B	302	19% 92% 7% ..

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 5537 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called glucose-6-phosphate isomerase, conjectural.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	301	Total	C	N	O	S	2	26	0
			2434	1558	427	442	7			
1	B	300	Total	C	N	O	S	0	26	0
			2426	1554	422	444	6			

- Molecule 2 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	B	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 3 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	331	Total	O	1	0
			331	331		
4	B	294	Total	O	2	0
			294	294		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: glucose-6-phosphate isomerase, conjectural



- Molecule 1: glucose-6-phosphate isomerase, conjectural



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	55.10Å 100.80Å 55.80Å 90.00° 113.20° 90.00°	Depositor
Resolution (Å)	36.00 – 1.16 36.00 – 1.16	Depositor EDS
% Data completeness (in resolution range)	87.7 (36.00-1.16) 87.6 (36.00-1.16)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.92 (at 1.16Å)	Xtriage
Refinement program	REFMAC 5.1.24	Depositor
R, R_{free}	0.149 , 0.165 0.174 , 0.188	Depositor DCC
R_{free} test set	8508 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å ²)	10.3	Xtriage
Anisotropy	0.450	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 40.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.016 for l,-k,h	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	5537	wwPDB-VP
Average B, all atoms (Å ²)	14.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.27% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.75	3/2592 (0.1%)	0.73	3/3520 (0.1%)
1	B	0.46	0/2597	0.69	0/3528
All	All	0.62	3/5189 (0.1%)	0.71	3/7048 (0.0%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	161	GLU	CD-OE2	27.88	1.78	1.25
1	A	161	GLU	CB-CG	10.01	1.82	1.52
1	A	161	GLU	CD-OE1	5.11	1.35	1.25

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	161	GLU	CG-CD-OE1	-12.81	88.93	118.40
1	A	161	GLU	OE1-CD-OE2	6.26	137.93	122.90
1	A	161	GLU	CG-CD-OE2	5.74	131.61	118.40

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2434	0	2486	28	0
1	B	2426	0	2458	28	0
2	A	20	0	0	1	0
2	B	20	0	0	1	0
3	A	12	0	16	2	0
4	A	331	0	0	1	0
4	B	294	0	0	1	0
All	All	5537	0	4960	43	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (43) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:249[B]:VAL:HG21	1:B:256[B]:ILE:HD11	1.34	1.06
1:B:249[B]:VAL:CG2	1:B:256[B]:ILE:HD11	2.13	0.76
1:B:249[B]:VAL:HG21	1:B:256[B]:ILE:CD1	2.17	0.69
1:A:223[B]:GLU:HG2	1:B:206[B]:LYS:NZ	2.11	0.65
1:A:223[B]:GLU:CG	1:B:206[B]:LYS:HZ2	2.10	0.65
1:A:223[B]:GLU:HG2	1:B:206[B]:LYS:HZ2	1.64	0.63
1:A:3[B]:GLN:HG2	1:A:132[B]:SER:OG	2.03	0.58
1:A:195:TYR:CE2	1:A:211[B]:VAL:HG21	2.39	0.58
1:A:223[A]:GLU:HG2	1:B:294:ILE:HG13	1.85	0.58
1:B:249[B]:VAL:HG11	1:B:256[B]:ILE:CG1	2.34	0.58
1:A:242[A]:GLN:NE2	3:A:802:GOL:O3	2.33	0.57
1:B:195:TYR:CE2	1:B:211[B]:VAL:HG11	2.40	0.56
1:A:151:LYS:N	1:A:151:LYS:HD3	2.23	0.53
1:B:53:ASP:HB3	1:B:192:GLY:CA	2.42	0.50
1:A:64:TRP:O	1:B:77[A]:LYS:NZ	2.45	0.50
1:A:41:TYR:CZ	1:A:78:ALA:HB2	2.47	0.49
1:A:184:ILE:HB	1:A:211[B]:VAL:HG12	1.96	0.48
1:B:3:GLN:HG3	1:B:132[B]:SER:OG	2.15	0.47
1:A:223[B]:GLU:CG	1:B:206[B]:LYS:NZ	2.75	0.47
1:A:222[B]:ILE:HD13	1:B:297:LEU:HD11	1.96	0.46
1:B:53:ASP:HB3	1:B:192:GLY:HA2	1.97	0.46
1:B:92:THR:OG1	2:B:705:SO4:O4	2.32	0.46
1:A:53:ASP:HB3	1:A:192:GLY:CA	2.47	0.45
1:A:223[A]:GLU:HG3	1:B:294:ILE:HD12	1.97	0.45
1:A:155:ILE:HD11	4:A:1045:HOH:O	2.15	0.45
1:A:100:GLU:HG2	1:A:121[B]:MET:HE1	1.99	0.45
1:B:195:TYR:CZ	1:B:211[B]:VAL:HG11	2.52	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:223[A]:GLU:HG2	1:B:294:ILE:CD1	2.46	0.45
1:B:179:GLN:HE21	1:B:179:GLN:HA	1.82	0.45
1:A:53:ASP:HB3	1:A:192:GLY:HA2	1.99	0.44
1:A:195:TYR:CE1	3:A:801:GOL:H31	2.51	0.44
1:B:25:TYR:CD2	1:B:124:PRO:HD3	2.52	0.44
1:B:249[B]:VAL:HG11	1:B:256[B]:ILE:HG13	1.99	0.44
1:A:223[A]:GLU:CG	1:B:294:ILE:HD12	2.48	0.44
1:A:92[A]:THR:OG1	2:A:709:SO4:O2	2.33	0.43
1:A:223[A]:GLU:HG2	1:B:294:ILE:CG1	2.49	0.43
1:A:3[B]:GLN:CG	1:A:132[B]:SER:OG	2.68	0.42
1:A:198:LYS:HD3	1:A:211[A]:VAL:HG23	2.02	0.42
1:A:251:ILE:HG22	1:B:293[C]:ARG:NH2	2.35	0.42
1:B:155:ILE:HD11	4:B:1002:HOH:O	2.20	0.41
1:B:249[B]:VAL:HG11	1:B:256[B]:ILE:HD11	2.02	0.41
1:A:252:VAL:HG12	1:B:293[C]:ARG:NH2	2.36	0.41
1:A:195:TYR:CZ	1:A:211[B]:VAL:HG21	2.54	0.41

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	321/302 (106%)	314 (98%)	7 (2%)	0	100	100
1	B	323/302 (107%)	318 (98%)	5 (2%)	0	100	100
All	All	644/604 (107%)	632 (98%)	12 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	274/253 (108%)	262 (96%)	12 (4%)	25	1
1	B	275/253 (109%)	267 (97%)	8 (3%)	37	4
All	All	549/506 (108%)	529 (96%)	20 (4%)	36	2

All (20) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	28	GLU
1	A	65[A]	GLU
1	A	65[B]	GLU
1	A	106	ARG
1	A	121[A]	MET
1	A	121[B]	MET
1	A	151	LYS
1	A	220	ASN
1	A	226[A]	GLU
1	A	226[B]	GLU
1	A	256[A]	ILE
1	A	256[B]	ILE
1	B	77[A]	LYS
1	B	77[B]	LYS
1	B	176	GLU
1	B	179	GLN
1	B	220	ASN
1	B	239	LYS
1	B	271	ARG
1	B	299	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	63	ASN
1	B	3	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	6	GLN
1	B	179	GLN
1	B	242	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

10 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	SO4	B	708	-	4,4,4	0.26	0	6,6,6	0.21	0
2	SO4	A	706	-	4,4,4	0.23	0	6,6,6	0.21	0
2	SO4	A	704	-	4,4,4	0.23	0	6,6,6	0.12	0
3	GOL	A	802	-	5,5,5	0.35	0	5,5,5	0.29	0
2	SO4	B	701	-	4,4,4	0.24	0	6,6,6	0.14	0
2	SO4	B	705	-	4,4,4	0.21	0	6,6,6	0.19	0
2	SO4	A	709	-	4,4,4	0.26	0	6,6,6	0.16	0
2	SO4	A	702	-	4,4,4	0.23	0	6,6,6	0.09	0
2	SO4	B	707	-	4,4,4	0.22	0	6,6,6	0.12	0
3	GOL	A	801	-	5,5,5	0.38	0	5,5,5	0.28	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	A	802	-	-	0/4/4/4	-
3	GOL	A	801	-	-	2/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	801	GOL	O1-C1-C2-C3
3	A	801	GOL	O1-C1-C2-O2

There are no ring outliers.

4 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	802	GOL	1	0
2	B	705	SO4	1	0
2	A	709	SO4	1	0
3	A	801	GOL	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	301/302 (99%)	1.13	59 (19%) 3 8	6, 11, 17, 21	44 (14%)
1	B	300/302 (99%)	1.22	58 (19%) 3 8	5, 11, 18, 27	44 (14%)
All	All	601/604 (99%)	1.17	117 (19%) 3 8	5, 11, 18, 27	88 (14%)

All (117) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	29[A]	GLY	7.5
1	B	65[A]	GLU	6.7
1	B	298	LYS	6.5
1	B	296	ALA	6.1
1	B	299	ARG	5.8
1	B	66[A]	VAL	4.5
1	B	26	VAL	4.4
1	A	301	LEU	4.4
1	B	159	ILE	4.4
1	A	76	LEU	4.3
1	B	251	ILE	4.3
1	B	14	TYR	4.3
1	A	26	VAL	4.2
1	B	301	LEU	4.2
1	B	78	ALA	4.2
1	A	66[A]	VAL	4.1
1	A	296	ALA	4.1
1	A	65[A]	GLU	4.0
1	A	253	GLY	3.8
1	A	23	THR	3.8
1	A	27	VAL	3.8
1	A	252	VAL	3.8
1	B	28[A]	GLU	3.7
1	A	157	VAL	3.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	31	VAL	3.6
1	B	294	ILE	3.6
1	B	297	LEU	3.5
1	A	254	GLY	3.5
1	B	290	ALA	3.5
1	B	291	THR	3.5
1	A	251	ILE	3.5
1	B	289	LEU	3.4
1	B	20	ASP	3.3
1	B	157	VAL	3.2
1	A	302	GLN	3.2
1	A	77[A]	LYS	3.2
1	A	19	VAL	3.1
1	B	123	VAL	3.1
1	B	295	ASP	3.1
1	A	159	ILE	3.0
1	A	290	ALA	3.0
1	B	252	VAL	2.9
1	A	119	ALA	2.9
1	B	76	LEU	2.9
1	A	287	ASN	2.9
1	A	99	VAL	2.9
1	A	297	LEU	2.8
1	B	155	ILE	2.8
1	A	162	GLY	2.8
1	A	299	ARG	2.8
1	B	62	TRP	2.8
1	B	2	SER	2.8
1	A	152	VAL	2.8
1	B	63	ASN	2.7
1	A	28	GLU	2.7
1	A	63	ASN	2.6
1	B	27	VAL	2.6
1	B	32	VAL	2.6
1	A	64	TRP	2.6
1	B	64	TRP	2.6
1	A	291	THR	2.6
1	A	102	ALA	2.6
1	A	18	ARG	2.6
1	B	292	PRO	2.6
1	B	36	ALA	2.5
1	B	122	GLY	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	107	ILE	2.5
1	B	79	ARG	2.5
1	B	120	GLN	2.5
1	A	21	PHE	2.5
1	B	22	PRO	2.4
1	A	255[A]	SER	2.4
1	B	31	VAL	2.4
1	B	3	GLN	2.4
1	A	289	LEU	2.4
1	B	156[A]	ASP	2.4
1	A	60	LEU	2.4
1	A	170[A]	LEU	2.4
1	A	122	GLY	2.4
1	B	119	ALA	2.3
1	B	257	TYR	2.3
1	A	37[A]	MET	2.3
1	A	300[A]	ARG	2.3
1	A	107	ILE	2.3
1	A	106	ARG	2.3
1	A	169	ALA	2.3
1	B	75	PHE	2.3
1	B	253	GLY	2.3
1	A	108[A]	PRO	2.3
1	A	25	TYR	2.2
1	B	300	ARG	2.2
1	A	29	GLY	2.2
1	A	154	GLY	2.2
1	A	33	ARG	2.2
1	A	155	ILE	2.2
1	B	150	ALA	2.2
1	B	124	PRO	2.2
1	B	77[A]	LYS	2.2
1	A	32	VAL	2.2
1	B	249[A]	VAL	2.2
1	B	23	THR	2.1
1	B	247	ALA	2.1
1	A	148	VAL	2.1
1	A	62	TRP	2.1
1	A	97	TYR	2.1
1	B	81	GLY	2.1
1	A	79	ARG	2.1
1	B	15	ILE	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	161	GLU	2.1
1	B	148	VAL	2.1
1	A	20	ASP	2.0
1	A	256[A]	ILE	2.0
1	A	17	ARG	2.0
1	A	286	VAL	2.0
1	B	19	VAL	2.0
1	B	293[B]	ARG	2.0
1	A	24	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	GOL	A	801	6/6	0.64	0.21	25,26,27,27	6
2	SO4	B	708	5/5	0.73	0.17	17,18,19,20	5
2	SO4	A	702	5/5	0.74	0.20	36,36,36,36	5
2	SO4	A	704	5/5	0.74	0.21	32,32,32,33	5
2	SO4	B	701	5/5	0.87	0.13	31,31,31,31	5
2	SO4	B	707	5/5	0.87	0.13	21,21,22,22	5
2	SO4	B	705	5/5	0.88	0.13	13,14,16,16	5
3	GOL	A	802	6/6	0.88	0.12	18,19,19,19	6
2	SO4	A	709	5/5	0.94	0.10	6,8,9,9	5
2	SO4	A	706	5/5	0.95	0.09	17,18,19,19	5

6.5 Other polymers [i](#)

There are no such residues in this entry.