



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 8, 2026 – 01:36 PM UTC

PDB ID : 1TZN / pdb_00001tzn
Title : Crystal Structure of the Anthrax Toxin Protective Antigen Heptameric Pre-pore bound to the VWA domain of CMG2, an anthrax toxin receptor
Authors : Lacy, D.B.; Wigelsworth, D.J.; Melnyk, R.A.; Collier, R.J.
Deposited on : 2004-07-10
Resolution : 4.30 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 4.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1052 (4.70-3.90)
Clashscore	190562	1097 (4.70-3.90)
Ramachandran outliers	187476	1001 (4.70-3.90)
RSRZ outliers	180081	1049 (4.70-3.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	562	
1	B	562	
1	C	562	
1	D	562	
1	E	562	
1	F	562	
1	G	562	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	H	562	
1	I	562	
1	J	562	
1	K	562	
1	L	562	
1	M	562	
1	O	562	
2	a	181	
2	b	181	
2	c	181	
2	d	181	
2	e	181	
2	f	181	
2	g	181	
2	h	181	
2	i	181	
2	j	181	
2	k	181	
2	l	181	
2	m	181	
2	o	181	

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 80570 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Protective antigen.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	552	Total	C	N	O	S	0	0	0
			4356	2726	752	872	6			
1	B	552	Total	C	N	O	S	0	0	0
			4356	2726	752	872	6			
1	C	552	Total	C	N	O	S	0	0	0
			4356	2726	752	872	6			
1	D	552	Total	C	N	O	S	0	0	0
			4356	2726	752	872	6			
1	E	552	Total	C	N	O	S	0	0	0
			4356	2726	752	872	6			
1	F	552	Total	C	N	O	S	0	0	0
			4356	2726	752	872	6			
1	G	552	Total	C	N	O	S	0	0	0
			4356	2726	752	872	6			
1	H	552	Total	C	N	O	S	0	0	0
			4356	2726	752	872	6			
1	I	552	Total	C	N	O	S	0	0	0
			4356	2726	752	872	6			
1	J	552	Total	C	N	O	S	0	0	0
			4356	2726	752	872	6			
1	K	552	Total	C	N	O	S	0	0	0
			4356	2726	752	872	6			
1	L	552	Total	C	N	O	S	0	0	0
			4356	2726	752	872	6			
1	M	552	Total	C	N	O	S	0	0	0
			4356	2726	752	872	6			
1	O	552	Total	C	N	O	S	0	0	0
			4356	2726	752	872	6			

- Molecule 2 is a protein called Anthrax toxin receptor 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	a	181	Total	C	N	O	S	0	0	0
			1396	891	236	265	4			
2	b	181	Total	C	N	O	S	0	0	0
			1396	891	236	265	4			
2	c	181	Total	C	N	O	S	0	0	0
			1396	891	236	265	4			
2	d	181	Total	C	N	O	S	0	0	0
			1396	891	236	265	4			
2	e	181	Total	C	N	O	S	0	0	0
			1396	891	236	265	4			
2	f	181	Total	C	N	O	S	0	0	0
			1396	891	236	265	4			
2	g	181	Total	C	N	O	S	0	0	0
			1396	891	236	265	4			
2	h	181	Total	C	N	O	S	0	0	0
			1396	891	236	265	4			
2	i	181	Total	C	N	O	S	0	0	0
			1396	891	236	265	4			
2	j	181	Total	C	N	O	S	0	0	0
			1396	891	236	265	4			
2	k	181	Total	C	N	O	S	0	0	0
			1396	891	236	265	4			
2	l	181	Total	C	N	O	S	0	0	0
			1396	891	236	265	4			
2	m	181	Total	C	N	O	S	0	0	0
			1396	891	236	265	4			
2	o	181	Total	C	N	O	S	0	0	0
			1396	891	236	265	4			

- Molecule 3 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	2	Total	Ca	0	0
			2	2		
3	B	2	Total	Ca	0	0
			2	2		
3	C	2	Total	Ca	0	0
			2	2		
3	D	2	Total	Ca	0	0
			2	2		
3	E	2	Total	Ca	0	0
			2	2		
3	F	2	Total	Ca	0	0
			2	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	G	2	Total 2	Ca 2	0	0
3	H	2	Total 2	Ca 2	0	0
3	I	2	Total 2	Ca 2	0	0
3	J	2	Total 2	Ca 2	0	0
3	K	2	Total 2	Ca 2	0	0
3	L	2	Total 2	Ca 2	0	0
3	M	2	Total 2	Ca 2	0	0
3	O	2	Total 2	Ca 2	0	0

- Molecule 4 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total 1	Mg 1	0	0
4	B	1	Total 1	Mg 1	0	0
4	C	1	Total 1	Mg 1	0	0
4	D	1	Total 1	Mg 1	0	0
4	E	1	Total 1	Mg 1	0	0
4	F	1	Total 1	Mg 1	0	0
4	G	1	Total 1	Mg 1	0	0
4	H	1	Total 1	Mg 1	0	0
4	I	1	Total 1	Mg 1	0	0
4	J	1	Total 1	Mg 1	0	0
4	K	1	Total 1	Mg 1	0	0

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	L	1	Total 1	Mg 1	0	0
4	M	1	Total 1	Mg 1	0	0
4	O	1	Total 1	Mg 1	0	0

3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Protective antigen





• Molecule 1: Protective antigen






• Molecule 1: Protective antigen







• Molecule 1: Protective antigen




• Molecule 1: Protective antigen




• Molecule 1: Protective antigen








• Molecule 1: Protective antigen




• Molecule 1: Protective antigen








• Molecule 1: Protective antigen





• Molecule 1: Protective antigen







• Molecule 1: Protective antigen








• Molecule 2: Anthrax toxin receptor 2




• Molecule 2: Anthrax toxin receptor 2




• Molecule 2: Anthrax toxin receptor 2




• Molecule 2: Anthrax toxin receptor 2



• Molecule 2: Anthrax toxin receptor 2




• Molecule 2: Anthrax toxin receptor 2




• Molecule 2: Anthrax toxin receptor 2







• Molecule 2: Anthrax toxin receptor 2




• Molecule 2: Anthrax toxin receptor 2




• Molecule 2: Anthrax toxin receptor 2







• Molecule 2: Anthrax toxin receptor 2



• Molecule 2: Anthrax toxin receptor 2





• Molecule 2: Anthrax toxin receptor 2





• Molecule 2: Anthrax toxin receptor 2



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	158.80Å 158.85Å 214.08Å 69.58° 69.07° 65.58°	Depositor
Resolution (Å)	20.00 – 4.30 20.00 – 4.30	Depositor EDS
% Data completeness (in resolution range)	97.2 (20.00-4.30) 96.2 (20.00-4.30)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.17	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.97 (at 4.20Å)	Xtriage
Refinement program	CNS	Depositor
R, R_{free}	0.322 , 0.330 0.305 , 0.310	Depositor DCC
R_{free} test set	5704 reflections (4.76%)	wwPDB-VP
Wilson B-factor (Å ²)	81.4	Xtriage
Anisotropy	0.270	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.20 , 71.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.44$, $\langle L^2 \rangle = 0.26$	Xtriage
Estimated twinning fraction	0.125 for -k,-h,-l	Xtriage
F_o, F_c correlation	0.77	EDS
Total number of atoms	80570	wwPDB-VP
Average B, all atoms (Å ²)	65.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.22% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CA, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.32	0/4430	0.78	7/6007 (0.1%)
1	B	0.32	0/4430	0.78	7/6007 (0.1%)
1	C	0.32	0/4430	0.78	7/6007 (0.1%)
1	D	0.32	0/4430	0.78	7/6007 (0.1%)
1	E	0.32	0/4430	0.78	7/6007 (0.1%)
1	F	0.32	0/4430	0.78	7/6007 (0.1%)
1	G	0.32	0/4430	0.78	7/6007 (0.1%)
1	H	0.32	0/4430	0.78	7/6007 (0.1%)
1	I	0.32	0/4430	0.78	7/6007 (0.1%)
1	J	0.32	0/4430	0.78	7/6007 (0.1%)
1	K	0.32	0/4430	0.78	7/6007 (0.1%)
1	L	0.32	0/4430	0.78	7/6007 (0.1%)
1	M	0.32	0/4430	0.78	7/6007 (0.1%)
1	O	0.32	0/4430	0.78	7/6007 (0.1%)
2	a	0.27	0/1417	0.80	1/1907 (0.1%)
2	b	0.27	0/1417	0.80	1/1907 (0.1%)
2	c	0.27	0/1417	0.80	1/1907 (0.1%)
2	d	0.27	0/1417	0.80	1/1907 (0.1%)
2	e	0.27	0/1417	0.80	1/1907 (0.1%)
2	f	0.27	0/1417	0.80	1/1907 (0.1%)
2	g	0.27	0/1417	0.80	1/1907 (0.1%)
2	h	0.27	0/1417	0.80	1/1907 (0.1%)
2	i	0.27	0/1417	0.80	1/1907 (0.1%)
2	j	0.27	0/1417	0.80	1/1907 (0.1%)
2	k	0.27	0/1417	0.80	1/1907 (0.1%)
2	l	0.27	0/1417	0.80	1/1907 (0.1%)
2	m	0.27	0/1417	0.80	1/1907 (0.1%)
2	o	0.27	0/1417	0.80	1/1907 (0.1%)
All	All	0.31	0/81858	0.79	112/110796 (0.1%)

There are no bond length outliers.

The worst 5 of 112 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	J	424	GLN	OE1-CD-NE2	-9.87	112.73	122.60
1	G	424	GLN	OE1-CD-NE2	-9.86	112.74	122.60
1	E	424	GLN	OE1-CD-NE2	-9.85	112.75	122.60
1	B	424	GLN	OE1-CD-NE2	-9.85	112.75	122.60
1	C	424	GLN	OE1-CD-NE2	-9.84	112.76	122.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4356	0	4314	442	5
1	B	4356	0	4314	438	0
1	C	4356	0	4314	430	2
1	D	4356	0	4314	426	2
1	E	4356	0	4314	419	2
1	F	4356	0	4314	417	9
1	G	4356	0	4314	474	6
1	H	4356	0	4314	483	5
1	I	4356	0	4314	478	0
1	J	4356	0	4314	444	1
1	K	4356	0	4314	451	2
1	L	4356	0	4314	460	0
1	M	4356	0	4314	474	8
1	O	4356	0	4314	447	2
2	a	1396	0	1426	122	2
2	b	1396	0	1426	121	2
2	c	1396	0	1426	120	0
2	d	1396	0	1426	118	0
2	e	1396	0	1426	120	5
2	f	1396	0	1426	117	0
2	g	1396	0	1426	119	0
2	h	1396	0	1426	119	1
2	i	1396	0	1426	120	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	j	1396	0	1426	120	0
2	k	1396	0	1426	117	5
2	l	1396	0	1426	120	4
2	m	1396	0	1426	117	1
2	o	1396	0	1426	119	0
3	A	2	0	0	0	0
3	B	2	0	0	0	0
3	C	2	0	0	0	0
3	D	2	0	0	0	0
3	E	2	0	0	0	0
3	F	2	0	0	0	0
3	G	2	0	0	0	0
3	H	2	0	0	0	0
3	I	2	0	0	0	0
3	J	2	0	0	0	0
3	K	2	0	0	0	0
3	L	2	0	0	0	0
3	M	2	0	0	0	0
3	O	2	0	0	0	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
4	C	1	0	0	0	0
4	D	1	0	0	0	0
4	E	1	0	0	0	0
4	F	1	0	0	0	0
4	G	1	0	0	0	0
4	H	1	0	0	0	0
4	I	1	0	0	0	0
4	J	1	0	0	0	0
4	K	1	0	0	0	0
4	L	1	0	0	0	0
4	M	1	0	0	0	0
4	O	1	0	0	0	0
All	All	80570	0	80360	7282	32

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 45.

The worst 5 of 7282 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:G:414:LYS:HG3	1:M:319:SER:N	1.35	1.39
1:G:305:GLY:HA2	1:H:670:GLN:NE2	1.42	1.34
1:G:414:LYS:CG	1:M:319:SER:H	1.41	1.32
1:H:483:GLN:NE2	1:I:469:VAL:HG21	1.47	1.28
1:L:305:GLY:HA2	1:M:670:GLN:CD	1.58	1.26

The worst 5 of 32 symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:H:546:ASP:OD2	2:l:1203:PHE:CE1[1_455]	0.85	1.35
1:F:570:ASN:CG	1:G:197:LYS:NZ[1_544]	1.33	0.87
2:b:1067:GLN:OE1	1:E:537:ASN:OD1[1_455]	1.36	0.84
1:A:197:LYS:NZ	1:M:570:ASN:ND2[1_544]	1.37	0.83
1:A:197:LYS:NZ	1:M:570:ASN:CG[1_544]	1.45	0.75

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	546/562 (97%)	374 (68%)	132 (24%)	40 (7%)	1	11
1	B	546/562 (97%)	374 (68%)	132 (24%)	40 (7%)	1	11
1	C	546/562 (97%)	373 (68%)	133 (24%)	40 (7%)	1	11
1	D	546/562 (97%)	374 (68%)	132 (24%)	40 (7%)	1	11
1	E	546/562 (97%)	374 (68%)	132 (24%)	40 (7%)	1	11
1	F	546/562 (97%)	373 (68%)	133 (24%)	40 (7%)	1	11
1	G	546/562 (97%)	374 (68%)	132 (24%)	40 (7%)	1	11
1	H	546/562 (97%)	374 (68%)	132 (24%)	40 (7%)	1	11
1	I	546/562 (97%)	374 (68%)	132 (24%)	40 (7%)	1	11
1	J	546/562 (97%)	374 (68%)	132 (24%)	40 (7%)	1	11

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	K	546/562 (97%)	374 (68%)	132 (24%)	40 (7%)	1	11
1	L	546/562 (97%)	374 (68%)	132 (24%)	40 (7%)	1	11
1	M	546/562 (97%)	373 (68%)	133 (24%)	40 (7%)	1	11
1	O	546/562 (97%)	374 (68%)	132 (24%)	40 (7%)	1	11
2	a	179/181 (99%)	144 (80%)	29 (16%)	6 (3%)	3	21
2	b	179/181 (99%)	144 (80%)	29 (16%)	6 (3%)	3	21
2	c	179/181 (99%)	143 (80%)	30 (17%)	6 (3%)	3	21
2	d	179/181 (99%)	144 (80%)	29 (16%)	6 (3%)	3	21
2	e	179/181 (99%)	144 (80%)	29 (16%)	6 (3%)	3	21
2	f	179/181 (99%)	144 (80%)	29 (16%)	6 (3%)	3	21
2	g	179/181 (99%)	144 (80%)	29 (16%)	6 (3%)	3	21
2	h	179/181 (99%)	144 (80%)	29 (16%)	6 (3%)	3	21
2	i	179/181 (99%)	144 (80%)	29 (16%)	6 (3%)	3	21
2	j	179/181 (99%)	144 (80%)	29 (16%)	6 (3%)	3	21
2	k	179/181 (99%)	144 (80%)	29 (16%)	6 (3%)	3	21
2	l	179/181 (99%)	144 (80%)	29 (16%)	6 (3%)	3	21
2	m	179/181 (99%)	144 (80%)	29 (16%)	6 (3%)	3	21
2	o	179/181 (99%)	144 (80%)	29 (16%)	6 (3%)	3	21
All	All	10150/10402 (98%)	7248 (71%)	2258 (22%)	644 (6%)	1	13

5 of 644 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	197	LYS
1	A	209	ASN
1	A	422	ASN
1	A	656	ILE
2	a	1086	SER

5.3.2 Protein sidechains ⓘ

There are no protein residues with a non-rotameric sidechain to report in this entry.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 42 ligands modelled in this entry, 42 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å²)	Q<0.9	
1	A	552/562 (98%)	0.81	45 (8%)	17	21	1, 36, 166, 200	0
1	B	552/562 (98%)	0.70	32 (5%)	29	28	1, 36, 166, 200	0
1	C	552/562 (98%)	0.97	69 (12%)	8	12	1, 36, 166, 200	0
1	D	552/562 (98%)	0.71	41 (7%)	20	23	1, 36, 166, 200	0
1	E	552/562 (98%)	0.72	29 (5%)	32	30	1, 36, 166, 200	0
1	F	552/562 (98%)	0.79	39 (7%)	22	23	1, 36, 166, 200	0
1	G	552/562 (98%)	0.80	48 (8%)	16	19	1, 36, 166, 200	0
1	H	552/562 (98%)	0.80	43 (7%)	19	21	1, 36, 166, 200	0
1	I	552/562 (98%)	0.71	44 (7%)	18	21	1, 36, 166, 200	0
1	J	552/562 (98%)	0.83	45 (8%)	17	21	1, 36, 166, 200	0
1	K	552/562 (98%)	0.84	58 (10%)	11	15	1, 36, 166, 200	0
1	L	552/562 (98%)	0.80	48 (8%)	16	19	1, 36, 166, 200	0
1	M	552/562 (98%)	0.82	40 (7%)	21	23	1, 36, 166, 200	0
1	O	552/562 (98%)	0.77	48 (8%)	16	19	1, 36, 166, 200	0
2	a	181/181 (100%)	1.50	54 (29%)	1	4	6, 92, 179, 200	0
2	b	181/181 (100%)	0.91	22 (12%)	8	12	6, 92, 179, 200	0
2	c	181/181 (100%)	1.55	50 (27%)	1	4	6, 92, 179, 200	0
2	d	181/181 (100%)	1.10	24 (13%)	7	11	6, 92, 179, 200	0
2	e	181/181 (100%)	1.18	24 (13%)	7	11	6, 92, 179, 200	0
2	f	181/181 (100%)	1.21	21 (11%)	9	13	6, 92, 179, 200	0
2	g	181/181 (100%)	1.24	33 (18%)	3	7	6, 92, 179, 200	0
2	h	181/181 (100%)	1.14	29 (16%)	5	9	6, 92, 179, 200	0
2	i	181/181 (100%)	1.08	26 (14%)	6	10	6, 92, 179, 200	0
2	j	181/181 (100%)	1.87	67 (37%)	1	3	6, 92, 179, 200	0

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
2	k	181/181 (100%)	1.36	45 (24%) 2 5	6, 92, 179, 200	0
2	l	181/181 (100%)	1.74	58 (32%) 1 3	6, 92, 179, 200	0
2	m	181/181 (100%)	0.99	17 (9%) 14 18	6, 92, 179, 200	0
2	o	181/181 (100%)	1.25	39 (21%) 2 6	6, 92, 179, 200	0
All	All	10262/10402 (98%)	0.92	1138 (11%) 10 14	1, 54, 174, 200	0

The worst 5 of 1138 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	M	319	SER	12.2
1	E	319	SER	9.1
1	J	424	GLN	7.8
1	I	424	GLN	7.8
1	J	319	SER	7.8

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	MG	F	738	1/1	0.50	0.31	0,0,0,0	0
4	MG	L	738	1/1	0.89	0.27	0,0,0,0	0
4	MG	B	738	1/1	0.90	0.11	0,0,0,0	0
4	MG	D	738	1/1	0.90	0.09	0,0,0,0	0
4	MG	J	738	1/1	0.91	0.12	0,0,0,0	0
3	CA	G	737	1/1	0.94	0.11	0,0,0,0	0
4	MG	E	738	1/1	0.96	0.15	0,0,0,0	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	CA	F	737	1/1	0.96	0.15	0,0,0,0	0
3	CA	F	736	1/1	0.97	0.05	0,0,0,0	0
3	CA	A	737	1/1	0.97	0.11	0,0,0,0	0
3	CA	B	737	1/1	0.97	0.17	0,0,0,0	0
3	CA	J	737	1/1	0.97	0.18	0,0,0,0	0
4	MG	G	738	1/1	0.97	0.14	0,0,0,0	0
4	MG	H	738	1/1	0.97	0.12	0,0,0,0	0
3	CA	M	737	1/1	0.97	0.15	0,0,0,0	0
4	MG	A	738	1/1	0.97	0.14	0,0,0,0	0
3	CA	E	737	1/1	0.98	0.18	0,0,0,0	0
3	CA	H	737	1/1	0.98	0.07	0,0,0,0	0
4	MG	I	738	1/1	0.98	0.08	0,0,0,0	0
3	CA	O	737	1/1	0.98	0.07	0,0,0,0	0
3	CA	I	736	1/1	0.98	0.11	0,0,0,0	0
4	MG	M	738	1/1	0.98	0.15	0,0,0,0	0
3	CA	C	737	1/1	0.99	0.10	0,0,0,0	0
3	CA	G	736	1/1	0.99	0.06	0,0,0,0	0
3	CA	D	737	1/1	0.99	0.05	0,0,0,0	0
3	CA	H	736	1/1	0.99	0.06	0,0,0,0	0
4	MG	C	738	1/1	0.99	0.05	0,0,0,0	0
3	CA	E	736	1/1	0.99	0.07	0,0,0,0	0
3	CA	B	736	1/1	0.99	0.05	0,0,0,0	0
3	CA	I	737	1/1	0.99	0.09	0,0,0,0	0
3	CA	J	736	1/1	0.99	0.05	0,0,0,0	0
3	CA	C	736	1/1	0.99	0.06	0,0,0,0	0
3	CA	K	736	1/1	0.99	0.04	0,0,0,0	0
3	CA	K	737	1/1	0.99	0.05	0,0,0,0	0
4	MG	K	738	1/1	0.99	0.12	0,0,0,0	0
3	CA	L	736	1/1	0.99	0.05	0,0,0,0	0
3	CA	L	737	1/1	0.99	0.11	0,0,0,0	0
3	CA	D	736	1/1	1.00	0.05	0,0,0,0	0
3	CA	O	736	1/1	1.00	0.03	0,0,0,0	0
3	CA	A	736	1/1	1.00	0.01	0,0,0,0	0
3	CA	M	736	1/1	1.00	0.02	0,0,0,0	0
4	MG	O	738	1/1	1.00	0.09	0,0,0,0	0

6.5 Other polymers

There are no such residues in this entry.