



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 09:48 PM UTC

PDB ID : 4TZ0 / pdb_00004tz0
Title : DEAD-box helicase Mss116 bound to ssRNA and GDP-BeF
Authors : Mallam, A.L.; Sidote, D.J.; Lambowitz, A.M.
Deposited on : 2014-07-09
Resolution : 2.35 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.35 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1596 (2.36-2.36)
Clashscore	190562	1663 (2.36-2.36)
Ramachandran outliers	187476	1646 (2.36-2.36)
Sidechain outliers	187428	1646 (2.36-2.36)
RSRZ outliers	180081	1598 (2.36-2.36)
RNA backbone	3983	1027 (2.66-2.06)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	509	22% 94% 6%
2	B	7	29% 100%

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 8057 atoms, of which 3927 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called ATP-dependent RNA helicase MSS116, mitochondrial.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	508	Total	C	H	N	O	S	4	0	0
			7723	2486	3838	653	732	14			

- Molecule 2 is a RNA chain called RNA (5'-R(*AP*AP*AP*AP*AP*AP*A)-3').


Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	B	7	Total	C	H	N	O	P	0	0	0
			230	70	79	35	40	6			

- Molecule 3 is GUANOSINE-5'-DIPHOSPHATE (CCD ID: GDP) (formula: C₁₀H₁₅N₅O₁₁P₂).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
3	A	1	Total	C	H	N	O	P	0	0
			38	10	10	5	11	2		

- Molecule 4 is BERYLLIUM TRIFLUORIDE ION (CCD ID: BEF) (formula: BeF₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	Be	F	0	0
			4	1	3		

- Molecule 5 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	Mg	0	0
			1	1		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	58	Total	O	0	0
			58	58		
6	B	3	Total	O	0	0
			3	3		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: ATP-dependent RNA helicase MSS116, mitochondrial



- Molecule 2: RNA (5'-R(*AP*AP*AP*AP*AP*AP*A)-3')



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	89.98Å 126.61Å 55.55Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.27 – 2.35 47.27 – 2.35	Depositor EDS
% Data completeness (in resolution range)	99.9 (47.27-2.35) 99.6 (47.27-2.35)	Depositor EDS
R_{merge}	0.13	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.60 (at 2.34Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8.4_1496)	Depositor
R, R_{free}	0.231 , 0.260 0.242 , 0.268	Depositor DCC
R_{free} test set	1363 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	44.6	Xtriage
Anisotropy	0.835	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 71.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	8057	wwPDB-VP
Average B, all atoms (Å ²)	69.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.84% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MG, BEF, GDP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.30	0/3950	0.65	0/5348
2	B	0.09	0/171	0.29	0/265
All	All	0.29	0/4121	0.64	0/5613

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3885	3838	3847	11	0
2	B	151	79	79	0	0
3	A	28	10	12	0	0
4	A	4	0	0	0	0
5	A	1	0	0	0	0
6	A	58	0	0	0	0
6	B	3	0	0	0	0
All	All	4130	3927	3938	11	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (11) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:341:ARG:NH1	1:A:472:LYS:O	2.22	0.73
1:A:383:VAL:O	1:A:386:THR:OG1	2.25	0.49
1:A:173:THR:O	1:A:182:LYS:NZ	2.45	0.49
1:A:572:VAL:HG21	1:A:592:PHE:CD1	2.49	0.47
1:A:371:ASP:O	1:A:372:SER:OG	2.30	0.46
1:A:522:GLU:OE1	1:A:522:GLU:N	2.45	0.46
1:A:164:ILE:HB	1:A:165:PRO:HD3	1.98	0.45
1:A:485:LEU:N	1:A:486:PRO:CD	2.80	0.45
1:A:144:GLU:CB	1:A:145:ASP:HA	2.48	0.43
1:A:109:GLU:C	1:A:111:GLY:H	2.28	0.41
1:A:142:SER:O	1:A:300:LYS:NZ	2.54	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	506/509 (99%)	490 (97%)	16 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	411/458 (90%)	402 (98%)	9 (2%)	45 59

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	163	LEU
1	A	170	LEU
1	A	194	LEU
1	A	256	LYS
1	A	290	GLU
1	A	302	LEU
1	A	308	LEU
1	A	366	GLN
1	A	466	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	178	GLN
1	A	204	HIS
1	A	362	HIS
1	A	444	ASN

5.3.3 RNA ⓘ

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
2	B	6/7 (85%)	0	0

There are no RNA backbone outliers to report.

There are no RNA pucker outliers to report.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

Of 3 ligands modelled in this entry, 1 is monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	BEF	A	602	-	0,3,3	-	-	-		
3	GDP	A	601	5	29,30,30	1.14	2 (6%)	45,47,47	1.79	7 (15%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GDP	A	601	5	-	5/16/32/32	0/3/3/3

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	601	GDP	C5-C4	3.11	1.47	1.38
3	A	601	GDP	C6-N1	-2.36	1.34	1.38

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	601	GDP	C5-C4-N3	-6.18	118.56	128.39
3	A	601	GDP	C2-N3-C4	5.08	121.05	112.30
3	A	601	GDP	N9-C4-N3	4.40	134.75	125.95
3	A	601	GDP	C6-C5-N7	3.43	136.53	130.29
3	A	601	GDP	C4-C5-N7	-2.76	106.29	110.67
3	A	601	GDP	O6-C6-C5	-2.18	120.77	126.53
3	A	601	GDP	C8-N7-C5	2.01	107.84	104.26

There are no chirality outliers.


All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	601	GDP	C5'-O5'-PA-O3A
3	A	601	GDP	C5'-O5'-PA-O1A
3	A	601	GDP	C5'-O5'-PA-O2A
3	A	601	GDP	O4'-C4'-C5'-O5'
3	A	601	GDP	C3'-C4'-C5'-O5'

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	508/509 (99%)	1.23	111 (21%) 2 2	37, 66, 108, 142	1 (0%)
2	B	7/7 (100%)	0.94	2 (28%) 1 1	45, 50, 116, 125	0
All	All	515/516 (99%)	1.23	113 (21%) 2 2	37, 66, 109, 142	1 (0%)

All (113) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	91	ILE	5.4
1	A	482	LYS	5.3
1	A	308	LEU	5.0
1	A	139	PRO	5.0
1	A	92	HIS	4.9
1	A	334	ASN	4.8
1	A	90	LEU	4.7
1	A	310	ASP	4.5
1	A	98	ASN	4.5
1	A	518	ALA	4.3
1	A	328	LEU	4.3
1	A	330	THR	4.3
1	A	326	LEU	4.3
1	A	576	PHE	4.3
1	A	327	PHE	4.1
1	A	88	SER	4.0
1	A	146	HIS	3.9
1	A	93	VAL	3.8
1	A	487	PHE	3.7
1	A	520	THR	3.7
1	A	333	LYS	3.7
1	A	572	VAL	3.6
1	A	117	ILE	3.6
1	A	499	ILE	3.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	143	SER	3.5
1	A	348	ILE	3.5
1	A	148	VAL	3.4
1	A	570	ILE	3.4
1	A	149	ILE	3.3
1	A	175	PHE	3.3
1	A	582	LEU	3.2
1	A	544	TYR	3.1
1	A	219	VAL	3.1
1	A	151	ARG	3.1
2	B	1	A	3.1
1	A	144	GLU	3.0
1	A	156	THR	3.0
1	A	594	ILE	3.0
1	A	312	VAL	3.0
1	A	331	VAL	3.0
1	A	145	ASP	3.0
1	A	113	LEU	3.0
1	A	519	VAL	3.0
1	A	546	PHE	2.9
1	A	325	CYS	2.9
1	A	573	SER	2.9
1	A	336	PRO	2.9
1	A	109	GLU	2.9
1	A	578	ASP	2.9
1	A	577	LEU	2.8
1	A	111	GLY	2.8
1	A	491	LEU	2.7
2	B	7	A	2.7
1	A	273	LEU	2.7
1	A	595	ARG	2.7
1	A	94	PRO	2.7
1	A	127	PRO	2.7
1	A	138	LYS	2.7
1	A	249	VAL	2.7
1	A	453	VAL	2.7
1	A	579	LYS	2.7
1	A	575	ARG	2.7
1	A	500	ALA	2.6
1	A	498	VAL	2.6
1	A	574	ARG	2.6
1	A	551	ILE	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	141	LEU	2.6
1	A	497	ILE	2.6
1	A	150	ALA	2.5
1	A	354	ASN	2.5
1	A	254	SER	2.5
1	A	461	ILE	2.5
1	A	324	GLU	2.5
1	A	350	GLU	2.5
1	A	315	LEU	2.5
1	A	293	SER	2.5
1	A	515	VAL	2.5
1	A	418	LEU	2.4
1	A	516	LEU	2.4
1	A	337	GLU	2.4
1	A	496	ASN	2.3
1	A	95	LYS	2.3
1	A	342	ILE	2.3
1	A	112	VAL	2.3
1	A	485	LEU	2.3
1	A	586	PRO	2.3
1	A	590	ALA	2.3
1	A	547	SER	2.3
1	A	521	GLU	2.3
1	A	108	LEU	2.3
1	A	473	GLU	2.3
1	A	460	TYR	2.2
1	A	592	PHE	2.2
1	A	480	ILE	2.2
1	A	369	GLU	2.2
1	A	478	LEU	2.1
1	A	457	LEU	2.1
1	A	372	SER	2.1
1	A	532	SER	2.1
1	A	311	LYS	2.1
1	A	343	ASP	2.1
1	A	155	GLY	2.1
1	A	581	GLY	2.1
1	A	153	LYS	2.0
1	A	353	ALA	2.0
1	A	396	GLU	2.0
1	A	526	ILE	2.0
1	A	340	GLU	2.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	455	SER	2.0
1	A	395	ASN	2.0
1	A	137	ILE	2.0
1	A	489	ARG	2.0
1	A	531	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	GDP	A	601	28/28	0.90	0.12	1,63,84,95	0
4	BEF	A	602	4/4	0.96	0.08	44,45,46,48	0
5	MG	A	603	1/1	0.99	0.05	46,46,46,46	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around GDP A 601:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.