



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 27, 2026 – 10:56 PM UTC

PDB ID : 4TZE / pdb_00004tze
Title : Structure of metallo-beta-lactamase
Authors : Ferguson, J.A.; Makena, A.; Brem, J.; McDonough, M.A.; Schofield, C.J.
Deposited on : 2014-07-10
Resolution : 1.57 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.57 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1094 (1.58-1.58)
Clashscore	190562	1105 (1.58-1.58)
Ramachandran outliers	187476	1082 (1.58-1.58)
Sidechain outliers	187428	1081 (1.58-1.58)
RSRZ outliers	180081	1094 (1.58-1.58)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	231	15% 94% 6%
1	B	231	18% 95% • •

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 3704 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Class B carbapenemase NDM-5.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	230	Total	C	N	O	S	0	9	0
			1732	1092	301	329	10			
1	B	229	Total	C	N	O	S	0	3	0
			1699	1073	295	322	9			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	40	GLY	-	expression tag	UNP G3K399
A	41	PRO	-	expression tag	UNP G3K399
B	40	GLY	-	expression tag	UNP G3K399
B	41	PRO	-	expression tag	UNP G3K399

- Molecule 2 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total	Zn	0	0
			2	2		
2	B	2	Total	Zn	0	0
			2	2		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	131	Total	O	0	0
			131	131		
3	B	138	Total	O	0	0
			138	138		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Class B carbapenemase NDM-5



- Molecule 1: Class B carbapenemase NDM-5



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	53.77Å 59.22Å 62.97Å 90.00° 92.19° 90.00°	Depositor
Resolution (Å)	34.07 – 1.57 34.07 – 1.57	Depositor EDS
% Data completeness (in resolution range)	99.5 (34.07-1.57) 99.6 (34.07-1.57)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.45 (at 1.58Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8.2_1309)	Depositor
R, R_{free}	0.195 , 0.234 0.203 , 0.243	Depositor DCC
R_{free} test set	1992 reflections (3.65%)	wwPDB-VP
Wilson B-factor (Å ²)	16.9	Xtriage
Anisotropy	0.455	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 43.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.044 for h,-k,-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	3704	wwPDB-VP
Average B, all atoms (Å ²)	25.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 89.33 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 3.5452e-08. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.36	0/1793	0.66	1/2448 (0.0%)
1	B	0.36	0/1749	0.64	0/2388
All	All	0.36	0/3542	0.65	1/4836 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	41	PRO	N-CA-CB	6.62	110.29	103.00

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	66	ASP	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1732	0	1644	8	0
1	B	1699	0	1623	6	0
2	A	2	0	0	0	0
2	B	2	0	0	0	0
3	A	131	0	0	2	0
3	B	138	0	0	2	0
All	All	3704	0	3267	14	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (14) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:41:PRO:N	3:A:404:HOH:O	2.32	0.62
1:A:211:LYS:HA	1:A:248:MET:HE3	1.87	0.57
1:A:164:ALA:O	3:A:401:HOH:O	2.17	0.56
1:B:176:ASN:ND2	3:B:401:HOH:O	2.32	0.56
1:A:129[A]:MET:HG3	1:A:133:HIS:CE1	2.46	0.51
1:A:217:SER:HA	1:A:265:MET:HE1	1.95	0.48
1:B:129[A]:MET:HG3	1:B:133:HIS:CE1	2.50	0.46
1:B:217:SER:HA	1:B:265:MET:HE1	1.99	0.44
1:B:264:ARG:NH1	3:B:409:HOH:O	2.51	0.43
1:B:78:LEU:HD13	1:B:198:ILE:HD11	2.01	0.42
1:A:149:ALA:HB3	1:A:150:PRO:HD3	2.02	0.41
1:A:248:MET:HE2	1:A:248:MET:HB2	1.83	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	237/231 (103%)	235 (99%)	2 (1%)	0	100	100
1	B	230/231 (100%)	227 (99%)	3 (1%)	0	100	100
All	All	467/462 (101%)	462 (99%)	5 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	172/172 (100%)	168 (98%)	4 (2%)	44	14
1	B	168/172 (98%)	167 (99%)	1 (1%)	78	65
All	All	340/344 (99%)	335 (98%)	5 (2%)	63	31

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	70[A]	PHE
1	A	70[B]	PHE
1	A	193	ASN
1	A	251	SER
1	B	251	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	97	GLN
1	B	97	GLN
1	B	193	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 4 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	230/231 (99%)	0.87	35 (15%) 5 9	8, 22, 39, 66	9 (3%)
1	B	229/231 (99%)	0.90	42 (18%) 3 6	7, 23, 44, 62	3 (1%)
All	All	459/462 (99%)	0.88	77 (16%) 4 7	7, 22, 43, 66	12 (2%)

All (77) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	68[A]	PRO	6.9
1	B	269	LEU	5.6
1	B	69	GLY	5.1
1	A	70[A]	PHE	4.8
1	B	70	PHE	4.8
1	A	270	ARG	4.7
1	A	69[A]	GLY	4.6
1	A	41	PRO	4.6
1	B	42	GLY	4.4
1	A	263	ALA	4.0
1	B	259	ILE	3.7
1	B	266	ALA	3.6
1	A	230	ALA	3.5
1	A	221	LEU	3.4
1	B	68	PRO	3.4
1	B	265	MET	3.4
1	A	168	TRP	3.3
1	B	247	VAL	3.2
1	A	229	TYR	3.2
1	B	241	PRO	3.1
1	B	270	ARG	3.1
1	A	217	SER	3.0
1	A	233	ALA	2.9
1	B	198	ILE	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	213	SER	2.9
1	B	264	ARG	2.9
1	A	218	LEU	2.9
1	B	258	ALA	2.8
1	B	199	ASP	2.8
1	B	262	THR	2.7
1	B	261	HIS	2.7
1	B	263	ALA	2.7
1	B	260	THR	2.6
1	A	165	ALA	2.6
1	B	243	ALA	2.6
1	B	257	ALA	2.6
1	A	67[A]	MET	2.6
1	A	269	LEU	2.6
1	B	212	ASP	2.6
1	B	231	ALA	2.5
1	B	221	LEU	2.5
1	A	225	ASP	2.5
1	A	228	HIS	2.5
1	A	187	PRO	2.5
1	B	238	ALA	2.5
1	A	240	PHE	2.5
1	B	210	ILE	2.4
1	B	230	ALA	2.4
1	B	236	PHE	2.4
1	A	259	ILE	2.4
1	B	165	ALA	2.3
1	B	228	HIS	2.3
1	A	226	THR	2.3
1	A	262	THR	2.3
1	B	219	GLY	2.3
1	B	253	PRO	2.3
1	B	229	TYR	2.3
1	A	231	ALA	2.3
1	B	252	ALA	2.3
1	A	186	GLY	2.3
1	A	236	PHE	2.2
1	A	200	GLY	2.2
1	A	222	GLY	2.2
1	B	226	THR	2.2
1	B	242	LYS	2.2
1	A	268	LYS	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	260	THR	2.2
1	A	251	SER	2.1
1	A	55	ALA	2.1
1	B	240	PHE	2.1
1	B	220	ASN	2.1
1	B	227	GLU	2.1
1	B	168	TRP	2.1
1	A	212	ASP	2.0
1	B	201	THR	2.0
1	A	183	PHE	2.0
1	B	218	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	ZN	A	302	1/1	0.97	0.11	22,22,22,22	1
2	ZN	B	302	1/1	0.98	0.10	24,24,24,24	1
2	ZN	B	301	1/1	0.99	0.02	17,17,17,17	0
2	ZN	A	301	1/1	0.99	0.03	18,18,18,18	0

6.5 Other polymers [i](#)

There are no such residues in this entry.