



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 05:48 PM UTC

PDB ID : 6TZH / pdb_00006tzh
Title : ADC-7 in complex with boronic acid transition state inhibitor S06015
Authors : Fish, E.R.; Powers, R.A.; Wallar, B.J.
Deposited on : 2019-08-12
Resolution : 2.04 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.04 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2260 (2.04-2.04)
Clashscore	190562	2333 (2.04-2.04)
Ramachandran outliers	187476	2318 (2.04-2.04)
Sidechain outliers	187428	2318 (2.04-2.04)
RSRZ outliers	180081	2260 (2.04-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	361	7% 93% 5% ..
1	B	361	93% 5% ..
1	C	361	11% 92% 6% ..
1	D	361	14% 94% ..

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	GLY	A	402	-	X	-	-

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 11315 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Beta-lactamase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	354	Total	C	N	O	S	0	0	0
			2744	1762	455	518	9			
1	B	358	Total	C	N	O	S	0	0	0
			2811	1806	466	530	9			
1	C	354	Total	C	N	O	S	0	0	0
			2706	1739	451	507	9			
1	D	356	Total	C	N	O	S	0	0	0
			2714	1742	447	516	9			

There are 4 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	MET	-	expression tag	UNP Q6DRA1
B	-1	MET	-	expression tag	UNP Q6DRA1
C	-1	MET	-	expression tag	UNP Q6DRA1
D	-1	MET	-	expression tag	UNP Q6DRA1

- Molecule 2 is phosphonoxy-[(4-thiophen-3-yl-1,2,3-triazol-1-yl)methyl]borinic acid (CCD ID: ERF) (formula: C₇H₉BN₃O₅PS) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms							ZeroOcc	AltConf
2	A	1	Total	B	C	N	O	P	S	0	0
			18	1	7	3	5	1	1		
2	B	1	Total	B	C	N	O	P	S	0	0
			18	1	7	3	5	1	1		
2	C	1	Total	B	C	N	O	P	S	0	0
			18	1	7	3	5	1	1		
2	D	1	Total	B	C	N	O	P	S	0	0
			18	1	7	3	5	1	1		

- Molecule 3 is GLYCINE (CCD ID: GLY) (formula: $C_2H_5NO_2$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			5	2	1	2		
3	B	1	Total	C	N	O	0	0
			5	2	1	2		
3	B	1	Total	C	N	O	0	0
			5	2	1	2		
3	C	1	Total	C	N	O	0	0
			5	2	1	2		
3	D	1	Total	C	N	O	0	0
			5	2	1	2		

- Molecule 4 is PHOSPHATE ION (CCD ID: PO4) (formula: O₄P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	O	P	0	0
			5	4	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	58	Total	O	0	0
			58	58		
5	B	112	Total	O	0	1
			113	113		
5	C	41	Total	O	0	1
			42	42		
5	D	25	Total	O	0	0
			25	25		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Beta-lactamase




• Molecule 1: Beta-lactamase



• Molecule 1: Beta-lactamase



• Molecule 1: Beta-lactamase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	88.55Å 81.46Å 105.67Å 90.00° 113.10° 90.00°	Depositor
Resolution (Å)	97.19 – 2.04 97.19 – 2.04	Depositor EDS
% Data completeness (in resolution range)	98.0 (97.19-2.04) 98.0 (97.19-2.04)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.08 (at 2.03Å)	Xtriage
Refinement program	REFMAC 5.8.0253	Depositor
R, R_{free}	0.221 , 0.254 (Not available) , 0.262	Depositor DCC
R_{free} test set	4321 reflections (4.87%)	wwPDB-VP
Wilson B-factor (Å ²)	38.6	Xtriage
Anisotropy	0.116	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 32.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.016 for h,-k,-h-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	11315	wwPDB-VP
Average B, all atoms (Å ²)	46.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 41.10 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 2.4809e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PO4, ERF

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.03	0/2811	1.34	0/3831
1	B	1.03	0/2879	1.34	1/3915 (0.0%)
1	C	1.03	0/2773	1.33	0/3784
1	D	1.03	0/2781	1.35	2/3800 (0.1%)
All	All	1.03	0/11244	1.34	3/15330 (0.0%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	193	LYS	CA-C-N	5.89	128.98	120.38
1	D	193	LYS	C-N-CA	5.89	128.98	120.38
1	B	5	ASP	CA-CB-CG	5.71	118.31	112.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2744	0	2664	8	0
1	B	2811	0	2778	8	0
1	C	2706	0	2592	10	0
1	D	2714	0	2575	13	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	A	18	0	0	0	0
2	B	18	0	0	0	0
2	C	18	0	0	0	0
2	D	18	0	0	0	0
3	A	5	0	2	0	0
3	B	10	0	4	0	0
3	C	5	0	2	0	0
3	D	5	0	2	0	0
4	B	5	0	0	0	0
5	A	58	0	0	0	0
5	B	113	0	0	0	0
5	C	42	0	0	0	0
5	D	25	0	0	0	0
All	All	11315	0	10619	38	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (38) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:248:ILE:O	1:D:252:ILE:HG23	1.80	0.82
1:D:238:ASN:HA	1:D:252:ILE:HD12	1.64	0.79
1:C:99:THR:HG22	1:C:100:PRO:HD2	1.65	0.78
1:D:238:ASN:HA	1:D:252:ILE:CD1	2.23	0.68
1:B:4:LYS:HE3	1:B:34:GLN:OE1	1.96	0.66
1:D:241:PRO:HB3	1:D:252:ILE:HG12	1.78	0.64
1:B:208:GLN:OE1	1:C:51:LYS:HA	2.04	0.57
1:A:46:GLN:HG3	1:A:200:ALA:HA	1.91	0.51
1:C:99:THR:HG21	1:C:136:LYS:HA	1.91	0.51
1:D:241:PRO:HB3	1:D:252:ILE:CG1	2.40	0.51
1:A:87:ASP:OD2	1:A:92:TYR:OH	2.26	0.51
1:D:241:PRO:HB3	1:D:252:ILE:CD1	2.41	0.51
1:B:87:ASP:OD2	1:B:92:TYR:OH	2.26	0.51
1:B:46:GLN:HG3	1:B:200:ALA:HA	1.94	0.48
1:C:46:GLN:HG3	1:C:200:ALA:HA	1.95	0.48
1:A:172:GLN:OE1	1:A:176:LYS:HE3	2.13	0.48
1:D:276:TYR:CD1	1:D:277:PRO:HA	2.50	0.47
1:D:46:GLN:HG3	1:D:200:ALA:HA	1.97	0.47
1:C:264:THR:OG1	1:C:282:THR:HG23	2.15	0.46
1:C:276:TYR:CD1	1:C:277:PRO:HA	2.51	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:95:GLU:HG3	1:B:132:LEU:HD21	1.97	0.45
1:A:29:ALA:HB1	1:A:228:LEU:HG	1.98	0.45
1:D:248:ILE:O	1:D:252:ILE:CG2	2.58	0.45
1:A:28:MET:HE2	1:A:337:THR:CG2	2.47	0.44
1:A:276:TYR:CD1	1:A:277:PRO:HA	2.53	0.44
1:B:276:TYR:CD1	1:B:277:PRO:HA	2.52	0.43
1:D:197:GLN:HE21	1:D:197:GLN:HB3	1.62	0.43
1:C:87:ASP:OD2	1:C:92:TYR:OH	2.26	0.43
1:A:290:GLN:OE1	1:A:294:LYS:HD2	2.18	0.43
1:C:99:THR:CG2	1:C:100:PRO:HD2	2.40	0.42
1:A:241:PRO:HB2	1:A:249:GLN:HG3	2.02	0.42
1:D:237:ALA:O	1:D:252:ILE:HD13	2.20	0.42
1:B:35:ASN:HA	1:B:330:ASN:ND2	2.35	0.41
1:D:241:PRO:HB3	1:D:252:ILE:HD11	2.03	0.41
1:D:354:LEU:HD12	1:D:354:LEU:HA	1.87	0.41
1:B:179:PHE:HB2	1:B:180:PRO:HD3	2.02	0.40
1:C:152:ASN:N	1:C:153:PRO:HD2	2.37	0.40
1:C:219:ALA:HB3	1:C:220:PRO:HD3	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	352/361 (98%)	343 (97%)	9 (3%)	0	100	100
1	B	356/361 (99%)	348 (98%)	8 (2%)	0	100	100
1	C	352/361 (98%)	345 (98%)	7 (2%)	0	100	100
1	D	354/361 (98%)	346 (98%)	8 (2%)	0	100	100
All	All	1414/1444 (98%)	1382 (98%)	32 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	293/319 (92%)	288 (98%)	5 (2%)	53	53
1	B	306/319 (96%)	298 (97%)	8 (3%)	40	37
1	C	281/319 (88%)	272 (97%)	9 (3%)	34	29
1	D	282/319 (88%)	275 (98%)	7 (2%)	42	39
All	All	1162/1276 (91%)	1133 (98%)	29 (2%)	42	39

All (29) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	46	GLN
1	A	93	TRP
1	A	228	LEU
1	A	269	LEU
1	A	311	HIS
1	B	4	LYS
1	B	41	MET
1	B	46	GLN
1	B	95	GLU
1	B	97	LYS
1	B	206	GLU
1	B	269	LEU
1	B	311	HIS
1	C	46	GLN
1	C	80	LYS
1	C	99	THR
1	C	197	GLN
1	C	206	GLU
1	C	269	LEU
1	C	302	SER
1	C	308	LYS
1	C	311	HIS
1	D	41	MET
1	D	197	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	D	252	ILE
1	D	269	LEU
1	D	301	ILE
1	D	311	HIS
1	D	354	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (20) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	14	GLN
1	A	195	GLN
1	A	198	ASN
1	A	205	GLN
1	A	330	ASN
1	B	126	GLN
1	C	34	GLN
1	C	79	ASN
1	C	120	GLN
1	C	152	ASN
1	C	290	GLN
1	C	330	ASN
1	D	34	GLN
1	D	120	GLN
1	D	142	ASN
1	D	152	ASN
1	D	197	GLN
1	D	281	GLN
1	D	290	GLN
1	D	355	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

10 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	ERF	A	401	1	17,19,19	2.14	5 (29%)	18,27,27	2.82	7 (38%)
3	GLY	C	402	-	4,4,4	1.03	0	3,4,4	1.48	0
3	GLY	B	404	-	4,4,4	1.26	1 (25%)	3,4,4	1.40	0
4	PO4	B	402	-	4,4,4	0.62	0	6,6,6	0.49	0
3	GLY	A	402	-	4,4,4	1.19	1 (25%)	3,4,4	1.66	1 (33%)
3	GLY	D	402	-	4,4,4	1.08	1 (25%)	3,4,4	1.26	0
2	ERF	D	401	1	17,19,19	2.73	4 (23%)	18,27,27	3.01	6 (33%)
2	ERF	C	401	1	17,19,19	2.66	4 (23%)	18,27,27	3.05	6 (33%)
3	GLY	B	403	-	4,4,4	1.14	1 (25%)	3,4,4	1.30	0
2	ERF	B	401	1	17,19,19	2.24	4 (23%)	18,27,27	2.96	8 (44%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ERF	A	401	1	-	1/1/13/13	0/2/2/2
3	GLY	C	402	-	-	2/2/2/2	-
3	GLY	B	404	-	-	2/2/2/2	-
3	GLY	A	402	-	-	2/2/2/2	-
3	GLY	D	402	-	-	2/2/2/2	-
2	ERF	D	401	1	-	0/1/13/13	0/2/2/2
2	ERF	C	401	1	-	1/1/13/13	0/2/2/2

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GLY	B	403	-	-	2/2/2/2	-
2	ERF	B	401	1	-	0/1/13/13	0/2/2/2

All (21) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	401	ERF	P1-O10	7.69	1.63	1.56
2	D	401	ERF	P1-O10	7.36	1.63	1.56
2	D	401	ERF	N3-N2	6.18	1.42	1.32
2	B	401	ERF	N3-N2	5.90	1.41	1.32
2	C	401	ERF	N3-N2	5.62	1.41	1.32
2	A	401	ERF	N3-N2	5.17	1.40	1.32
2	B	401	ERF	N1-N2	4.92	1.42	1.34
2	D	401	ERF	N1-N2	4.34	1.41	1.34
2	A	401	ERF	P1-O10	4.29	1.60	1.56
2	A	401	ERF	N1-N2	4.00	1.41	1.34
2	C	401	ERF	N1-N2	3.76	1.40	1.34
2	B	401	ERF	P1-O10	3.14	1.59	1.56
2	D	401	ERF	B7-C6	2.42	1.61	1.57
3	B	404	GLY	OXT-C	-2.41	1.22	1.30
2	A	401	ERF	C2-C3	2.34	1.40	1.37
3	A	402	GLY	OXT-C	-2.32	1.23	1.30
2	B	401	ERF	C2-C3	2.29	1.40	1.37
3	B	403	GLY	OXT-C	-2.16	1.23	1.30
2	C	401	ERF	B7-C6	2.07	1.60	1.57
2	A	401	ERF	B7-C6	2.04	1.60	1.57
3	D	402	GLY	OXT-C	-2.03	1.24	1.30

All (28) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	401	ERF	C2-C3-N3	6.73	115.06	108.33
2	C	401	ERF	C3-N3-N2	-6.71	103.12	108.84
2	D	401	ERF	C3-N3-N2	-6.42	103.37	108.84
2	B	401	ERF	C2-C3-N3	6.21	114.54	108.33
2	A	401	ERF	C2-C3-N3	5.97	114.31	108.33
2	D	401	ERF	C2-C3-N3	5.96	114.29	108.33
2	D	401	ERF	C2-C3-C7	-5.92	119.96	130.55
2	B	401	ERF	C3-N3-N2	-5.41	104.23	108.84
2	A	401	ERF	C6-N1-N2	-5.26	113.28	119.86
2	A	401	ERF	C3-N3-N2	-5.19	104.41	108.84
2	C	401	ERF	C11-C7-C8	5.13	118.48	111.24

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	401	ERF	C2-C3-C7	-4.58	122.36	130.55
2	B	401	ERF	C2-C3-C7	-4.46	122.56	130.55
2	D	401	ERF	C11-C7-C8	4.44	117.51	111.24
2	B	401	ERF	C6-N1-N2	-4.33	114.44	119.86
2	B	401	ERF	C11-C7-C8	4.30	117.32	111.24
2	B	401	ERF	C7-C8-S1	-4.30	105.06	112.23
2	A	401	ERF	C11-C7-C8	4.28	117.28	111.24
2	D	401	ERF	C7-C8-S1	-4.20	105.22	112.23
2	C	401	ERF	C7-C8-S1	-3.45	106.48	112.23
2	A	401	ERF	C2-C3-C7	-3.44	124.39	130.55
2	C	401	ERF	C6-N1-N2	-3.15	115.92	119.86
2	A	401	ERF	C7-C8-S1	-3.12	107.03	112.23
2	D	401	ERF	C7-C3-N3	2.49	125.52	121.11
3	A	402	GLY	OXT-C-O	-2.35	117.29	123.33
2	A	401	ERF	C3-C2-N1	-2.27	102.69	104.95
2	B	401	ERF	C3-C2-N1	-2.07	102.89	104.95
2	B	401	ERF	C2-N1-N2	2.04	112.75	110.75

There are no chirality outliers.

All (12) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	C	401	ERF	C2-C3-C7-C8
3	A	402	GLY	O-C-CA-N
3	A	402	GLY	OXT-C-CA-N
3	B	403	GLY	O-C-CA-N
3	B	403	GLY	OXT-C-CA-N
3	B	404	GLY	OXT-C-CA-N
3	C	402	GLY	O-C-CA-N
3	C	402	GLY	OXT-C-CA-N
3	D	402	GLY	O-C-CA-N
3	B	404	GLY	O-C-CA-N
3	D	402	GLY	OXT-C-CA-N
2	A	401	ERF	C2-C3-C7-C8


There are no ring outliers.


No monomer is involved in short contacts.


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is

within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	354/361 (98%)	0.57	25 (7%) 22 22	27, 41, 62, 84	0
1	B	358/361 (99%)	-0.02	1 (0%) 90 92	24, 34, 49, 63	0
1	C	354/361 (98%)	0.89	39 (11%) 10 10	28, 51, 75, 88	0
1	D	356/361 (98%)	1.14	49 (13%) 6 6	36, 56, 77, 104	0
All	All	1422/1444 (98%)	0.64	114 (8%) 18 18	24, 45, 72, 104	0

All (114) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	144	ILE	4.1
1	D	331	ILE	3.9
1	A	358	LYS	3.7
1	A	133	THR	3.6
1	D	241	PRO	3.6
1	A	101	ILE	3.6
1	D	11	LEU	3.5
1	A	132	LEU	3.5
1	A	5	ASP	3.5
1	D	252	ILE	3.4
1	A	93	TRP	3.4
1	C	147	TYR	3.3
1	D	3	PRO	3.3
1	C	140	PRO	3.2
1	C	5	ASP	3.2
1	D	325	PHE	3.2
1	C	98	ASN	3.2
1	C	96	LEU	3.1
1	C	145	GLY	3.1
1	D	269	LEU	3.1
1	C	106	LEU	3.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	301	ILE	3.1
1	C	107	LEU	3.0
1	C	298	VAL	3.0
1	A	125	VAL	3.0
1	C	280	LEU	2.9
1	D	357	ILE	2.9
1	D	85	PHE	2.9
1	A	131	VAL	2.9
1	C	110	ALA	2.9
1	C	138	TRP	2.9
1	C	132	LEU	2.8
1	C	264	THR	2.8
1	C	330	ASN	2.8
1	C	115	GLY	2.8
1	A	138	TRP	2.8
1	D	248	ILE	2.8
1	D	326	ILE	2.8
1	C	136	LYS	2.8
1	D	36	ASN	2.7
1	C	161	VAL	2.7
1	C	167	ASN	2.6
1	D	277	PRO	2.6
1	C	164	LEU	2.6
1	D	310	TYR	2.6
1	D	327	PRO	2.6
1	B	1	ASN	2.6
1	C	104	VAL	2.5
1	C	301	ILE	2.5
1	D	274	PHE	2.5
1	C	248	ILE	2.5
1	D	59	ILE	2.5
1	D	307	VAL	2.5
1	C	357	ILE	2.5
1	D	32	VAL	2.5
1	A	357	ILE	2.5
1	D	83	ILE	2.5
1	D	300	ALA	2.4
1	A	263	ASN	2.4
1	D	35	ASN	2.4
1	D	76	TYR	2.4
1	D	328	LYS	2.4
1	D	247	ASP	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	248	ILE	2.4
1	D	298	VAL	2.4
1	D	263	ASN	2.4
1	C	131	VAL	2.4
1	C	85	PHE	2.3
1	C	93	TRP	2.3
1	A	116	ASN	2.3
1	C	81	GLY	2.3
1	C	139	LYS	2.3
1	D	303	LYS	2.3
1	D	178	ILE	2.3
1	C	356	ALA	2.3
1	D	8	ILE	2.3
1	A	129	GLN	2.2
1	A	96	LEU	2.2
1	C	135	PHE	2.2
1	A	144	ILE	2.2
1	D	352	VAL	2.2
1	D	5	ASP	2.2
1	D	239	LEU	2.2
1	D	186	HIS	2.2
1	C	142	ASN	2.2
1	A	293	MET	2.2
1	A	117	LEU	2.2
1	D	93	TRP	2.2
1	A	99	THR	2.2
1	C	279	THR	2.2
1	D	194	THR	2.2
1	A	134	PHE	2.1
1	C	257	GLN	2.1
1	D	278	ALA	2.1
1	A	306	SER	2.1
1	C	39	TYR	2.1
1	C	92	TYR	2.1
1	D	246	THR	2.1
1	D	305	PRO	2.1
1	A	135	PHE	2.1
1	A	140	PRO	2.1
1	A	121	PHE	2.1
1	D	234	PHE	2.1
1	C	262	VAL	2.1
1	D	75	GLY	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	159	GLY	2.1
1	D	356	ALA	2.1
1	D	20	LEU	2.0
1	D	249	GLN	2.0
1	A	7	GLU	2.0
1	D	304	GLU	2.0
1	D	330	ASN	2.0
1	C	129	GLN	2.0
1	D	164	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	PO4	B	402	5/5	0.77	0.14	65,67,69,73	5
3	GLY	B	404	5/5	0.83	0.18	69,69,73,75	0
3	GLY	B	403	5/5	0.84	0.17	47,48,70,71	0
3	GLY	A	402	5/5	0.85	0.13	53,55,65,65	0
3	GLY	C	402	5/5	0.86	0.14	55,55,60,64	0
3	GLY	D	402	5/5	0.87	0.13	53,53,59,62	0
2	ERF	A	401	18/18	0.94	0.09	34,53,62,62	0
2	ERF	C	401	18/18	0.94	0.08	37,50,56,59	0
2	ERF	D	401	18/18	0.94	0.08	39,49,59,64	0
2	ERF	B	401	18/18	0.95	0.07	31,42,57,67	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around ERF A 401:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around ERF C 401:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around ERF D 401:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around ERF B 401:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.