



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 01:52 PM UTC

PDB ID : 3U09 / pdb_00003u09
Title : Crystal structure of 3-ketoacyl-(acyl-carrier-protein) reductase (FabG)(G92D) from *Vibrio cholerae*
Authors : Hou, J.; Chruszcz, M.; Zheng, H.; Grabowski, M.; Fratzczak, Z.; Anderson, W.F.; Minor, W.; Center for Structural Genomics of Infectious Diseases (CS-GID)
Deposited on : 2011-09-28
Resolution : 1.75 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3183 (1.76-1.76)
Clashscore	190562	3299 (1.76-1.76)
Ramachandran outliers	187476	3274 (1.76-1.76)
Sidechain outliers	187428	3274 (1.76-1.76)
RSRZ outliers	180081	3183 (1.76-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	251	 90% 9% .
1	B	251	 88% 10% .

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 4039 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 3-oxoacyl-[acyl-carrier-protein] reductase FabG.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	248	Total	C	N	O	S	0	2	0
			1825	1134	322	357	12			
1	B	248	Total	C	N	O	S	0	2	0
			1814	1131	316	355	12			

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	SER	-	expression tag	UNP Q9KQH7
A	-1	ASN	-	expression tag	UNP Q9KQH7
A	0	ALA	-	expression tag	UNP Q9KQH7
A	1	MET	-	expression tag	UNP Q9KQH7
A	2	SER	-	expression tag	UNP Q9KQH7
A	3	GLN	-	expression tag	UNP Q9KQH7
A	4	PHE	-	expression tag	UNP Q9KQH7
A	92	ASP	GLY	engineered mutation	UNP Q9KQH7
B	-2	SER	-	expression tag	UNP Q9KQH7
B	-1	ASN	-	expression tag	UNP Q9KQH7
B	0	ALA	-	expression tag	UNP Q9KQH7
B	1	MET	-	expression tag	UNP Q9KQH7
B	2	SER	-	expression tag	UNP Q9KQH7
B	3	GLN	-	expression tag	UNP Q9KQH7
B	4	PHE	-	expression tag	UNP Q9KQH7
B	92	ASP	GLY	engineered mutation	UNP Q9KQH7

- Molecule 2 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 3 is UNKNOWN ATOM OR ION (CCD ID: UNX) (formula: X).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	5	Total	X	0	0
			5	5		
3	B	5	Total	X	0	0
			5	5		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	171	Total 171	O 171	0	0
4	B	179	Total 179	O 179	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: 3-oxoacyl-[acyl-carrier-protein] reductase FabG

Chain A:  90% 9% .



- Molecule 1: 3-oxoacyl-[acyl-carrier-protein] reductase FabG

Chain B:  88% 10% .



4 Data and refinement statistics

Property	Value	Source
Space group	P 62	Depositor
Cell constants a, b, c, α , β , γ	63.57Å 63.57Å 191.06Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	27.00 – 1.75 27.00 – 1.75	Depositor EDS
% Data completeness (in resolution range)	97.1 (27.00-1.75) 97.2 (27.00-1.75)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.20 (at 1.75Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.160 , 0.187 0.163 , 0.188	Depositor DCC
R_{free} test set	2131 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	21.6	Xtriage
Anisotropy	0.131	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 28.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	0.225 for h,-h-k,-l	Xtriage
Reported twinning fraction	0.833 for H, K, L 0.167 for K, H, -L	Depositor
Outliers	0 of 42620 reflections	Xtriage
F_o, F_c correlation	0.98	EDS
Total number of atoms	4039	wwPDB-VP
Average B, all atoms (Å ²)	27.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.71% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: UNX, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.69	0/1849	0.82	0/2492
1	B	0.69	0/1838	0.83	0/2479
All	All	0.69	0/3687	0.83	0/4971

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1825	0	1851	19	0
1	B	1814	0	1835	28	0
2	A	20	0	0	0	0
2	B	20	0	0	1	0
3	A	5	0	0	0	0
3	B	5	0	0	0	0
4	A	171	0	0	0	0
4	B	179	0	0	1	0
All	All	4039	0	3686	45	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (45) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:187[B]:PHE:CD2	1:A:245:MET:HE1	1.83	1.14
1:A:5:MET:HE2	1:B:5:MET:HE2	1.29	1.11
1:B:187[B]:PHE:CD2	1:B:245:MET:HE1	1.92	1.03
1:A:187[B]:PHE:HD2	1:A:245:MET:HE1	1.46	0.79
1:B:126:LEU:HG	1:B:130:MET:HE2	1.68	0.75
1:A:187[B]:PHE:CE2	1:A:245:MET:HE1	2.25	0.70
1:A:126:LEU:HG	1:A:130:MET:HE2	1.77	0.67
1:B:187[B]:PHE:CD2	1:B:245:MET:CE	2.78	0.59
1:A:126:LEU:HD22	1:A:169[A]:MET:HE3	1.84	0.59
1:B:118:ILE:HD12	1:B:162:VAL:HG21	1.85	0.59
1:A:5:MET:CE	1:B:5:MET:HE2	2.20	0.58
1:A:208:VAL:HG22	1:A:245:MET:HE3	1.85	0.58
1:B:64:VAL:HG21	4:B:363:HOH:O	2.03	0.58
1:A:166:THR:HG23	1:A:180:VAL:HG12	1.88	0.56
1:B:169[B]:MET:HE3	1:B:169[B]:MET:HA	1.88	0.54
1:B:187[B]:PHE:HD2	1:B:245:MET:HE1	1.62	0.53
1:B:205:LEU:HD21	1:B:213:LEU:HD12	1.91	0.52
1:A:35:LYS:HG2	1:A:81:PHE:CD1	2.46	0.51
1:B:166:THR:HG23	1:B:180:VAL:HG12	1.93	0.50
1:B:2:SER:OG	2:B:249:SO4:O1	2.28	0.45
1:B:4:PHE:CE2	1:B:5:MET:HG3	2.51	0.45
1:B:187[B]:PHE:CE2	1:B:204:THR:HB	2.51	0.45
1:A:118:ILE:HD12	1:A:162:VAL:HG21	1.99	0.44
1:B:35:LYS:NZ	1:B:55:ASP:O	2.50	0.44
1:B:126:LEU:HD21	1:B:173:VAL:HG11	1.99	0.44
1:A:36:VAL:O	1:A:57:GLY:HA3	2.18	0.43
1:B:169[B]:MET:HE3	1:B:169[B]:MET:CA	2.49	0.43
1:B:37:ILE:HD12	1:B:37:ILE:N	2.34	0.43
1:B:126:LEU:HD22	1:B:169[A]:MET:HE3	2.00	0.43
1:B:118:ILE:HD12	1:B:162:VAL:CG2	2.48	0.43
1:B:111:MET:HA	1:B:111:MET:HE2	1.99	0.43
1:B:187[B]:PHE:HD2	1:B:245:MET:CE	2.27	0.43
1:A:133:ARG:HA	1:A:176:ARG:O	2.19	0.42
1:A:4:PHE:CE2	1:A:5:MET:HG3	2.54	0.42
1:B:133:ARG:HA	1:B:176:ARG:O	2.20	0.42
1:B:36:VAL:O	1:B:57:GLY:HA3	2.20	0.41
1:B:187[B]:PHE:CE2	1:B:245:MET:HE1	2.50	0.41
1:B:208:VAL:HG22	1:B:245:MET:HE3	2.01	0.41
1:A:137:ILE:CD1	1:A:169[A]:MET:CE	2.99	0.41
1:A:187[B]:PHE:HD2	1:A:245:MET:CE	2.23	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:169[B]:MET:HA	1:B:169[B]:MET:CE	2.47	0.40
1:A:187[B]:PHE:CD2	1:A:245:MET:CE	2.76	0.40
1:A:111:MET:HA	1:A:111:MET:HE2	2.04	0.40
1:A:137:ILE:CD1	1:A:169[A]:MET:HE1	2.52	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	248/251 (99%)	243 (98%)	5 (2%)	0	100	100
1	B	248/251 (99%)	243 (98%)	5 (2%)	0	100	100
All	All	496/502 (99%)	486 (98%)	10 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	186/190 (98%)	184 (99%)	2 (1%)	65	52
1	B	183/190 (96%)	183 (100%)	0	100	100
All	All	369/380 (97%)	367 (100%)	2 (0%)	81	75

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	105	GLU
1	A	217	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	134	GLN
1	B	149	ASN
1	B	152	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 18 ligands modelled in this entry, 10 are unknown - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	SO4	A	249	-	4,4,4	0.47	0	6,6,6	0.18	0
2	SO4	A	251	-	4,4,4	0.42	0	6,6,6	0.19	0
2	SO4	B	251	-	4,4,4	0.36	0	6,6,6	0.30	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	SO4	A	252	-	4,4,4	0.43	0	6,6,6	0.07	0
2	SO4	B	250	-	4,4,4	0.47	0	6,6,6	0.19	0
2	SO4	B	252	-	4,4,4	0.49	0	6,6,6	0.36	0
2	SO4	B	249	-	4,4,4	0.35	0	6,6,6	0.43	0
2	SO4	A	250	-	4,4,4	0.42	0	6,6,6	0.24	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	249	SO4	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	248/251 (98%)	-1.33	0 100 100	10, 22, 42, 57	3 (1%)
1	B	248/251 (98%)	-1.29	0 100 100	13, 24, 45, 63	4 (1%)
All	All	496/502 (98%)	-1.31	0 100 100	10, 23, 44, 63	7 (1%)

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	SO4	A	250	5/5	0.97	0.04	54,57,60,61	0
2	SO4	A	249	5/5	0.98	0.04	55,59,64,64	0
2	SO4	A	251	5/5	0.98	0.06	33,36,37,39	5
2	SO4	B	252	5/5	0.98	0.04	49,53,57,57	0
3	UNX	A	337	1/1	0.98	0.06	39,39,39,39	0
3	UNX	B	327	1/1	0.98	0.08	44,44,44,44	0
3	UNX	B	328	1/1	0.98	0.08	40,40,40,40	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	UNX	B	338	1/1	0.98	0.07	43,43,43,43	0
3	UNX	A	323	1/1	0.99	0.10	41,41,41,41	0
3	UNX	A	324	1/1	0.99	0.13	47,47,47,47	0
3	UNX	A	325	1/1	0.99	0.09	40,40,40,40	0
2	SO4	B	250	5/5	0.99	0.03	51,55,59,61	0
3	UNX	B	326	1/1	0.99	0.07	38,38,38,38	0
2	SO4	B	251	5/5	0.99	0.03	50,51,52,53	0
2	SO4	A	252	5/5	0.99	0.02	53,54,55,55	5
3	UNX	A	300	1/1	0.99	0.06	36,36,36,36	0
3	UNX	B	339	1/1	0.99	0.05	38,38,38,38	0
2	SO4	B	249	5/5	1.00	0.04	35,38,40,41	0

6.5 Other polymers [i](#)

There are no such residues in this entry.