



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 12, 2026 – 06:00 PM UTC

PDB ID : 7U1C / pdb_00007u1c
Title : Structure of EstG crystalized with SO4 and Tris
Authors : Gabelli, S.B.; Chen, Z.
Deposited on : 2022-02-20
Resolution : 2.09 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.09 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6658 (2.10-2.10)
Clashscore	190562	7164 (2.10-2.10)
Ramachandran outliers	187476	7099 (2.10-2.10)
Sidechain outliers	187428	7100 (2.10-2.10)
RSRZ outliers	180081	6662 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	469	

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 3317 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Beta-lactamase domain-containing protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	401	3017	1914	527	559	17	0	0	0

There are 7 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	463	LYS	-	expression tag	UNP Q9A800
A	464	LEU	-	expression tag	UNP Q9A800
A	465	ALA	-	expression tag	UNP Q9A800
A	466	ALA	-	expression tag	UNP Q9A800
A	467	ALA	-	expression tag	UNP Q9A800
A	468	LEU	-	expression tag	UNP Q9A800
A	469	GLU	-	expression tag	UNP Q9A800

- Molecule 2 is SULFATE ION (CCD ID: SO4) (formula: O₄S).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		

- Molecule 3 is SODIUM ION (CCD ID: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Na	0	0
			1	1		

- Molecule 4 is 2-AMINO-2-HYDROXYMETHYL-PROPANE-1,3-DIOL (CCD ID: TRS) (formula: C₄H₁₂NO₃).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			8	4	1	3		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	286	Total	O	0	0
			286	286		

- Molecule 1: Beta-lactamase domain-containing protein



4 Data and refinement statistics

Property	Value	Source
Space group	P 41	Depositor
Cell constants a, b, c, α , β , γ	111.25Å 111.25Å 56.87Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	27.83 – 2.09 27.83 – 2.09	Depositor EDS
% Data completeness (in resolution range)	99.2 (27.83-2.09) 99.6 (27.83-2.09)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.09 (at 2.08Å)	Xtriage
Refinement program	REFMAC 5.8.0267	Depositor
R, R_{free}	0.172 , 0.210 (Not available) , 0.221	Depositor DCC
R_{free} test set	2092 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	37.6	Xtriage
Anisotropy	0.284	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 31.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.025 for h,-k,-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	3317	wwPDB-VP
Average B, all atoms (Å ²)	46.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.49% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: TRS, SO4, NA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.10	2/3088 (0.1%)	1.35	6/4196 (0.1%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	396	TRP	C-O	6.81	1.30	1.24
1	A	107	THR	C-O	5.62	1.30	1.24

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	175	ASN	CB-CA-C	-13.74	90.59	112.03
1	A	198	ASP	CA-CB-CG	5.93	118.53	112.60
1	A	151	THR	CB-CA-C	5.53	118.00	109.15
1	A	218	TYR	CB-CA-C	5.52	118.12	109.84
1	A	346	PHE	CA-C-O	-5.35	115.50	121.81
1	A	310	PHE	CA-CB-CG	-5.31	108.49	113.80

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3017	0	3000	70	0
2	A	5	0	0	0	0
3	A	1	0	0	0	0
4	A	8	0	12	2	0
5	A	286	0	0	35	0
All	All	3317	0	3012	70	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

All (70) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:278:GLN:NE2	1:A:278:GLN:HA	1.64	1.09
1:A:136:LEU:HB3	1:A:208:MET:HE2	1.36	1.07
1:A:140:LYS:HA	5:A:708:HOH:O	1.56	1.06
1:A:141:GLY:HA2	5:A:606:HOH:O	1.65	0.95
1:A:351:ASN:HB2	5:A:679:HOH:O	1.68	0.92
1:A:278:GLN:HA	1:A:278:GLN:HE21	1.34	0.92
1:A:77:GLN:HG3	5:A:775:HOH:O	1.74	0.86
1:A:136:LEU:HB3	1:A:208:MET:CE	2.06	0.85
1:A:222:ALA:HB1	5:A:782:HOH:O	1.77	0.84
1:A:153:VAL:HB	1:A:208:MET:HE1	1.58	0.84
1:A:141:GLY:CA	5:A:606:HOH:O	2.24	0.83
1:A:370:MET:HG3	5:A:844:HOH:O	1.79	0.81
1:A:31:LEU:HB2	5:A:819:HOH:O	1.82	0.79
1:A:370:MET:SD	5:A:844:HOH:O	2.42	0.77
1:A:140:LYS:HD2	5:A:708:HOH:O	1.88	0.73
1:A:30:ALA:CB	5:A:685:HOH:O	2.37	0.72
1:A:30:ALA:HB1	5:A:685:HOH:O	1.89	0.71
1:A:222:ALA:CB	5:A:782:HOH:O	2.33	0.71
1:A:403:TRP:CH2	5:A:609:HOH:O	2.43	0.70
1:A:49:LEU:HD12	5:A:741:HOH:O	1.92	0.69
1:A:370:MET:CG	5:A:844:HOH:O	2.35	0.69
1:A:104:LYS:HA	5:A:782:HOH:O	1.93	0.68
1:A:98:ARG:H	1:A:419:GLN:HE22	1.41	0.66
1:A:218:TYR:OH	4:A:503:TRS:H11	1.95	0.66
1:A:98:ARG:H	1:A:419:GLN:NE2	1.96	0.62
1:A:209:PHE:HD1	5:A:627:HOH:O	1.84	0.61
1:A:110:ALA:HA	1:A:113:MET:HE2	1.81	0.61
1:A:231:LYS:HE2	5:A:853:HOH:O	2.01	0.60
1:A:53:MET:HE2	5:A:741:HOH:O	2.01	0.60

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:31:LEU:HD22	5:A:819:HOH:O	2.02	0.59
1:A:292:LYS:HE2	5:A:848:HOH:O	2.02	0.59
1:A:347:VAL:HG12	1:A:349:THR:HG22	1.85	0.58
1:A:278:GLN:HE21	1:A:278:GLN:CA	2.07	0.57
1:A:189:GLU:HA	1:A:192:GLN:HE21	1.70	0.56
1:A:281:PRO:HD2	5:A:845:HOH:O	2.06	0.55
1:A:403:TRP:CZ3	5:A:609:HOH:O	2.60	0.54
1:A:53:MET:HE1	1:A:428:LEU:HD22	1.92	0.51
1:A:153:VAL:CB	1:A:208:MET:HE1	2.35	0.51
1:A:412:LEU:HD12	1:A:412:LEU:C	2.36	0.50
1:A:189:GLU:HB3	1:A:192:GLN:HG2	1.95	0.49
1:A:209:PHE:CD1	5:A:627:HOH:O	2.55	0.48
1:A:247:THR:HB	1:A:248:PRO:HD3	1.95	0.48
1:A:126:VAL:HB	5:A:868:HOH:O	2.14	0.48
1:A:347:VAL:HG12	1:A:349:THR:CG2	2.44	0.47
1:A:408:PRO:HD2	5:A:666:HOH:O	2.16	0.46
1:A:431:GLN:NE2	1:A:431:GLN:HA	2.30	0.45
1:A:416:GLY:CA	5:A:609:HOH:O	2.64	0.45
1:A:39:GLY:O	1:A:71:ARG:HD3	2.16	0.45
1:A:106:VAL:HG12	1:A:241:MET:HE2	1.99	0.45
1:A:140:LYS:CD	5:A:708:HOH:O	2.56	0.45
1:A:230:GLU:HA	1:A:235:GLN:O	2.16	0.45
1:A:199:LEU:C	1:A:199:LEU:HD23	2.42	0.45
1:A:347:VAL:O	1:A:349:THR:HG23	2.17	0.44
1:A:165:HIS:HD2	1:A:219:SER:OG	2.00	0.44
1:A:417:MET:N	5:A:609:HOH:O	2.51	0.44
1:A:101:SER:O	1:A:104:LYS:HB3	2.17	0.44
1:A:159:MET:O	1:A:163:MET:HG2	2.16	0.43
1:A:79:ARG:HG2	1:A:81:PHE:CZ	2.53	0.43
1:A:218:TYR:OH	4:A:503:TRS:C1	2.64	0.43
1:A:104:LYS:HE2	1:A:165:HIS:HE1	1.84	0.43
1:A:140:LYS:C	5:A:606:HOH:O	2.61	0.43
1:A:140:LYS:CE	5:A:708:HOH:O	2.66	0.43
1:A:444:PRO:HD2	5:A:603:HOH:O	2.18	0.43
1:A:416:GLY:C	5:A:609:HOH:O	2.62	0.42
1:A:100:ARG:O	1:A:301:GLY:HA2	2.19	0.42
1:A:237:LEU:N	1:A:238:PRO:CD	2.82	0.42
1:A:86:LEU:HD23	1:A:86:LEU:HA	1.92	0.42
1:A:431:GLN:HA	1:A:431:GLN:HE21	1.85	0.41
1:A:30:ALA:HB3	5:A:685:HOH:O	2.14	0.41
1:A:377:ALA:O	1:A:394:VAL:HA	2.21	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	397/469 (85%)	388 (98%)	8 (2%)	1 (0%)	36	36

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	301	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	310/358 (87%)	294 (95%)	16 (5%)	21	20

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	31	LEU
1	A	37	GLU
1	A	76	VAL
1	A	79	ARG
1	A	83	ARG
1	A	84	ARG
1	A	142	VAL

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	151	THR
1	A	155	ARG
1	A	178	SER
1	A	183	GLN
1	A	200	VAL
1	A	263	GLN
1	A	278	GLN
1	A	332	GLU
1	A	370	MET

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	129	HIS
1	A	165	HIS
1	A	192	GLN
1	A	278	GLN
1	A	411	ASN
1	A	419	GLN
1	A	431	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 1 is monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	TRS	A	503	-	7,7,7	1.18	1 (14%)	9,9,9	1.62	2 (22%)
2	SO4	A	501	-	4,4,4	0.45	0	6,6,6	0.09	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	TRS	A	503	-	-	3/9/9/9	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	503	TRS	C-N	2.38	1.57	1.49

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	503	TRS	C2-C-N	2.71	115.08	108.17
4	A	503	TRS	C2-C-C1	-2.14	104.95	110.66

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	503	TRS	C1-C-C3-O3
4	A	503	TRS	N-C-C3-O3
4	A	503	TRS	C2-C-C3-O3

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	503	TRS	2	0

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	401/469 (85%)	0.35	31 (7%) 19 21	27, 41, 74, 139	0

All (31) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	30	ALA	6.4
1	A	285	GLY	5.8
1	A	352	GLY	5.4
1	A	276	THR	5.2
1	A	275	ALA	5.0
1	A	367	ALA	4.9
1	A	444	PRO	4.8
1	A	350	THR	4.6
1	A	277	GLY	4.2
1	A	274	PRO	4.0
1	A	351	ASN	3.7
1	A	273	ASP	3.3
1	A	348	VAL	3.2
1	A	81	PHE	3.2
1	A	345	GLY	3.1
1	A	177	PRO	3.0
1	A	349	THR	3.0
1	A	175	ASN	2.9
1	A	140	LYS	2.7
1	A	126	VAL	2.7
1	A	286	ALA	2.7
1	A	178	SER	2.5
1	A	344	GLU	2.5
1	A	33	LYS	2.4
1	A	147	GLN	2.3
1	A	142	VAL	2.3
1	A	284	GLU	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	48	ARG	2.3
1	A	347	VAL	2.3
1	A	278	GLN	2.0
1	A	176	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	TRS	A	503	8/8	0.88	0.12	28,32,42,43	0
2	SO4	A	501	5/5	0.93	0.13	27,36,45,47	0
3	NA	A	502	1/1	0.96	0.14	45,45,45,45	0

6.5 Other polymers [i](#)

There are no such residues in this entry.