



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 06:59 AM UTC

PDB ID : 1U28 / pdb_00001u28
Title : R. rubrum transhydrogenase asymmetric complex (dI.NAD⁺)₂(dIII.NADP⁺)₁
Authors : Mather, O.C.; Singh, A.; van Boxel, G.I.; White, S.A.; Jackson, J.B.
Deposited on : 2004-07-16
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	6919 (2.30-2.30)
Ramachandran outliers	187476	6854 (2.30-2.30)
Sidechain outliers	187428	6854 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	384	
1	B	384	
2	C	203	

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 6974 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called NAD(P) transhydrogenase subunit alpha part 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	378	Total	C	N	O	S	0	0	0
			2779	1753	479	529	18			
1	B	359	Total	C	N	O	S	0	0	0
			2635	1666	458	495	16			

- Molecule 2 is a protein called NAD(P) transhydrogenase subunit beta.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	174	Total	C	N	O	S	0	0	0
			1311	830	217	253	11			

- Molecule 3 is NICOTINAMIDE-ADENINE-DINUCLEOTIDE (CCD ID: NAD) (formula: C₂₁H₂₇N₇O₁₄P₂).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	1
			52	27	8	15	2		
3	B	1	Total	C	N	O	P	0	0
			27	10	5	10	2		

- Molecule 4 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	C	O	0	0
			6	3	3		
4	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 5 is NADP NICOTINAMIDE-ADENINE-DINUCLEOTIDE PHOSPHATE (CCD ID: NAP) (formula: $C_{21}H_{28}N_7O_{17}P_3$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	C	1	Total 48	C 21	N 7	O 17	P 3	0	0

- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	39	Total O 39 39	0	0
6	B	64	Total O 64 64	0	0
6	C	7	Total O 7 7	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.


- Molecule 1: NAD(P) transhydrogenase subunit alpha part 1

Chain A: 




- Molecule 1: NAD(P) transhydrogenase subunit alpha part 1

Chain B: 



- Molecule 2: NAD(P) transhydrogenase subunit beta

Chain C: 



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	71.66Å 73.96Å 204.17Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.00 – 2.30	Depositor
% Data completeness (in resolution range)	(Not available) (49.00-2.30)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	0.08	Depositor
Refinement program	REFMAC 5.0, TNT	Depositor
R, R_{free}	0.241 , 0.275	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	6974	wwPDB-VP
Average B, all atoms (Å ²)	63.0	wwPDB-VP

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAD, NAP, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.58	0/2816	0.79	1/3816 (0.0%)
1	B	0.62	0/2668	0.78	0/3616
2	C	0.55	0/1334	0.78	1/1803 (0.1%)
All	All	0.59	0/6818	0.79	2/9235 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	375	VAL	N-CA-C	5.20	120.15	109.34
2	C	104	VAL	N-CA-CB	5.10	115.69	110.23

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2779	0	2904	10	0
1	B	2635	0	2777	13	0
2	C	1311	0	1303	23	0
3	A	52	0	12	0	0
3	B	27	0	12	0	0
4	B	12	0	16	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	C	48	0	25	1	0
6	A	39	0	0	0	0
6	B	64	0	0	1	0
6	C	7	0	0	0	0
All	All	6974	0	7049	46	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (46) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:142:ASN:HD21	1:B:186:GLN:HE21	1.10	0.91
2:C:53:PRO:HB2	2:C:95:MET:HE2	1.57	0.84
2:C:43:MET:HE3	2:C:49:VAL:HG11	1.68	0.76
2:C:43:MET:HE1	2:C:69:MET:HE3	1.72	0.71
1:B:142:ASN:ND2	1:B:186:GLN:HE21	1.90	0.65
2:C:43:MET:HE2	2:C:73:LEU:HD21	1.81	0.62
1:B:97:MET:HE1	1:B:343:LEU:HD13	1.81	0.61
1:A:122:MET:HA	1:A:122:MET:HE2	1.82	0.61
2:C:63:GLN:HG2	2:C:98:LEU:HB3	1.85	0.58
2:C:31:VAL:HG11	2:C:167:MET:HE2	1.87	0.56
2:C:43:MET:HE1	2:C:69:MET:CE	2.37	0.53
1:A:97:MET:HE1	1:A:343:LEU:HD13	1.91	0.53
1:B:362:THR:HG21	6:B:810:HOH:O	2.09	0.52
1:A:23:VAL:HG12	1:A:33:VAL:HG11	1.92	0.52
2:C:91:MET:HE2	2:C:94:HIS:HA	1.92	0.51
1:B:255:LEU:HD11	1:B:284:MET:HE2	1.93	0.51
2:C:162:PHE:CD1	2:C:186:MET:HE1	2.46	0.51
1:A:332:SER:HB2	1:A:333:PRO:HD3	1.93	0.50
1:B:156:PHE:CZ	1:B:259:ASP:HB3	2.46	0.50
2:C:126:PHE:CZ	2:C:163:ILE:HD11	2.47	0.49
2:C:194:MET:HA	2:C:197:GLN:HE21	1.78	0.48
2:C:95:MET:HE3	2:C:95:MET:HA	1.96	0.48
2:C:85:HIS:CG	2:C:133:VAL:HG22	2.47	0.48
2:C:41:PHE:CD1	2:C:41:PHE:C	2.91	0.48
1:B:10:ARG:HH11	1:B:10:ARG:HG3	1.80	0.47
1:B:162:MET:HE3	1:B:172:PRO:HD3	1.98	0.45
1:B:264:THR:HG22	1:B:293:LEU:HD12	1.98	0.45
2:C:57:MET:HE2	2:C:66:LEU:CD2	2.47	0.44
1:A:4:ALA:HA	1:A:34:ILE:O	2.17	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:106:ARG:N	1:A:107:PRO:CD	2.80	0.44
2:C:184:THR:HG22	2:C:186:MET:CE	2.47	0.44
1:A:156:PHE:CZ	1:A:259:ASP:HB3	2.53	0.44
2:C:140:THR:O	2:C:141:ASP:C	2.61	0.43
1:B:342:PHE:CZ	1:B:366:THR:HG21	2.53	0.43
1:B:97:MET:HE1	1:B:343:LEU:CD1	2.47	0.43
1:B:106:ARG:N	1:B:107:PRO:CD	2.81	0.43
2:C:31:VAL:HG12	2:C:180:PHE:CD1	2.55	0.42
2:C:191:ALA:HB2	5:C:400:NAP:C2A	2.50	0.42
1:A:222:ASP:HA	1:A:226:MET:HG3	2.02	0.41
2:C:65:ALA:HA	2:C:68:GLU:HG2	2.02	0.41
1:B:323:VAL:N	1:B:324:PRO:CD	2.83	0.41
2:C:61:GLN:HA	2:C:63:GLN:OE1	2.20	0.41
2:C:63:GLN:HE21	2:C:63:GLN:HB2	1.66	0.41
2:C:184:THR:HG22	2:C:186:MET:HE3	2.02	0.41
1:A:161:PRO:HG2	1:A:163:MET:HE2	2.02	0.40
1:A:60:THR:OG1	1:A:61:ALA:N	2.54	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	376/384 (98%)	350 (93%)	19 (5%)	7 (2%)	6	5
1	B	353/384 (92%)	344 (98%)	9 (2%)	0	100	100
2	C	172/203 (85%)	166 (96%)	6 (4%)	0	100	100
All	All	901/971 (93%)	860 (95%)	34 (4%)	7 (1%)	16	20

All (7) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	102	ALA
1	A	222	ASP
1	A	349	LYS
1	A	366	THR
1	A	224	GLU
1	A	358	LEU
1	A	375	VAL

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	293/296 (99%)	279 (95%)	14 (5%)	23	35
1	B	280/296 (95%)	260 (93%)	20 (7%)	13	19
2	C	138/154 (90%)	127 (92%)	11 (8%)	11	15
All	All	711/746 (95%)	666 (94%)	45 (6%)	16	23

All (45) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1	MET
1	A	50	LEU
1	A	70	VAL
1	A	75	GLN
1	A	92	GLU
1	A	122	MET
1	A	164	MET
1	A	176	LEU
1	A	199	MET
1	A	266	LEU
1	A	327	VAL
1	A	337	LYS
1	A	353	THR
1	A	362	THR
1	B	35	VAL
1	B	37	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	50	LEU
1	B	65	LEU
1	B	74	VAL
1	B	89	LEU
1	B	137	LEU
1	B	147	ARG
1	B	176	LEU
1	B	207	THR
1	B	245	LYS
1	B	246	LYS
1	B	247	GLN
1	B	281	VAL
1	B	289	VAL
1	B	327	VAL
1	B	339	LEU
1	B	358	LEU
1	B	362	THR
1	B	366	THR
2	C	38	ASP
2	C	63	GLN
2	C	91	MET
2	C	103	ASN
2	C	104	VAL
2	C	123	ASP
2	C	133	VAL
2	C	144	SER
2	C	164	LYS
2	C	180	PHE
2	C	203	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (17) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	37	GLN
1	A	75	GLN
1	A	105	ASN
1	A	142	ASN
1	A	186	GLN
1	A	313	HIS
1	A	322	ASN
1	A	338	ASN
1	B	67	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	142	ASN
1	B	247	GLN
1	B	338	ASN
2	C	45	ASN
2	C	63	GLN
2	C	131	ASN
2	C	197	GLN
2	C	203	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	NAD	B	600	-	28,29,48	1.52	4 (14%)	43,45,73	1.87	11 (25%)
4	GOL	B	800	-	5,5,5	0.37	0	5,5,5	0.27	0
3	NAD	A	500[A]	-	46,48,48	1.82	6 (13%)	64,73,73	1.57	10 (15%)
5	NAP	C	400	-	50,52,52	1.74	7 (14%)	71,80,80	1.61	11 (15%)
3	NAD	A	500[B]	-	46,48,48	1.78	6 (13%)	64,73,73	1.59	11 (17%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	GOL	B	700	-	5,5,5	0.37	0	5,5,5	0.34	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAD	B	600	-	-	3/16/32/62	0/3/3/5
4	GOL	B	800	-	-	2/4/4/4	-
3	NAD	A	500[A]	-	-	6/30/62/62	0/5/5/5
5	NAP	C	400	-	-	6/35/67/67	0/5/5/5
3	NAD	A	500[B]	-	-	4/30/62/62	0/5/5/5
4	GOL	B	700	-	-	2/4/4/4	-

All (23) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	500[A]	NAD	O7N-C7N	9.62	1.42	1.24
3	A	500[B]	NAD	O7N-C7N	9.24	1.41	1.24
5	C	400	NAP	O7N-C7N	8.77	1.40	1.24
3	B	600	NAD	C5A-C4A	4.82	1.47	1.39
5	C	400	NAP	PN-O3	3.37	1.63	1.59
5	C	400	NAP	PA-O3	3.16	1.62	1.59
3	B	600	NAD	PA-O3	3.13	1.62	1.59
3	B	600	NAD	C5A-C6A	2.93	1.49	1.41
3	A	500[A]	NAD	C2A-N3A	2.81	1.38	1.33
3	A	500[B]	NAD	C2A-N3A	2.81	1.38	1.33
3	A	500[A]	NAD	C2A-N1A	2.76	1.38	1.33
3	A	500[B]	NAD	C2A-N1A	2.76	1.38	1.33
5	C	400	NAP	C2A-N1A	2.72	1.38	1.33
5	C	400	NAP	C2A-N3A	2.70	1.38	1.33
3	B	600	NAD	C8A-N7A	2.54	1.36	1.31
3	A	500[A]	NAD	C2N-N1N	2.52	1.37	1.35
3	A	500[B]	NAD	C2N-N1N	2.49	1.37	1.35
3	A	500[A]	NAD	PA-O3	2.47	1.62	1.59
3	A	500[B]	NAD	PA-O3	2.47	1.62	1.59
5	C	400	NAP	C8A-N7A	2.30	1.36	1.31
3	A	500[A]	NAD	PN-O3	2.03	1.61	1.59
3	A	500[B]	NAD	PN-O3	2.03	1.61	1.59
5	C	400	NAP	C2N-N1N	2.03	1.37	1.35

All (43) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	C	400	NAP	N3A-C2A-N1A	-6.21	119.19	128.58
3	A	500[A]	NAD	N3A-C2A-N1A	-5.75	119.88	128.58
3	A	500[B]	NAD	N3A-C2A-N1A	-5.75	119.88	128.58
3	B	600	NAD	C5A-C4A-N3A	-5.08	119.72	126.72
3	A	500[A]	NAD	C5A-C4A-N3A	-4.40	120.66	126.72
3	A	500[B]	NAD	C5A-C4A-N3A	-4.40	120.66	126.72
5	C	400	NAP	N9A-C8A-N7A	-4.29	107.85	113.94
3	B	600	NAD	N3A-C4A-N9A	4.15	134.22	127.17
3	A	500[A]	NAD	N9A-C8A-N7A	-4.13	108.07	113.94
3	A	500[B]	NAD	N9A-C8A-N7A	-4.13	108.07	113.94
3	B	600	NAD	N3A-C2A-N1A	-4.07	122.42	128.58
3	B	600	NAD	C2A-N3A-C4A	3.82	121.15	111.83
5	C	400	NAP	C5A-C4A-N3A	-3.75	121.55	126.72
3	A	500[A]	NAD	C5A-N7A-C8A	3.65	109.19	103.45
3	A	500[B]	NAD	C5A-N7A-C8A	3.65	109.19	103.45
3	B	600	NAD	C4A-C5A-N7A	-3.51	106.57	110.58
5	C	400	NAP	O4B-C1B-N9A	3.50	114.81	108.09
3	A	500[A]	NAD	C2A-N3A-C4A	3.47	120.30	111.83
3	A	500[B]	NAD	C2A-N3A-C4A	3.47	120.30	111.83
3	B	600	NAD	C4A-N9A-C8A	3.38	109.29	105.74
5	C	400	NAP	C2A-N3A-C4A	3.36	120.05	111.83
5	C	400	NAP	C5A-N7A-C8A	3.35	108.72	103.45
3	A	500[A]	NAD	N3A-C4A-N9A	3.04	132.33	127.17
3	A	500[B]	NAD	N3A-C4A-N9A	3.04	132.33	127.17
5	C	400	NAP	O2A-PA-O3	2.99	115.35	107.27
3	B	600	NAD	C5A-N7A-C8A	2.82	107.88	103.45
3	A	500[A]	NAD	O4B-C1B-N9A	2.77	113.40	108.09
3	A	500[B]	NAD	O4B-C1B-N9A	2.77	113.40	108.09
3	B	600	NAD	N9A-C8A-N7A	-2.63	110.20	113.94
3	B	600	NAD	C6A-C5A-N7A	2.57	137.05	132.09
3	A	500[A]	NAD	C4A-C5A-N7A	-2.54	107.68	110.58
3	A	500[B]	NAD	C4A-C5A-N7A	-2.54	107.68	110.58
3	B	600	NAD	C2A-N1A-C6A	2.49	122.82	118.73
5	C	400	NAP	N3A-C4A-N9A	2.41	131.26	127.17
5	C	400	NAP	C2B-C1B-N9A	-2.31	109.95	113.75
3	A	500[B]	NAD	C3N-C7N-N7N	2.27	120.54	117.74
5	C	400	NAP	C4A-N9A-C8A	2.25	108.10	105.74
3	A	500[A]	NAD	C4A-N9A-C8A	2.21	108.06	105.74
3	A	500[B]	NAD	C4A-N9A-C8A	2.21	108.06	105.74
5	C	400	NAP	C4A-C5A-N7A	-2.13	108.14	110.58
3	A	500[A]	NAD	C5A-C6A-N6A	-2.09	118.11	123.29
3	A	500[B]	NAD	C5A-C6A-N6A	-2.09	118.11	123.29

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	600	NAD	O4B-C1B-N9A	2.05	112.02	108.09

There are no chirality outliers.

All (23) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	500[A]	NAD	C5D-O5D-PN-O2N
3	A	500[A]	NAD	O4D-C1D-N1N-C6N
3	A	500[B]	NAD	C5D-O5D-PN-O2N
4	B	800	GOL	O1-C1-C2-C3
5	C	400	NAP	O4D-C1D-N1N-C2N
5	C	400	NAP	O4D-C1D-N1N-C6N
5	C	400	NAP	C2D-C1D-N1N-C6N
4	B	700	GOL	C1-C2-C3-O3
4	B	800	GOL	O1-C1-C2-O2
4	B	700	GOL	O2-C2-C3-O3
5	C	400	NAP	PN-O3-PA-O1A
3	A	500[A]	NAD	C5B-O5B-PA-O1A
3	A	500[A]	NAD	C5D-O5D-PN-O3
3	A	500[A]	NAD	C5D-O5D-PN-O1N
3	A	500[B]	NAD	C5B-O5B-PA-O1A
3	A	500[B]	NAD	C5D-O5D-PN-O3
3	A	500[B]	NAD	C5D-O5D-PN-O1N
3	B	600	NAD	C5B-O5B-PA-O1A
3	B	600	NAD	C5B-O5B-PA-O2A
3	B	600	NAD	C5B-O5B-PA-O3
5	C	400	NAP	C2D-C1D-N1N-C2N
3	A	500[A]	NAD	O4D-C1D-N1N-C2N
5	C	400	NAP	PN-O3-PA-O2A


There are no ring outliers.


1 monomer is involved in 1 short contact:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	C	400	NAP	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be

highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.