



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 6, 2026 – 02:52 AM UTC

PDB ID : 4U33 / pdb_00004u33
Title : Structure of Mtb GlgE bound to maltose
Authors : Ronning, D.R.; Lindenberger, J.J.
Deposited on : 2014-07-18
Resolution : 3.29 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.29 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1169 (3.32-3.28)
Clashscore	190562	1209 (3.32-3.28)
Ramachandran outliers	187476	1188 (3.32-3.28)
Sidechain outliers	187428	1187 (3.32-3.28)
RSRZ outliers	180081	1169 (3.32-3.28)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	723	
1	B	723	
1	C	723	
1	D	723	
1	E	723	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	723	63% 24% 9%
2	G	2	50% 50%
2	H	2	50% 50%
2	I	2	50% 50%
2	J	2	50% 50%
2	K	2	50% 50%
2	L	2	50% 50%

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 31596 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Alpha-1,4-glucan:maltose-1-phosphate maltosyltransferase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	661	Total	C	N	O	S	0	0	0
			5243	3365	913	951	14			
1	B	661	Total	C	N	O	S	0	0	0
			5243	3365	913	951	14			
1	C	661	Total	C	N	O	S	0	0	0
			5243	3365	913	951	14			
1	D	661	Total	C	N	O	S	0	0	0
			5243	3365	913	951	14			
1	E	661	Total	C	N	O	S	0	0	0
			5243	3365	913	951	14			
1	F	661	Total	C	N	O	S	0	0	0
			5243	3365	913	951	14			

There are 132 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-21	MET	-	expression tag	UNP P9WQ16
A	-20	GLY	-	expression tag	UNP P9WQ16
A	-19	SER	-	expression tag	UNP P9WQ16
A	-18	SER	-	expression tag	UNP P9WQ16
A	-17	HIS	-	expression tag	UNP P9WQ16
A	-16	HIS	-	expression tag	UNP P9WQ16
A	-15	HIS	-	expression tag	UNP P9WQ16
A	-14	HIS	-	expression tag	UNP P9WQ16
A	-13	HIS	-	expression tag	UNP P9WQ16
A	-12	HIS	-	expression tag	UNP P9WQ16
A	-11	SER	-	expression tag	UNP P9WQ16
A	-10	SER	-	expression tag	UNP P9WQ16
A	-9	GLY	-	expression tag	UNP P9WQ16
A	-8	LEU	-	expression tag	UNP P9WQ16
A	-7	GLU	-	expression tag	UNP P9WQ16
A	-6	VAL	-	expression tag	UNP P9WQ16
A	-5	LEU	-	expression tag	UNP P9WQ16

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	-4	PHE	-	expression tag	UNP P9WQ16
A	-3	GLN	-	expression tag	UNP P9WQ16
A	-2	GLY	-	expression tag	UNP P9WQ16
A	-1	PRO	-	expression tag	UNP P9WQ16
A	0	HIS	-	expression tag	UNP P9WQ16
B	-21	MET	-	expression tag	UNP P9WQ16
B	-20	GLY	-	expression tag	UNP P9WQ16
B	-19	SER	-	expression tag	UNP P9WQ16
B	-18	SER	-	expression tag	UNP P9WQ16
B	-17	HIS	-	expression tag	UNP P9WQ16
B	-16	HIS	-	expression tag	UNP P9WQ16
B	-15	HIS	-	expression tag	UNP P9WQ16
B	-14	HIS	-	expression tag	UNP P9WQ16
B	-13	HIS	-	expression tag	UNP P9WQ16
B	-12	HIS	-	expression tag	UNP P9WQ16
B	-11	SER	-	expression tag	UNP P9WQ16
B	-10	SER	-	expression tag	UNP P9WQ16
B	-9	GLY	-	expression tag	UNP P9WQ16
B	-8	LEU	-	expression tag	UNP P9WQ16
B	-7	GLU	-	expression tag	UNP P9WQ16
B	-6	VAL	-	expression tag	UNP P9WQ16
B	-5	LEU	-	expression tag	UNP P9WQ16
B	-4	PHE	-	expression tag	UNP P9WQ16
B	-3	GLN	-	expression tag	UNP P9WQ16
B	-2	GLY	-	expression tag	UNP P9WQ16
B	-1	PRO	-	expression tag	UNP P9WQ16
B	0	HIS	-	expression tag	UNP P9WQ16
C	-21	MET	-	expression tag	UNP P9WQ16
C	-20	GLY	-	expression tag	UNP P9WQ16
C	-19	SER	-	expression tag	UNP P9WQ16
C	-18	SER	-	expression tag	UNP P9WQ16
C	-17	HIS	-	expression tag	UNP P9WQ16
C	-16	HIS	-	expression tag	UNP P9WQ16
C	-15	HIS	-	expression tag	UNP P9WQ16
C	-14	HIS	-	expression tag	UNP P9WQ16
C	-13	HIS	-	expression tag	UNP P9WQ16
C	-12	HIS	-	expression tag	UNP P9WQ16
C	-11	SER	-	expression tag	UNP P9WQ16
C	-10	SER	-	expression tag	UNP P9WQ16
C	-9	GLY	-	expression tag	UNP P9WQ16
C	-8	LEU	-	expression tag	UNP P9WQ16
C	-7	GLU	-	expression tag	UNP P9WQ16

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	-6	VAL	-	expression tag	UNP P9WQ16
C	-5	LEU	-	expression tag	UNP P9WQ16
C	-4	PHE	-	expression tag	UNP P9WQ16
C	-3	GLN	-	expression tag	UNP P9WQ16
C	-2	GLY	-	expression tag	UNP P9WQ16
C	-1	PRO	-	expression tag	UNP P9WQ16
C	0	HIS	-	expression tag	UNP P9WQ16
D	-21	MET	-	expression tag	UNP P9WQ16
D	-20	GLY	-	expression tag	UNP P9WQ16
D	-19	SER	-	expression tag	UNP P9WQ16
D	-18	SER	-	expression tag	UNP P9WQ16
D	-17	HIS	-	expression tag	UNP P9WQ16
D	-16	HIS	-	expression tag	UNP P9WQ16
D	-15	HIS	-	expression tag	UNP P9WQ16
D	-14	HIS	-	expression tag	UNP P9WQ16
D	-13	HIS	-	expression tag	UNP P9WQ16
D	-12	HIS	-	expression tag	UNP P9WQ16
D	-11	SER	-	expression tag	UNP P9WQ16
D	-10	SER	-	expression tag	UNP P9WQ16
D	-9	GLY	-	expression tag	UNP P9WQ16
D	-8	LEU	-	expression tag	UNP P9WQ16
D	-7	GLU	-	expression tag	UNP P9WQ16
D	-6	VAL	-	expression tag	UNP P9WQ16
D	-5	LEU	-	expression tag	UNP P9WQ16
D	-4	PHE	-	expression tag	UNP P9WQ16
D	-3	GLN	-	expression tag	UNP P9WQ16
D	-2	GLY	-	expression tag	UNP P9WQ16
D	-1	PRO	-	expression tag	UNP P9WQ16
D	0	HIS	-	expression tag	UNP P9WQ16
E	-21	MET	-	expression tag	UNP P9WQ16
E	-20	GLY	-	expression tag	UNP P9WQ16
E	-19	SER	-	expression tag	UNP P9WQ16
E	-18	SER	-	expression tag	UNP P9WQ16
E	-17	HIS	-	expression tag	UNP P9WQ16
E	-16	HIS	-	expression tag	UNP P9WQ16
E	-15	HIS	-	expression tag	UNP P9WQ16
E	-14	HIS	-	expression tag	UNP P9WQ16
E	-13	HIS	-	expression tag	UNP P9WQ16
E	-12	HIS	-	expression tag	UNP P9WQ16
E	-11	SER	-	expression tag	UNP P9WQ16
E	-10	SER	-	expression tag	UNP P9WQ16
E	-9	GLY	-	expression tag	UNP P9WQ16

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
E	-8	LEU	-	expression tag	UNP P9WQ16
E	-7	GLU	-	expression tag	UNP P9WQ16
E	-6	VAL	-	expression tag	UNP P9WQ16
E	-5	LEU	-	expression tag	UNP P9WQ16
E	-4	PHE	-	expression tag	UNP P9WQ16
E	-3	GLN	-	expression tag	UNP P9WQ16
E	-2	GLY	-	expression tag	UNP P9WQ16
E	-1	PRO	-	expression tag	UNP P9WQ16
E	0	HIS	-	expression tag	UNP P9WQ16
F	-21	MET	-	expression tag	UNP P9WQ16
F	-20	GLY	-	expression tag	UNP P9WQ16
F	-19	SER	-	expression tag	UNP P9WQ16
F	-18	SER	-	expression tag	UNP P9WQ16
F	-17	HIS	-	expression tag	UNP P9WQ16
F	-16	HIS	-	expression tag	UNP P9WQ16
F	-15	HIS	-	expression tag	UNP P9WQ16
F	-14	HIS	-	expression tag	UNP P9WQ16
F	-13	HIS	-	expression tag	UNP P9WQ16
F	-12	HIS	-	expression tag	UNP P9WQ16
F	-11	SER	-	expression tag	UNP P9WQ16
F	-10	SER	-	expression tag	UNP P9WQ16
F	-9	GLY	-	expression tag	UNP P9WQ16
F	-8	LEU	-	expression tag	UNP P9WQ16
F	-7	GLU	-	expression tag	UNP P9WQ16
F	-6	VAL	-	expression tag	UNP P9WQ16
F	-5	LEU	-	expression tag	UNP P9WQ16
F	-4	PHE	-	expression tag	UNP P9WQ16
F	-3	GLN	-	expression tag	UNP P9WQ16
F	-2	GLY	-	expression tag	UNP P9WQ16
F	-1	PRO	-	expression tag	UNP P9WQ16
F	0	HIS	-	expression tag	UNP P9WQ16

- Molecule 2 is an oligosaccharide called alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose.



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
2	G	2	Total	C	O	0	0	0
			23	12	11			

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
2	H	2	Total 23	C 12	O 11	0	0	0
2	I	2	Total 23	C 12	O 11	0	0	0
2	J	2	Total 23	C 12	O 11	0	0	0
2	K	2	Total 23	C 12	O 11	0	0	0
2	L	2	Total 23	C 12	O 11	0	0	0


3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Alpha-1,4-glucan:maltose-1-phosphate maltosyltransferase




- Molecule 1: Alpha-1,4-glucan:maltose-1-phosphate maltosyltransferase






• Molecule 1: Alpha-1,4-glucan:maltose-1-phosphate maltosyltransferase

Chain C: 59% 28% 9%




• Molecule 1: Alpha-1,4-glucan:maltose-1-phosphate maltosyltransferase


Chain D:  60% 27% 9%



• Molecule 1: Alpha-1,4-glucan:maltose-1-phosphate maltosyltransferase


Chain E:  59% 28% 5% 9%





- Molecule 1: Alpha-1,4-glucan:maltose-1-phosphate maltosyltransferase

Chain F: 63% 24% 9%



- Molecule 2: alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose

Chain G: 50% 50%



- Molecule 2: alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose

Chain H: 50% 50%



- Molecule 2: alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose

Chain I: 50% 50%



- Molecule 2: alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose

Chain J:  50% 50%



- Molecule 2: alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose

Chain K:  50% 50%



- Molecule 2: alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose

Chain L:  50% 50%



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	343.23Å 242.60Å 243.67Å 90.00° 135.15° 90.00°	Depositor
Resolution (Å)	47.64 – 3.29 47.64 – 3.29	Depositor EDS
% Data completeness (in resolution range)	99.0 (47.64-3.29) 90.4 (47.64-3.29)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.41 (at 3.33Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.9_1692)	Depositor
R, R_{free}	0.194 , 0.222 0.208 , 0.231	Depositor DCC
R_{free} test set	10542 reflections (4.97%)	wwPDB-VP
Wilson B-factor (Å ²)	81.2	Xtriage
Anisotropy	0.067	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 52.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.013 for -h-2*k,l,h+1 0.013 for -h-k-l,l,k 0.013 for -h+k-l,-l,-k 0.014 for -k+l,-h-l,-l 0.014 for k+l,h+l,-l 0.398 for k-l,h+l,-k 0.397 for h+k+l,-l,-h-l 0.397 for -k-l,-h-l,k 0.397 for h-k+l,l,-h-l 0.013 for h,-k,-h-l 0.457 for h+2*k,l,-k,-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	31596	wwPDB-VP
Average B, all atoms (Å ²)	71.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.38% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: GLC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.02	4/5407 (0.1%)	0.99	16/7395 (0.2%)
1	B	1.02	2/5407 (0.0%)	1.00	22/7395 (0.3%)
1	C	1.01	2/5407 (0.0%)	0.99	26/7395 (0.4%)
1	D	1.02	4/5407 (0.1%)	1.01	26/7395 (0.4%)
1	E	1.00	1/5407 (0.0%)	1.00	21/7395 (0.3%)
1	F	1.02	3/5407 (0.1%)	0.96	17/7395 (0.2%)
All	All	1.01	16/32442 (0.0%)	0.99	128/44370 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	C	0	1

The worst 5 of 16 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	487	ALA	CA-C	-6.79	1.44	1.52
1	F	487	ALA	CA-CB	-6.75	1.42	1.53
1	D	487	ALA	CA-C	-6.60	1.44	1.52
1	D	487	ALA	CA-CB	-6.46	1.43	1.53
1	A	487	ALA	CA-C	-6.42	1.45	1.52

The worst 5 of 128 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	184	ARG	N-CA-C	-11.14	99.22	111.36
1	A	184	ARG	N-CA-C	-9.66	100.75	111.28

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	196	ALA	N-CA-C	-9.19	100.46	112.68
1	B	92	LYS	CA-C-N	-8.68	111.20	120.66
1	B	92	LYS	C-N-CA	-8.68	111.20	120.66

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	C	194	ALA	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5243	0	5051	170	0
1	B	5243	0	5051	168	0
1	C	5243	0	5051	161	0
1	D	5243	0	5051	168	0
1	E	5243	0	5052	159	1
1	F	5243	0	5052	141	1
2	G	23	0	21	1	0
2	H	23	0	21	1	0
2	I	23	0	21	1	0
2	J	23	0	21	1	0
2	K	23	0	21	2	0
2	L	23	0	21	2	0
All	All	31596	0	30434	945	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 15.

The worst 5 of 945 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:275:TYR:OH	1:D:416:ARG:NH1	1.98	0.96

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:275:TYR:OH	1:E:416:ARG:NH1	2.01	0.94
1:B:275:TYR:OH	1:B:416:ARG:NH1	2.03	0.91
1:C:275:TYR:OH	1:C:416:ARG:NH1	2.02	0.91
1:E:188:ASP:HB2	1:E:191:THR:HG23	1.55	0.89

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:29:CYS:CB	1:E:29:CYS:SG[2_656]	1.82	0.38
1:F:29:CYS:CB	1:F:29:CYS:SG[2_555]	1.84	0.36

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	655/723 (91%)	632 (96%)	18 (3%)	5 (1%)	16	45
1	B	655/723 (91%)	630 (96%)	16 (2%)	9 (1%)	9	33
1	C	655/723 (91%)	632 (96%)	17 (3%)	6 (1%)	14	43
1	D	655/723 (91%)	624 (95%)	25 (4%)	6 (1%)	14	43
1	E	655/723 (91%)	632 (96%)	18 (3%)	5 (1%)	16	45
1	F	655/723 (91%)	631 (96%)	18 (3%)	6 (1%)	14	43
All	All	3930/4338 (91%)	3781 (96%)	112 (3%)	37 (1%)	14	43

5 of 37 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	151	SER
1	A	170	ARG
1	B	151	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	488	GLU
1	C	68	ARG

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	543/597 (91%)	490 (90%)	53 (10%)	7	28
1	B	543/597 (91%)	485 (89%)	58 (11%)	6	24
1	C	543/597 (91%)	498 (92%)	45 (8%)	10	34
1	D	543/597 (91%)	496 (91%)	47 (9%)	9	32
1	E	543/597 (91%)	492 (91%)	51 (9%)	8	30
1	F	543/597 (91%)	497 (92%)	46 (8%)	10	33
All	All	3258/3582 (91%)	2958 (91%)	300 (9%)	8	30

5 of 300 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	323	THR
1	F	447	GLU
1	E	464	THR
1	F	154	LEU
1	B	486	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 21 such sidechains are listed below:

Mol	Chain	Res	Type
1	E	318	HIS
1	E	485	GLN
1	F	685	ASN
1	F	318	HIS
1	E	455	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

12 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	GLC	G	1	2	12,12,12	0.58	0	17,17,17	0.86	0
2	GLC	G	2	2	11,11,12	0.96	0	15,15,17	0.72	0
2	GLC	H	1	2	12,12,12	0.56	0	17,17,17	1.05	1 (5%)
2	GLC	H	2	2	11,11,12	1.02	0	15,15,17	0.82	0
2	GLC	I	1	2	12,12,12	0.56	0	17,17,17	0.91	0
2	GLC	I	2	2	11,11,12	0.92	0	15,15,17	0.76	0
2	GLC	J	1	2	12,12,12	0.52	0	17,17,17	1.04	1 (5%)
2	GLC	J	2	2	11,11,12	0.91	0	15,15,17	0.73	0
2	GLC	K	1	2	12,12,12	0.51	0	17,17,17	1.04	1 (5%)
2	GLC	K	2	2	11,11,12	1.00	0	15,15,17	0.86	0
2	GLC	L	1	2	12,12,12	0.53	0	17,17,17	1.00	1 (5%)
2	GLC	L	2	2	11,11,12	0.97	0	15,15,17	0.86	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GLC	G	1	2	-	1/2/22/22	0/1/1/1
2	GLC	G	2	2	-	0/2/19/22	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GLC	H	1	2	-	0/2/22/22	0/1/1/1
2	GLC	H	2	2	-	2/2/19/22	0/1/1/1
2	GLC	I	1	2	-	0/2/22/22	0/1/1/1
2	GLC	I	2	2	-	2/2/19/22	0/1/1/1
2	GLC	J	1	2	-	0/2/22/22	0/1/1/1
2	GLC	J	2	2	-	1/2/19/22	0/1/1/1
2	GLC	K	1	2	-	0/2/22/22	0/1/1/1
2	GLC	K	2	2	-	2/2/19/22	0/1/1/1
2	GLC	L	1	2	-	0/2/22/22	0/1/1/1
2	GLC	L	2	2	-	0/2/19/22	0/1/1/1

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	H	1	GLC	C4-C3-C2	-2.45	106.53	110.83
2	K	1	GLC	C4-C3-C2	-2.19	106.99	110.83
2	J	1	GLC	C1-O5-C5	2.08	117.68	113.65
2	L	1	GLC	C4-C3-C2	-2.02	107.29	110.83

There are no chirality outliers.

5 of 8 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	H	2	GLC	O5-C5-C6-O6
2	H	2	GLC	C4-C5-C6-O6
2	I	2	GLC	O5-C5-C6-O6
2	K	2	GLC	O5-C5-C6-O6
2	G	1	GLC	O5-C5-C6-O6

There are no ring outliers.

7 monomers are involved in 8 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	J	1	GLC	1	0
2	H	1	GLC	1	0
2	I	1	GLC	1	0
2	K	2	GLC	1	0
2	L	1	GLC	2	0
2	K	1	GLC	1	0


Continued on next page...


Continued from previous page...


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	G	1	GLC	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.














5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	661/723 (91%)	-1.36	1 (0%) 91 86	47, 65, 120, 146	0
1	B	661/723 (91%)	-1.38	1 (0%) 91 86	46, 66, 124, 148	0
1	C	661/723 (91%)	-1.35	2 (0%) 90 82	46, 65, 119, 137	0
1	D	661/723 (91%)	-1.37	3 (0%) 87 76	45, 65, 123, 144	0
1	E	661/723 (91%)	-1.37	1 (0%) 91 86	47, 65, 119, 148	0
1	F	661/723 (91%)	-1.41	1 (0%) 91 86	47, 65, 114, 136	0
All	All	3966/4338 (91%)	-1.37	9 (0%) 91 86	45, 65, 120, 148	0

The worst 5 of 9 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	15	PRO	3.1
1	A	15	PRO	3.1
1	D	15	PRO	2.8
1	B	169	PRO	2.7
1	D	197	LEU	2.6

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	GLC	H	1	12/12	0.99	0.04	76,83,87,89	0
2	GLC	I	2	11/12	0.99	0.03	76,80,81,82	0
2	GLC	L	1	12/12	0.99	0.03	74,80,88,88	0
2	GLC	L	2	11/12	0.99	0.04	74,79,82,82	0
2	GLC	I	1	12/12	1.00	0.03	78,83,87,89	0
2	GLC	G	2	11/12	1.00	0.04	77,83,84,85	0
2	GLC	J	1	12/12	1.00	0.03	76,82,87,87	0
2	GLC	J	2	11/12	1.00	0.03	77,81,83,84	0
2	GLC	K	1	12/12	1.00	0.03	79,85,90,92	0
2	GLC	K	2	11/12	1.00	0.04	76,81,83,83	0
2	GLC	G	1	12/12	1.00	0.03	75,83,87,89	0
2	GLC	H	2	11/12	1.00	0.03	75,80,82,82	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around Chain H:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around Chain I:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around Chain J:

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around Chain K:**

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)





6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.