



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 04:03 AM UTC

PDB ID : 5U3K / pdb_00005u3k
Title : Crystal Structure of DH511.2 Fab in Complex with HIV-1 gp41 MPER 662-683 Peptide
Authors : Ofek, G.; Wu, L.; Loughheed, C.S.; Williams, L.D.; Nicely, N.I.; Haynes, B.F.
Deposited on : 2016-12-02
Resolution : 2.64 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.64 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2053 (2.66-2.62)
Clashscore	190562	2097 (2.66-2.62)
Ramachandran outliers	187476	2066 (2.66-2.62)
Sidechain outliers	187428	2066 (2.66-2.62)
RSRZ outliers	180081	2052 (2.66-2.62)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	232	11% 92% 6%
1	H	232	11% 92% 6%
2	B	214	15% 89% 7%
2	L	214	12% 92% 8%
3	C	28	7% 68% 32%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	P	28	21% 68% 29%

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 13491 atoms, of which 6516 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DH511.2 Fab Heavy Chain.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	H	222	Total	C	H	N	O	S	0	0	0
			3279	1064	1605	282	319	9			
1	A	218	Total	C	H	N	O	S	0	0	0
			3244	1053	1595	278	309	9			

- Molecule 2 is a protein called DH511.2 Fab Light Chain.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	L	214	Total	C	H	N	O	S	0	0	0
			3237	1026	1591	282	333	5			
2	B	199	Total	C	H	N	O	S	0	0	0
			2908	934	1409	255	305	5			

- Molecule 3 is a protein called gp41 MPER peptide.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	P	20	Total	C	H	N	O	0	0	0
			350	128	165	31	26			
3	C	19	Total	C	H	N	O	0	0	0
			325	120	151	28	26			

- Molecule 4 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	H	1	Total	Ca	0	0
			1	1		

- Molecule 5 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	Cl	0	0
			1	1		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	H	29	Total 29	O 29	0	0
6	L	31	Total 31	O 31	0	0
6	A	52	Total 52	O 52	0	0
6	B	27	Total 27	O 27	0	0
6	P	2	Total 2	O 2	0	0
6	C	5	Total 5	O 5	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: DH511.2 Fab Heavy Chain




• Molecule 1: DH511.2 Fab Heavy Chain




• Molecule 2: DH511.2 Fab Light Chain




• Molecule 2: DH511.2 Fab Light Chain






● Molecule 3: gp41 MPER peptide



● Molecule 3: gp41 MPER peptide



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	109.40Å 75.76Å 122.63Å 90.00° 94.29° 90.00°	Depositor
Resolution (Å)	40.76 – 2.64 40.76 – 2.64	Depositor EDS
% Data completeness (in resolution range)	96.6 (40.76-2.64) 87.7 (40.76-2.64)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.71 (at 2.65Å)	Xtriage
Refinement program	PHENIX 1.11.1 _2575	Depositor
R, R_{free}	0.232 , 0.283 0.235 , 0.280	Depositor DCC
R_{free} test set	1343 reflections (3.94%)	wwPDB-VP
Wilson B-factor (Å ²)	37.1	Xtriage
Anisotropy	0.707	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 50.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.89	EDS
Total number of atoms	13491	wwPDB-VP
Average B, all atoms (Å ²)	59.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.08% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CA, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.13	0/1692	0.30	0/2300
1	H	0.12	0/1717	0.29	0/2333
2	B	0.11	0/1529	0.30	0/2084
2	L	0.11	0/1681	0.29	0/2288
3	C	0.11	0/183	0.25	0/253
3	P	0.10	0/195	0.30	0/269
All	All	0.12	0/6997	0.29	0/9527

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1649	1595	1594	3	0
1	H	1674	1605	1604	7	0
2	B	1499	1409	1406	4	0
2	L	1646	1591	1591	9	1
3	C	174	151	151	0	0
3	P	185	165	165	1	0
4	H	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	A	1	0	0	0	0
6	A	52	0	0	1	0
6	B	27	0	0	0	0
6	C	5	0	0	0	0
6	H	29	0	0	0	0
6	L	31	0	0	0	0
6	P	2	0	0	0	0
All	All	6975	6516	6511	21	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (21) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:H:66:ARG:NH2	1:H:86:ASP:OD2	2.28	0.65
1:H:11:LEU:HD21	1:H:147:PRO:HB3	1.87	0.56
2:L:65:SER:OG	2:L:72:THR:OG1	2.24	0.56
2:L:6:GLN:O	2:L:100:GLN:NE2	2.39	0.55
2:B:167:ASP:OD1	2:B:168:SER:N	2.43	0.52
2:L:187:GLU:O	2:L:211:ARG:NH2	2.45	0.49
1:A:197:ASN:O	6:A:401:HOH:O	2.20	0.49
2:B:6:GLN:O	2:B:100:GLN:NE2	2.46	0.48
2:L:189:HIS:O	2:L:211:ARG:NE	2.44	0.48
2:L:167:ASP:OD1	2:L:168:SER:N	2.47	0.47
1:H:103:TRP:HB2	2:L:43:SER:OG	2.16	0.45
1:H:100(E):GLU:OE1	2:L:30:ARG:NH1	2.50	0.43
2:L:39:LYS:HG2	2:L:84:ALA:HB2	2.00	0.43
1:H:30:ASP:N	1:H:30:ASP:OD1	2.51	0.43
2:B:39:LYS:HG2	2:B:84:ALA:HB2	2.02	0.42
2:B:50:ALA:O	2:B:51:ALA:HB3	2.19	0.42
2:L:50:ALA:O	2:L:51:ALA:HB3	2.20	0.42
1:A:30:ASP:OD1	1:A:30:ASP:N	2.53	0.41
1:A:124:LEU:O	1:A:125:ALA:C	2.63	0.41
1:H:100(M):MET:N	1:H:100(M):MET:SD	2.94	0.41
1:H:100(I):PHE:HB3	3:P:676:THR:HG21	2.03	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:L:9:SER:OG	2:L:195:GLU:OE2[2_556]	2.19	0.01

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	212/232 (91%)	208 (98%)	4 (2%)	0	100	100
1	H	216/232 (93%)	212 (98%)	4 (2%)	0	100	100
2	B	191/214 (89%)	184 (96%)	7 (4%)	0	100	100
2	L	212/214 (99%)	205 (97%)	7 (3%)	0	100	100
3	C	17/28 (61%)	16 (94%)	1 (6%)	0	100	100
3	P	18/28 (64%)	17 (94%)	1 (6%)	0	100	100
All	All	866/948 (91%)	842 (97%)	24 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	176/194 (91%)	175 (99%)	1 (1%)	78	87
1	H	179/194 (92%)	178 (99%)	1 (1%)	78	87
2	B	167/189 (88%)	166 (99%)	1 (1%)	78	87
2	L	188/189 (100%)	188 (100%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
3	C	16/27 (59%)	16 (100%)	0	100	100
3	P	17/27 (63%)	17 (100%)	0	100	100
All	All	743/820 (91%)	740 (100%)	3 (0%)	84	91

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	H	30	ASP
1	A	30	ASP
2	B	90	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
2	L	160	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	218/232 (93%)	0.87	25 (11%) 9 8	28, 49, 89, 118	0
1	H	222/232 (95%)	0.93	25 (11%) 10 8	24, 58, 97, 177	1 (0%)
2	B	199/214 (92%)	1.03	33 (16%) 4 3	31, 59, 97, 137	0
2	L	214/214 (100%)	0.96	25 (11%) 9 7	33, 58, 91, 124	0
3	C	19/28 (67%)	1.29	2 (10%) 11 9	35, 49, 81, 99	0
3	P	20/28 (71%)	1.38	6 (30%) 1 0	42, 60, 71, 83	0
All	All	892/948 (94%)	0.96	116 (13%) 7 6	24, 57, 95, 177	1 (0%)

All (116) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	C	667	ALA	6.4
1	A	186	SER	5.9
2	B	208	SER	5.7
2	B	191	VAL	4.9
2	L	1	ASP	4.6
1	A	9	GLY	4.0
1	H	7	SER	3.9
2	B	152	ASN	3.8
1	H	16	GLY	3.7
2	B	143	GLU	3.7
1	H	210	ARG	3.7
2	L	10	SER	3.6
2	B	188	LYS	3.6
2	L	143	GLU	3.5
3	P	666	TRP	3.5
1	H	204	ASN	3.5
1	H	9	GLY	3.5
1	A	134	GLY	3.5
1	A	133	GLY	3.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	55	ALA	3.4
1	H	138	LEU	3.3
1	A	121	VAL	3.3
2	B	192	TYR	3.3
1	H	100(C)	PHE	3.3
2	B	147	GLN	3.2
2	L	43	SER	3.2
3	C	668	SER	3.2
1	H	12	VAL	3.1
1	A	184	VAL	3.1
1	H	11	LEU	3.1
1	H	193	THR	3.0
1	A	211	VAL	3.0
2	L	183	LYS	2.9
2	L	184	ALA	2.9
1	A	122	PHE	2.8
2	B	148	TRP	2.8
2	B	77	ASN	2.8
1	H	191	THR	2.8
2	L	194	CYS	2.8
2	B	194	CYS	2.8
1	H	127	SER	2.8
2	L	168	SER	2.8
1	A	94	MET	2.8
3	P	678	TRP	2.7
1	A	125	ALA	2.7
2	B	15	VAL	2.7
1	A	165	THR	2.6
2	B	154	LEU	2.6
2	B	16	GLY	2.6
1	H	28	PHE	2.6
1	H	126	PRO	2.6
2	L	149	LYS	2.6
2	L	150	VAL	2.6
2	B	129	THR	2.5
1	H	206	LYS	2.5
2	L	47	LEU	2.5
1	A	135	THR	2.5
1	A	7	SER	2.5
1	A	1	GLN	2.5
2	L	95(A)	PRO	2.5
2	L	40	PRO	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	H	156	SER	2.4
2	B	43	SER	2.4
2	B	64	GLY	2.4
2	B	209	PHE	2.4
2	B	18	ARG	2.4
2	B	124	GLN	2.4
1	A	159	LEU	2.4
1	H	192	GLN	2.4
2	L	209	PHE	2.4
1	A	212	GLU	2.3
2	L	169	LYS	2.3
2	B	151	ASP	2.3
2	L	24	ARG	2.3
1	A	194	TYR	2.3
2	B	132	VAL	2.3
2	L	131	SER	2.3
2	B	42	GLY	2.3
1	A	160	THR	2.2
2	B	121	SER	2.2
2	B	92	TYR	2.2
1	A	185	PRO	2.2
1	H	105	ARG	2.2
2	B	1	ASP	2.2
2	B	14	SER	2.2
3	P	676	THR	2.2
1	A	213	PRO	2.2
2	B	56	THR	2.1
1	H	3	GLN	2.1
1	A	42	GLY	2.1
2	L	191	VAL	2.1
3	P	680	TRP	2.1
2	L	103	LYS	2.1
1	A	28	PHE	2.1
2	L	190	LYS	2.1
2	B	190	LYS	2.1
1	H	1	GLN	2.1
1	H	190	GLY	2.1
2	L	140	TYR	2.1
2	L	48	ILE	2.1
3	P	670	TRP	2.1
2	B	72	THR	2.1
1	H	189	LEU	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	78	LEU	2.1
1	A	208	ASP	2.0
2	B	105	ASP	2.0
2	B	142	ARG	2.0
1	H	183	THR	2.0
2	L	147	GLN	2.0
2	L	158	ASN	2.0
1	H	114	ALA	2.0
1	H	82(B)	THR	2.0
2	B	54	LEU	2.0
1	A	8	GLY	2.0
2	L	42	GLY	2.0
3	P	681	TYR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	CA	H	301	1/1	0.91	0.05	36,36,36,36	0
5	CL	A	301	1/1	0.99	0.06	26,26,26,26	0

6.5 Other polymers [i](#)

There are no such residues in this entry.