



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 12, 2026 – 06:51 AM UTC

PDB ID : 5U43 / pdb_00005u43
Title : Human PPARdelta ligand-binding domain in complexed with specific agonist 12
Authors : Wu, C.-C.; Baiga, T.J.; Downes, M.; La Clair, J.J.; Atkins, A.R.; Richard, S.B.; Stockley-Noel, T.A.; Bowman, M.E.; Evans, R.M.; Noel, J.P.
Deposited on : 2016-12-03
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7789 (1.90-1.90)
Clashscore	190562	8410 (1.90-1.90)
Ramachandran outliers	187476	8333 (1.90-1.90)
Sidechain outliers	187428	8333 (1.90-1.90)
RSRZ outliers	180081	7790 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	272	8% 91% 6%
1	B	272	11% 87% 9%

2 Entry composition [i](#)


There are 6 unique types of molecules in this entry. The entry contains 9398 atoms, of which 4639 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Peroxisome proliferator-activated receptor delta.


Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	266	Total	C	H	N	O	S	0	16	0
			4449	1424	2254	370	392	9			
1	B	262	Total	C	H	N	O	S	0	5	0
			4289	1374	2171	355	378	11			

- Molecule 2 is 6-(2-([cyclopropyl(4'-methoxy[1,1'-biphenyl]-4-carbonyl)amino)methyl}phenoxy)hexanoic acid (CCD ID: 7SS) (formula: C₃₀H₃₃NO₅).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	H	N	O	0	0
			68	30	32	1	5		
2	B	1	Total	C	H	N	O	0	0
			68	30	32	1	5		

- Molecule 3 is S-1,2-PROPANEDIOL (CCD ID: PGO) (formula: C₃H₈O₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	H	O	0	1
			26	6	16	4		
3	B	1	Total	C	H	O	0	0
			13	3	8	2		
3	B	1	Total	C	H	O	0	0
			13	3	8	2		

- Molecule 4 is DI(HYDROXYETHYL)ETHER (CCD ID: PEG) (formula: $C_4H_{10}O_3$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	H	O	0	1
			34	8	20	6		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	H	O	0	0
			17	4	10	3		
4	B	1	Total	C	H	O	0	0
			17	4	10	3		

- Molecule 5 is heptyl beta-D-glucopyranoside (CCD ID: B7G) (formula: C₁₃H₂₆O₆).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	H	O	0	0
			45	13	26	6		
5	A	1	Total	C	H	O	0	0
			45	13	26	6		
5	B	1	Total	C	H	O	0	0
			45	13	26	6		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	163	Total	O	0	0
			163	163		
6	B	106	Total	O	0	0
			106	106		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Peroxisome proliferator-activated receptor delta



- Molecule 1: Peroxisome proliferator-activated receptor delta



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	39.64Å 95.01Å 96.32Å 90.00° 97.60° 90.00°	Depositor
Resolution (Å)	35.41 – 1.90 35.41 – 1.90	Depositor EDS
% Data completeness (in resolution range)	97.0 (35.41-1.90) 97.0 (35.41-1.90)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.10 (at 1.81Å)	Xtriage
Refinement program	PHENIX (1.10.1_2155: ???)	Depositor
R, R_{free}	0.186 , 0.238 0.189 , 0.242	Depositor DCC
R_{free} test set	1846 reflections (2.82%)	wwPDB-VP
Wilson B-factor (Å ²)	23.9	Xtriage
Anisotropy	0.510	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.43 , 55.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	9398	wwPDB-VP
Average B, all atoms (Å ²)	42.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 10.21% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PGO, 7SS, PEG, B7G

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.60	0/2316	0.69	1/3131 (0.0%)
1	B	0.52	0/2202	0.62	0/2973
All	All	0.56	0/4518	0.66	1/6104 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	286	VAL	CG1-CB-CG2	6.63	125.39	110.80

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2195	2254	2171	9	2
1	B	2118	2171	2127	13	2
2	A	36	32	0	0	0
2	B	36	32	0	0	0
3	A	10	16	16	0	0
3	B	10	16	16	0	0
4	A	21	30	30	0	0
4	B	7	10	10	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	A	38	52	52	1	0
5	B	19	26	26	0	0
6	A	163	0	0	1	0
6	B	106	0	0	0	0
All	All	4759	4639	4448	22	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (22) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:249:CYS:O	1:A:253:THR:HG23	2.00	0.62
1:B:249:CYS:O	1:B:253:THR:HG23	2.07	0.54
1:A:251:CYS:SG	5:A:505:B7G:H82	2.50	0.52
1:A:375:ASP:O	1:A:379:ARG:HG3	2.10	0.51
1:B:216:ILE:HG12	1:B:241:ILE:HD12	1.95	0.49
1:B:220:TRP:CZ2	1:B:224:LYS:HD2	2.48	0.48
1:B:216:ILE:HG12	1:B:241:ILE:CD1	2.44	0.46
1:B:396:PHE:HB3	1:B:397:PRO:HD3	1.98	0.45
1:A:265:LYS:NZ	6:A:604:HOH:O	2.43	0.45
1:A:219:LEU:C	1:A:219:LEU:HD23	2.42	0.44
1:B:288:GLU:OE2	1:B:407:ARG:HG2	2.17	0.44
1:B:385:LEU:HD11	1:B:399:LEU:HD12	2.00	0.44
1:A:230:GLN:HG2	1:A:251:CYS:SG	2.57	0.44
1:B:324:PHE:O	1:B:327:ILE:HG22	2.20	0.42
1:B:327:ILE:HD11	1:B:417:MET:SD	2.59	0.42
1:A:329:GLU:N	1:A:330:PRO:CD	2.82	0.42
1:B:233:ASN:OD1	1:B:233:ASN:N	2.50	0.41
1:B:241:ILE:HD11	1:B:316:PHE:HZ	1.85	0.41
1:B:430:HIS:O	1:B:434:GLN:HG3	2.20	0.41
1:A:315[A]:GLU:OE2	1:A:318[A]:ARG:NH1	2.54	0.41
1:B:236:PRO:O	1:B:237:PRO:C	2.64	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:382:GLU:OE1	1:B:238:TYR:HH[2_547]	1.51	0.09
1:A:382:GLU:OE1	1:B:238:TYR:OH[2_547]	2.16	0.04

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	277/272 (102%)	274 (99%)	3 (1%)	0	100	100
1	B	263/272 (97%)	255 (97%)	8 (3%)	0	100	100
All	All	540/544 (99%)	529 (98%)	11 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	248/240 (103%)	244 (98%)	4 (2%)	55	54
1	B	237/240 (99%)	234 (99%)	3 (1%)	61	61
All	All	485/480 (101%)	478 (99%)	7 (1%)	57	59

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	286	VAL
1	A	303	LEU
1	A	426	GLU
1	A	439	ASP
1	B	198	ARG
1	B	303	LEU
1	B	352	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	191	ASN
1	A	374	GLN
1	A	386	GLN
1	A	408	GLN
1	A	418	GLN
1	B	374	GLN
1	B	408	GLN
1	B	418	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

13 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	B7G	B	501	-	19,19,19	0.92	0	24,24,24	1.39	4 (16%)
4	PEG	A	503[B]	-	6,6,6	0.61	0	5,5,5	0.28	0
4	PEG	B	504	-	6,6,6	0.58	0	5,5,5	0.47	0
2	7SS	A	501	-	39,39,39	0.93	3 (7%)	52,52,52	1.39	8 (15%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	B7G	A	505	-	19,19,19	0.97	1 (5%)	24,24,24	1.22	1 (4%)
3	PGO	A	502[A]	-	4,4,4	0.67	0	4,4,4	0.50	0
5	B7G	A	506	-	19,19,19	0.86	1 (5%)	24,24,24	1.32	2 (8%)
3	PGO	B	503	-	4,4,4	0.57	0	4,4,4	0.62	0
3	PGO	A	502[B]	-	4,4,4	0.65	0	4,4,4	0.60	0
4	PEG	A	503[A]	-	6,6,6	0.54	0	5,5,5	0.49	0
3	PGO	B	505	-	4,4,4	0.53	0	4,4,4	0.69	0
4	PEG	A	504	-	6,6,6	0.51	0	5,5,5	0.25	0
2	7SS	B	502	-	39,39,39	0.83	1 (2%)	52,52,52	1.81	6 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	B7G	B	501	-	-	2/10/30/30	0/1/1/1
4	PEG	A	503[B]	-	-	2/4/4/4	-
4	PEG	B	504	-	-	2/4/4/4	-
2	7SS	A	501	-	-	4/31/33/33	0/4/4/4
5	B7G	A	505	-	-	8/10/30/30	0/1/1/1
3	PGO	A	502[A]	-	-	2/2/2/2	-
5	B7G	A	506	-	-	3/10/30/30	0/1/1/1
3	PGO	B	503	-	-	2/2/2/2	-
3	PGO	A	502[B]	-	-	2/2/2/2	-
4	PEG	A	503[A]	-	-	1/4/4/4	-
3	PGO	B	505	-	-	1/2/2/2	-
4	PEG	A	504	-	-	2/4/4/4	-
2	7SS	B	502	-	-	4/31/33/33	0/4/4/4

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	502	7SS	C16-C20	2.38	1.42	1.38
5	A	505	B7G	O5-C1	2.28	1.47	1.41
2	A	501	7SS	C23-C13	2.22	1.53	1.48
2	A	501	7SS	C14-N	-2.22	1.29	1.34
2	A	501	7SS	C16-C15	-2.12	1.36	1.39
5	A	506	B7G	O5-C1	2.05	1.47	1.41

All (21) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	502	7SS	C23-C13-N	-8.96	107.32	118.46
5	A	505	B7G	C7-O1-C1	4.30	121.03	113.68
5	A	506	B7G	O1-C1-C2	3.96	114.28	108.27
5	B	501	B7G	O1-C1-C2	3.77	114.00	108.27
2	A	501	7SS	C23-C13-N	-3.66	113.91	118.46
2	A	501	7SS	C6-N-C14	3.51	126.72	116.97
2	B	502	7SS	C6-N-C14	3.48	126.65	116.97
2	B	502	7SS	O3-C14-C15	-3.23	113.90	120.29
2	A	501	7SS	C13-N-C14	-3.16	113.66	123.21
2	A	501	7SS	C15-C14-N	3.04	124.74	118.72
2	B	502	7SS	C15-C14-N	3.00	124.66	118.72
2	B	502	7SS	C13-N-C14	-2.95	114.32	123.21
5	B	501	B7G	C6-C5-C4	-2.65	106.50	113.02
2	A	501	7SS	O3-C14-C15	-2.53	115.28	120.29
5	A	506	B7G	O5-C5-C4	2.26	113.77	109.70
2	A	501	7SS	C16-C20-C19	-2.25	118.23	121.12
2	A	501	7SS	C16-C15-C17	2.24	121.41	118.57
5	B	501	B7G	O2-C2-C1	-2.15	104.96	110.08
2	B	502	7SS	C4-C5-C	2.13	121.29	118.37
2	A	501	7SS	C16-C15-C14	-2.04	115.06	120.28
5	B	501	B7G	O5-C5-C4	2.03	113.36	109.70

There are no chirality outliers.

All (35) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	501	7SS	C22-C13-N-C6
2	A	501	7SS	C22-C13-N-C14
2	A	501	7SS	C23-C13-N-C14
2	B	502	7SS	C22-C13-N-C6
2	B	502	7SS	C22-C13-N-C14
2	B	502	7SS	C23-C13-N-C14
3	A	502[A]	PGO	O1-C1-C2-C3
3	A	502[A]	PGO	O1-C1-C2-O2
3	A	502[B]	PGO	O1-C1-C2-C3
3	A	502[B]	PGO	O1-C1-C2-O2
5	A	505	B7G	O5-C1-O1-C7
5	A	505	B7G	C4-C5-C6-O6
5	A	505	B7G	C2-C1-O1-C7
5	A	505	B7G	O5-C5-C6-O6
5	A	506	B7G	O1-C7-C8-C9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
4	A	503[B]	PEG	O1-C1-C2-O2
4	A	503[A]	PEG	O1-C1-C2-O2
5	A	506	B7G	C7-C8-C9-C10
4	A	503[B]	PEG	O2-C3-C4-O4
5	B	501	B7G	C9-C10-C11-C12
5	A	505	B7G	C7-C8-C9-C10
4	A	504	PEG	O1-C1-C2-O2
5	A	505	B7G	C11-C10-C9-C8
5	A	505	B7G	O1-C7-C8-C9
5	A	506	B7G	C4-C5-C6-O6
5	A	505	B7G	C10-C11-C12-C13
3	B	503	PGO	O1-C1-C2-C3
4	B	504	PEG	C1-C2-O2-C3
4	A	504	PEG	C1-C2-O2-C3
3	B	503	PGO	O1-C1-C2-O2
2	A	501	7SS	C23-C13-N-C6
5	B	501	B7G	C7-C8-C9-C10
2	B	502	7SS	C7-C8-C9-C10
4	B	504	PEG	O1-C1-C2-O2
3	B	505	PGO	O1-C1-C2-O2


There are no ring outliers.

1 monomer is involved in 1 short contact:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	505	B7G	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.


Ligand B7G B 501




Ligand 7SS A 501




Ligand B7G A 505



Ligand B7G A 506





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	266/272 (97%)	0.25	22 (8%) 17 18	11, 33, 73, 101	6 (2%)
1	B	262/272 (96%)	0.62	31 (11%) 9 9	15, 39, 83, 144	1 (0%)
All	All	528/544 (97%)	0.43	53 (10%) 12 13	11, 35, 80, 144	7 (1%)

All (53) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	394	TYR	5.6
1	B	203	GLY	5.2
1	A	171	GLN	4.8
1	A	181	HIS	4.7
1	B	439	ASP	4.5
1	B	394	TYR	4.4
1	B	426	GLU	4.2
1	B	234	GLY	4.1
1	B	241	ILE	4.1
1	B	392	ALA	4.0
1	B	268	PRO	3.8
1	A	172	VAL	3.5
1	A	439	ASP	3.5
1	B	209	ALA	3.5
1	A	170	PRO	3.5
1	B	191	ASN	3.5
1	A	383	PHE	3.3
1	A	204	LYS	3.1
1	B	387	ALA	3.1
1	B	195	LYS	3.1
1	B	419	ARG	3.0
1	B	418	GLN	2.8
1	B	440	MET	2.8
1	B	393	GLN	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	176	LYS	2.7
1	B	232	VAL	2.7
1	B	307	ASN	2.6
1	B	383	PHE	2.5
1	A	177	ALA	2.5
1	A	387	ALA	2.5
1	B	235	LEU	2.5
1	A	309[A]	SER	2.5
1	A	429[A]	LEU	2.5
1	A	209	ALA	2.5
1	B	434	GLN	2.4
1	A	233	ASN	2.3
1	B	265	LYS	2.3
1	A	382	GLU	2.3
1	B	258	ARG	2.2
1	A	386	GLN	2.2
1	A	175	LEU	2.2
1	A	173	ALA	2.2
1	A	258	ARG	2.1
1	B	239[A]	LYS	2.1
1	A	393	GLN	2.1
1	B	251	CYS	2.1
1	B	357	LEU	2.1
1	A	326	ASP	2.1
1	B	369	ARG	2.1
1	B	229	LYS	2.0
1	A	229	LYS	2.0
1	B	267	ILE	2.0
1	B	233	ASN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

6.4 Ligands




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	B7G	A	505	19/19	0.80	0.22	45,71,125,145	0
4	PEG	A	503[B]	7/7	0.83	0.23	41,58,69,69	17
4	PEG	A	503[A]	7/7	0.83	0.23	29,54,70,70	17
4	PEG	B	504	7/7	0.84	0.28	70,85,113,113	0
3	PGO	B	503	5/5	0.86	0.17	51,67,94,113	0
3	PGO	B	505	5/5	0.86	0.14	49,65,73,84	0
3	PGO	A	502[A]	5/5	0.87	0.16	32,44,53,53	13
3	PGO	A	502[B]	5/5	0.87	0.16	38,46,51,52	13
4	PEG	A	504	7/7	0.90	0.15	48,62,116,116	0
5	B7G	A	506	19/19	0.92	0.11	28,44,56,66	0
2	7SS	B	502	36/36	0.93	0.10	18,37,70,84	0
5	B7G	B	501	19/19	0.93	0.10	29,41,62,63	0
2	7SS	A	501	36/36	0.96	0.06	15,24,38,38	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around B7G A 505:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around B7G A 506:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around 7SS B 502:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around B7G B 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around 7SS A 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.