



wwPDB X-ray Structure Validation Summary Report ⓘ

Apr 22, 2026 – 11:24 AM EDT

PDB ID : 7U40 / pdb_00007u40
Title : [L244] Self-assembling tensegrity triangle with two turns, four turns and four turns of DNA per axis by linker addition with P1 symmetry
Authors : Woloszyn, K.; Vecchioni, S.; Seeman, N.C.; Sha, R.; Ohayon, Y.P.
Deposited on : 2022-02-28
Resolution : 7.55 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	FAILED
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 7.55 Å.







Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	1000 (10.00-4.06)


The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Note EDS failed to run properly.

Mol	Chain	Length	Quality of chain
1	A	21	 24% 76%
2	E	14	 21% 79%
3	D	14	 36% 64%
4	B	21	 43% 57%
5	F	14	 14% 86%
6	C	21	 24% 76%
7	M	21	 38% 62%
8	X	21	 24% 76%
9	Y	21	 38% 62%
10	U	21	 38% 62%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
11	V	21	 A horizontal bar chart showing the quality of chain 11. The bar is divided into two segments: a green segment on the left representing 24% and a yellow segment on the right representing 76%. The percentages are labeled below the bar.

2 Entry composition [i](#)

There are 11 unique types of molecules in this entry. The entry contains 4299 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a DNA chain called DNA (5'-D(P*CP*AP*CP*GP*AP*GP*CP*CP*TP*GP*AP*TP*CP*GP*GP*AP*CP*AP*AP*GP*A)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	21	Total	C	N	O	P	0	0	0
			433	204	87	121	21			

- Molecule 2 is a DNA chain called DNA (5'-D(*TP*TP*AP*GP*TP*CP*GP*TP*GP*GP*CP*TP*CP*G)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	E	14	Total	C	N	O	P	0	0	0
			285	137	49	86	13			

- Molecule 3 is a DNA chain called DNA (5'-D(P*GP*AP*TP*CP*TP*TP*GP*TP*GP*GP*CP*TP*GP*C)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	D	14	Total	C	N	O	P	0	0	0
			288	137	49	88	14			

- Molecule 4 is a DNA chain called DNA (5'-D(P*AP*GP*GP*CP*AP*GP*CP*CP*TP*GP*TP*AP*CP*GP*GP*AP*CP*AP*TP*CP*A)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	B	21	Total	C	N	O	P	0	0	0
			432	204	84	123	21			

- Molecule 5 is a DNA chain called DNA (5'-D(P*AP*CP*TP*GP*AP*TP*GP*TP*GP*GP*TP*AP*GP*G)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
5	F	14	Total	C	N	O	P	0	0	0
			294	139	56	85	14			

- Molecule 6 is a DNA chain called DNA (5'-D(*AP*AP*CP*CP*TP*AP*CP*CP*TP*GP*GP*CP*AP*GP*GP*AP*CP*GP*AP*CP*T)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
6	C	21	Total	C	N	O	P	0	0	0
			426	203	82	121	20			

- Molecule 7 is a DNA chain called DNA (5'-D(P*TP*CP*AP*CP*CP*TP*GP*CP*CP*AP*CP*CP*GP*TP*AP*CP*AP*CP*CP*GP*A)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
7	M	21	Total	C	N	O	P	0	0	0
			421	200	76	124	21			

- Molecule 8 is a DNA chain called DNA (5'-D(P*GP*TP*CP*TP*CP*TP*GP*CP*TP*AP*GP*AP*TP*CP*GP*TP*CP*AP*GP*CP*A)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
8	X	21	Total	C	N	O	P	0	0	0
			428	204	75	128	21			

- Molecule 9 is a DNA chain called DNA (5'-D(P*CP*TP*TP*GP*CP*TP*GP*AP*CP*GP*AP*TP*CP*TP*AP*GP*CP*AP*GP*AP*G)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
9	Y	21	Total	C	N	O	P	0	0	0
			432	205	80	126	21			

- Molecule 10 is a DNA chain called DNA (5'-D(P*TP*GP*CP*TP*AP*GP*GP*CP*TP*A*P*AP*CP*TP*CP*GP*CP*TP*AP*GP*CP*G)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
10	U	21	Total	C	N	O	P	0	0	0
			430	204	78	127	21			

- Molecule 11 is a DNA chain called DNA (5'-D(P*TP*CP*CP*GP*CP*TP*AP*GP*CP*G*P*AP*GP*TP*TP*AP*GP*CP*CP*TP*AP*G)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
11	V	21	Total	C	N	O	P	0	0	0
			430	204	78	127	21			


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS failed to run properly.


- Molecule 1: DNA (5'-D(P*CP*AP*CP*GP*AP*GP*CP*CP*TP*GP*AP*TP*CP*GP*GP*AP*CP*AP*AP*GP*A)-3')

Chain A: 



- Molecule 2: DNA (5'-D(*TP*TP*AP*GP*TP*CP*GP*TP*GP*GP*CP*TP*CP*G)-3')

Chain E: 




- Molecule 3: DNA (5'-D(P*GP*AP*TP*CP*TP*TP*GP*TP*GP*GP*CP*TP*GP*C)-3')

Chain D: 



- Molecule 4: DNA (5'-D(P*AP*GP*GP*CP*AP*GP*CP*CP*TP*GP*TP*AP*CP*GP*GP*AP*CP*AP*TP*CP*A)-3')

Chain B: 




- Molecule 5: DNA (5'-D(P*AP*CP*TP*GP*AP*TP*GP*TP*GP*GP*TP*AP*GP*G)-3')

Chain F: 



- Molecule 6: DNA (5'-D(*AP*AP*CP*CP*TP*AP*CP*CP*TP*GP*GP*CP*AP*GP*GP*AP*CP*GP*AP*CP*T)-3')

Chain C:  24% 76%



- Molecule 7: DNA (5'-D(P*TP*CP*AP*CP*CP*TP*GP*CP*CP*AP*CP*CP*GP*TP*AP*CP*AP*CP*CP*GP*A)-3')

Chain M:  38% 62%





- Molecule 8: DNA (5'-D(P*GP*TP*CP*TP*CP*TP*GP*CP*TP*AP*GP*AP*TP*CP*GP*TP*CP*AP*GP*CP*A)-3')

Chain X:  24% 76%




- Molecule 9: DNA (5'-D(P*CP*TP*TP*GP*CP*TP*GP*AP*CP*GP*AP*TP*CP*TP*AP*GP*CP*AP*GP*AP*G)-3')

Chain Y:  38% 62%





- Molecule 10: DNA (5'-D(P*TP*GP*CP*TP*AP*GP*GP*CP*TP*AP*AP*CP*TP*CP*GP*CP*TP*AP*GP*CP*G)-3')

Chain U:  38% 62%



- Molecule 11: DNA (5'-D(P*TP*CP*CP*GP*CP*TP*AP*GP*CP*GP*AP*GP*TP*TP*AP*GP*CP*CP*TP*AP*G)-3')

Chain V:  24% 76%



4 Data and refinement statistics

EDS failed to run properly - this section is therefore incomplete.

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	67.64Å 135.54Å 136.11Å 100.06° 103.07° 96.36°	Depositor
Resolution (Å)	63.63 – 7.55	Depositor
% Data completeness (in resolution range)	59.1 (63.63-7.55)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.97 (at 7.41Å)	Xtriage
Refinement program	PHENIX 1.18.2_3874	Depositor
R, R_{free}	0.161 , 0.208	Depositor
Wilson B-factor (Å ²)	195.7	Xtriage
Anisotropy	0.273	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.38$, $\langle L^2 \rangle = 0.20$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	4299	wwPDB-VP
Average B, all atoms (Å ²)	939.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 18.38% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.29	0/487	0.57	0/749
2	E	0.31	0/318	0.70	0/490
3	D	0.28	0/321	0.63	0/494
4	B	0.28	0/485	0.53	0/746
5	F	0.32	0/330	0.66	0/509
6	C	0.27	0/478	0.51	0/735
7	M	0.21	0/470	0.41	0/720
8	X	0.28	0/478	0.57	0/735
9	Y	0.28	0/484	0.46	0/745
10	U	0.25	0/481	0.52	0/740
11	V	0.26	0/481	0.50	0/740
All	All	0.27	0/4813	0.55	0/7403

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	433	0	234	16	0
2	E	285	0	161	9	0
3	D	288	0	160	8	0
4	B	432	0	235	14	0
5	F	294	0	159	17	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	C	426	0	236	14	0
7	M	421	0	235	7	0
8	X	428	0	238	17	0
9	Y	432	0	237	11	0
10	U	430	0	237	12	0
11	V	430	0	237	11	0
All	All	4299	0	2369	129	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 20.

The worst 5 of 129 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
11:V:109:DC:H2'	11:V:110:DG:C8	2.25	0.72
7:M:119:DC:H2''	7:M:120:DG:C5	2.27	0.70
5:F:112:DA:H2'	5:F:113:DG:H8	1.57	0.69
10:U:113:DT:H2''	10:U:114:DC:C5	2.28	0.69
6:C:112:DC:H2'	6:C:113:DA:C8	2.27	0.69

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

There are no protein molecules in this entry.

5.3.2 Protein sidechains [i](#)

There are no protein molecules in this entry.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS failed to run properly - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS failed to run properly - this section is therefore empty.

6.3 Carbohydrates

EDS failed to run properly - this section is therefore empty.

6.4 Ligands

EDS failed to run properly - this section is therefore empty.

6.5 Other polymers

EDS failed to run properly - this section is therefore empty.