



wwPDB NMR Structure Validation Summary Report ⓘ

Mar 8, 2026 – 04:42 AM UTC

PDB ID : 1U5S / pdb_00001u5s
Title : NMR structure of the complex between Nck-2 SH3 domain and PINCH-1 LIM4 domain
Authors : Vaynberg, J.; Fukuda, T.; Vinogradova, O.; Velyvis, A.; Ng, L.; Wu, C.; Qin, J.
Deposited on : 2004-07-28

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
wwPDB-RCI	:	v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV	:	Wang et al. (2010)
wwPDB-ShiftChecker	:	v1.2
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment was not calculated.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	229148	14424
Ramachandran outliers	224038	12848
Sidechain outliers	223484	12823

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Mol	Chain	Length	Quality of chain
1	A	71	35% 45% 18% .
2	B	66	32% 58% 9% .

2 Ensemble composition and analysis

This entry contains 18 models. Model 3 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *closest to the average*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:1-A:71 (71)	0.00	3
2	B:72-B:137 (66)	0.00	7

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 3 clusters and 1 single-model cluster was found.

Cluster number	Models
1	2, 3, 7, 9, 10, 12, 13, 16
2	4, 6, 8, 11, 15, 18
3	5, 14, 17
Single-model clusters	1

3 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 2150 atoms, of which 1057 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Cytoplasmic protein NCK2.

Mol	Chain	Residues	Atoms						Trace
1	A	71	Total	C	H	N	O	S	0
			1121	361	555	95	108	2	

- Molecule 2 is a protein called PINCH protein.

Mol	Chain	Residues	Atoms						Trace
2	B	66	Total	C	H	N	O	S	0
			1027	331	502	100	87	7	

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	72	GLY	-	cloning artifact	UNP P48059
B	73	SER	-	cloning artifact	UNP P48059

- Molecule 3 is ZINC ION (CCD ID: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms	
3	B	2	Total	Zn
			2	2

4 Residue-property plots


4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: Cytoplasmic protein NCK2




- Molecule 2: PINCH protein



4.2 Residue scores for the representative (medoid) model from the NMR ensemble

The representative model is number 3. Colouring as in section 4.1 above.

- Molecule 1: Cytoplasmic protein NCK2



- Molecule 2: PINCH protein



D136	G72	R84	R89	R98	H98	V100	E101	F102	G114	R116	H117	E119	R120	G122	L123	A124	Y125	C126	E127	T128	H129	Y130	M131	Q132	L133	F134	C135	V137
	S73	P85	V90		H99	E100	E102		G113	R115	H118	E118	R120	G122	L123	A124	Y125	C126	E127	T128	H129	Y130	M131	Q132	L133	F134	C135	V137
	M74	I86	V91			E101	F102		G114	R116	H117	E119	R120	G122	L123	A124	Y125	C126	E127	T128	H129	Y130	M131	Q132	L133	F134	C135	V137
	G75	P85	V91			E101	F102		G114	R116	H117	E119	R120	G122	L123	A124	Y125	C126	E127	T128	H129	Y130	M131	Q132	L133	F134	C135	V137
	P77	I86	V91			E101	F102		G114	R116	H117	E119	R120	G122	L123	A124	Y125	C126	E127	T128	H129	Y130	M131	Q132	L133	F134	C135	V137
	I78	I86	V91			E101	F102		G114	R116	H117	E119	R120	G122	L123	A124	Y125	C126	E127	T128	H129	Y130	M131	Q132	L133	F134	C135	V137
	C79	I86	V91			E101	F102		G114	R116	H117	E119	R120	G122	L123	A124	Y125	C126	E127	T128	H129	Y130	M131	Q132	L133	F134	C135	V137
	G80	I86	V91			E101	F102		G114	R116	H117	E119	R120	G122	L123	A124	Y125	C126	E127	T128	H129	Y130	M131	Q132	L133	F134	C135	V137
	A81	I86	V91			E101	F102		G114	R116	H117	E119	R120	G122	L123	A124	Y125	C126	E127	T128	H129	Y130	M131	Q132	L133	F134	C135	V137

5 Refinement protocol and experimental data overview

The models were refined using the following method: *simulated annealing, molecular dynamics, rigid body/torsion angle dynamics*.

Of the 50 calculated structures, 18 were deposited, based on the following criterion: *structures with the most representative side chains orientations on the complex interface*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
X-PLOR	structure solution	3.1
Xplor-NIH	structure solution	
Xplor-NIH	refinement	

No chemical shift data was provided.

6 Model quality i

6.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section:
ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	1.51±0.00	0±0/580 (0.0± 0.0%)	1.30±0.00	1±0/789 (0.1± 0.0%)
2	B	3.04±0.00	3±0/540 (0.6± 0.0%)	2.64±0.00	7±0/725 (1.0± 0.0%)
All	All	2.37	54/20160 (0.3%)	2.05	144/27252 (0.5%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
2	B	0.0±0.0	1.0±0.0
All	All	0	18

All unique bond outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)	Models	
								Worst	Total
2	B	112	PHE	CA-CB	41.59	2.15	1.53	5	18
2	B	137	VAL	C-O	-37.91	0.47	1.23	9	18
2	B	112	PHE	CB-CG	26.98	2.12	1.50	10	18

5 of 8 unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
2	B	112	PHE	CA-CB-CG	-42.76	71.04	113.80	5	18
2	B	137	VAL	CA-C-O	-29.42	70.79	120.80	6	18
2	B	112	PHE	N-CA-CB	21.45	135.02	110.35	3	18
2	B	112	PHE	CB-CG-CD2	-16.48	92.69	120.70	7	18

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
2	B	112	PHE	CB-CG-CD1	14.67	145.63	120.70	12	18

There are no chirality outliers.

All unique planar outliers are listed below.

Mol	Chain	Res	Type	Group	Models (Total)
2	B	112	PHE	Sidechain	18

6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	566	555	555	52±3
2	B	525	502	501	57±2
All	All	19674	19026	19008	1960

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 51.

5 of 164 unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
2:B:112:PHE:CB	2:B:112:PHE:CG	1.33	2.12	16	18
2:B:112:PHE:CB	2:B:112:PHE:CA	1.25	2.15	2	18
2:B:112:PHE:CG	2:B:112:PHE:CA	0.97	2.48	7	18
1:A:19:THR:HG22	1:A:20:GLU:OE1	0.91	1.65	6	2
2:B:112:PHE:CB	2:B:112:PHE:CD2	0.86	2.59	7	18

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation

was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	69/71 (97%)	50±0 (73±1%)	13±0 (18±1%)	6±0 (9±0%)	1	12
2	B	64/66 (97%)	49±0 (77±0%)	14±0 (22±0%)	1±0 (2±0%)	10	55
All	All	2394/2466 (97%)	1789 (75%)	479 (20%)	126 (5%)	2	22

5 of 7 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	33	VAL	18
1	A	39	ASN	18
1	A	40	ASP	18
1	A	41	PRO	18
1	A	42	GLU	18

6.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	64/64 (100%)	51±1 (80±1%)	13±1 (20±1%)	3	33
2	B	54/54 (100%)	47±0 (87±0%)	7±0 (13±0%)	6	46
All	All	2124/2124 (100%)	1769 (83%)	355 (17%)	4	39

5 of 22 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	17	SER	18
1	A	22	GLU	18
1	A	23	LEU	18
1	A	27	LYS	18
1	A	29	GLU	18

6.3.3 RNA ⓘ

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation

No chemical shift data were provided