



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 5, 2026 – 08:42 AM UTC

PDB ID : 4U5C / pdb_00004u5c
Title : Crystal structure of GluA2, con-ikot-ikot snail toxin, partial agonist FW and positive modulator (R,R)-2b complex
Authors : Chen, L.; Gouaux, E.
Deposited on : 2014-07-25
Resolution : 3.69 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.69 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1263 (3.80-3.56)
Clashscore	190562	1305 (3.80-3.56)
Ramachandran outliers	187476	1259 (3.80-3.56)
Sidechain outliers	187428	1256 (3.80-3.56)
RSRZ outliers	180081	1262 (3.80-3.56)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	814	69% 22% • 8%
1	B	814	68% 22% • 8%
1	C	814	69% 21% • 8%
1	D	814	67% 23% • 7%
2	E	90	69% 23% • 6%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	F	90	86% 8% • 6%

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 24451 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Glutamate receptor 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	747	Total	C	N	O	S	0	0	0
			5766	3715	941	1084	26			
1	B	746	Total	C	N	O	S	0	0	0
			5706	3665	940	1075	26			
1	C	745	Total	C	N	O	S	0	0	0
			5702	3663	933	1080	26			
1	D	753	Total	C	N	O	S	0	0	0
			5815	3737	961	1091	26			

There are 120 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	184	GLY	LYS	engineered mutation	UNP P19491
A	237	GLU	ASN	engineered mutation	UNP P19491
A	?	-	LEU	deletion	UNP P19491
A	?	-	PRO	deletion	UNP P19491
A	385	ASP	ASN	engineered mutation	UNP P19491
A	392	GLN	ASN	engineered mutation	UNP P19491
A	461	ASP	ASN	engineered mutation	UNP P19491
A	528	ALA	CYS	engineered mutation	UNP P19491
A	535	LEU	GLY	engineered mutation	UNP P19491
A	?	-	ARG	deletion	UNP P19491
A	?	-	GLU	deletion	UNP P19491
A	?	-	THR	deletion	UNP P19491
A	?	-	GLN	deletion	UNP P19491
A	?	-	SER	deletion	UNP P19491
A	565	GLU	SER	engineered mutation	UNP P19491
A	577	PHE	LEU	engineered mutation	UNP P19491
A	580	ALA	SER	engineered mutation	UNP P19491
A	582	LYS	GLY	engineered mutation	UNP P19491
A	583	LEU	ALA	engineered mutation	UNP P19491
A	585	PHE	MET	engineered mutation	UNP P19491
A	589	ALA	CYS	engineered mutation	UNP P19491

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	598	ALA	GLY	engineered mutation	UNP P19491
A	602	ALA	GLY	engineered mutation	UNP P19491
A	815	ALA	CYS	engineered mutation	UNP P19491
A	818	ARG	SER	engineered mutation	UNP P19491
A	819	MET	ARG	engineered mutation	UNP P19491
A	820	LYS	ALA	engineered mutation	UNP P19491
A	821	LEU	GLU	engineered mutation	UNP P19491
A	822	VAL	ALA	engineered mutation	UNP P19491
A	823	PRO	LYS	engineered mutation	UNP P19491
B	184	GLY	LYS	engineered mutation	UNP P19491
B	237	GLU	ASN	engineered mutation	UNP P19491
B	?	-	LEU	deletion	UNP P19491
B	?	-	PRO	deletion	UNP P19491
B	385	ASP	ASN	engineered mutation	UNP P19491
B	392	GLN	ASN	engineered mutation	UNP P19491
B	461	ASP	ASN	engineered mutation	UNP P19491
B	528	ALA	CYS	engineered mutation	UNP P19491
B	535	LEU	GLY	engineered mutation	UNP P19491
B	?	-	ARG	deletion	UNP P19491
B	?	-	GLU	deletion	UNP P19491
B	?	-	THR	deletion	UNP P19491
B	?	-	GLN	deletion	UNP P19491
B	?	-	SER	deletion	UNP P19491
B	565	GLU	SER	engineered mutation	UNP P19491
B	577	PHE	LEU	engineered mutation	UNP P19491
B	580	ALA	SER	engineered mutation	UNP P19491
B	582	LYS	GLY	engineered mutation	UNP P19491
B	583	LEU	ALA	engineered mutation	UNP P19491
B	585	PHE	MET	engineered mutation	UNP P19491
B	589	ALA	CYS	engineered mutation	UNP P19491
B	598	ALA	GLY	engineered mutation	UNP P19491
B	602	ALA	GLY	engineered mutation	UNP P19491
B	815	ALA	CYS	engineered mutation	UNP P19491
B	818	ARG	SER	engineered mutation	UNP P19491
B	819	MET	ARG	engineered mutation	UNP P19491
B	820	LYS	ALA	engineered mutation	UNP P19491
B	821	LEU	GLU	engineered mutation	UNP P19491
B	822	VAL	ALA	engineered mutation	UNP P19491
B	823	PRO	LYS	engineered mutation	UNP P19491
C	184	GLY	LYS	engineered mutation	UNP P19491
C	237	GLU	ASN	engineered mutation	UNP P19491
C	?	-	LEU	deletion	UNP P19491

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	?	-	PRO	deletion	UNP P19491
C	385	ASP	ASN	engineered mutation	UNP P19491
C	392	GLN	ASN	engineered mutation	UNP P19491
C	461	ASP	ASN	engineered mutation	UNP P19491
C	528	ALA	CYS	engineered mutation	UNP P19491
C	535	LEU	GLY	engineered mutation	UNP P19491
C	?	-	ARG	deletion	UNP P19491
C	?	-	GLU	deletion	UNP P19491
C	?	-	THR	deletion	UNP P19491
C	?	-	GLN	deletion	UNP P19491
C	?	-	SER	deletion	UNP P19491
C	565	GLU	SER	engineered mutation	UNP P19491
C	577	PHE	LEU	engineered mutation	UNP P19491
C	580	ALA	SER	engineered mutation	UNP P19491
C	582	LYS	GLY	engineered mutation	UNP P19491
C	583	LEU	ALA	engineered mutation	UNP P19491
C	585	PHE	MET	engineered mutation	UNP P19491
C	589	ALA	CYS	engineered mutation	UNP P19491
C	598	ALA	GLY	engineered mutation	UNP P19491
C	602	ALA	GLY	engineered mutation	UNP P19491
C	815	ALA	CYS	engineered mutation	UNP P19491
C	818	ARG	SER	engineered mutation	UNP P19491
C	819	MET	ARG	engineered mutation	UNP P19491
C	820	LYS	ALA	engineered mutation	UNP P19491
C	821	LEU	GLU	engineered mutation	UNP P19491
C	822	VAL	ALA	engineered mutation	UNP P19491
C	823	PRO	LYS	engineered mutation	UNP P19491
D	184	GLY	LYS	engineered mutation	UNP P19491
D	237	GLU	ASN	engineered mutation	UNP P19491
D	?	-	LEU	deletion	UNP P19491
D	?	-	PRO	deletion	UNP P19491
D	385	ASP	ASN	engineered mutation	UNP P19491
D	392	GLN	ASN	engineered mutation	UNP P19491
D	461	ASP	ASN	engineered mutation	UNP P19491
D	528	ALA	CYS	engineered mutation	UNP P19491
D	535	LEU	GLY	engineered mutation	UNP P19491
D	?	-	ARG	deletion	UNP P19491
D	?	-	GLU	deletion	UNP P19491
D	?	-	THR	deletion	UNP P19491
D	?	-	GLN	deletion	UNP P19491
D	?	-	SER	deletion	UNP P19491
D	565	GLU	SER	engineered mutation	UNP P19491

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	577	PHE	LEU	engineered mutation	UNP P19491
D	580	ALA	SER	engineered mutation	UNP P19491
D	582	LYS	GLY	engineered mutation	UNP P19491
D	583	LEU	ALA	engineered mutation	UNP P19491
D	585	PHE	MET	engineered mutation	UNP P19491
D	589	ALA	CYS	engineered mutation	UNP P19491
D	598	ALA	GLY	engineered mutation	UNP P19491
D	602	ALA	GLY	engineered mutation	UNP P19491
D	815	ALA	CYS	engineered mutation	UNP P19491
D	818	ARG	SER	engineered mutation	UNP P19491
D	819	MET	ARG	engineered mutation	UNP P19491
D	820	LYS	ALA	engineered mutation	UNP P19491
D	821	LEU	GLU	engineered mutation	UNP P19491
D	822	VAL	ALA	engineered mutation	UNP P19491
D	823	PRO	LYS	engineered mutation	UNP P19491


- Molecule 2 is a protein called Con-ikot-ikot.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	E	85	Total	C	N	O	S	0	0	0
			641	387	113	125	16			
2	F	85	Total	C	N	O	S	0	0	0
			641	387	113	125	16			

There are 8 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
E	-3	GLY	-	expression tag	UNP P0CB20
E	-2	PRO	-	expression tag	UNP P0CB20
E	-1	GLY	-	expression tag	UNP P0CB20
E	0	SER	-	expression tag	UNP P0CB20
F	-3	GLY	-	expression tag	UNP P0CB20
F	-2	PRO	-	expression tag	UNP P0CB20
F	-1	GLY	-	expression tag	UNP P0CB20
F	0	SER	-	expression tag	UNP P0CB20

- Molecule 3 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: C₈H₁₅NO₆).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	B	1	Total	C	N	O	0	0
			14	8	1	5		
3	C	1	Total	C	N	O	0	0
			14	8	1	5		
3	D	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 4 is 2-AMINO-3-(5-FLUORO-2,4-DIOXO-3,4-DIHYDRO-2H-PYRIMIDIN-1-YL)-PROPIONIC ACID (CCD ID: FWD) (formula: $C_7H_8FN_3O_4$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	F	N	O	0	0
			15	7	1	3	4		
4	B	1	Total	C	F	N	O	0	0
			15	7	1	3	4		
4	C	1	Total	C	F	N	O	0	0
			15	7	1	3	4		
4	D	1	Total	C	F	N	O	0	0
			15	7	1	3	4		

- Molecule 5 is N,N'-[biphenyl-4,4'-diyl]di(2R)propane-2,1-diyl]dipropane-2-sulfonamide (CCD ID: FWF) (formula: $C_{24}H_{36}N_2O_4S_2$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	B	1	Total	C	N	O	S	0	0
			32	24	2	4	2		
5	D	1	Total	C	N	O	S	0	0
			32	24	2	4	2		


3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Glutamate receptor 2




• Molecule 1: Glutamate receptor 2





• Molecule 1: Glutamate receptor 2

Chain C: 69% 21% 8%



• Molecule 1: Glutamate receptor 2

Chain D: 67% 23% 7%





4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	163.07Å 364.78Å 109.75Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.98 – 3.69 19.98 – 3.69	Depositor EDS
% Data completeness (in resolution range)	98.1 (19.98-3.69) 97.3 (19.98-3.69)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.70 (at 3.67Å)	Xtriage
Refinement program	PHENIX	Depositor
R, R_{free}	0.207 , 0.254 0.215 , 0.258	Depositor DCC
R_{free} test set	3554 reflections (4.97%)	wwPDB-VP
Wilson B-factor (Å ²)	116.9	Xtriage
Anisotropy	0.020	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.24 , 103.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.44$, $\langle L^2 \rangle = 0.27$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	24451	wwPDB-VP
Average B, all atoms (Å ²)	139.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.03% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FWD, NAG, FWF

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.32	0/5884	0.79	4/7970 (0.1%)
1	B	0.33	0/5823	0.82	11/7900 (0.1%)
1	C	0.32	0/5819	0.78	6/7897 (0.1%)
1	D	0.33	0/5933	0.82	9/8037 (0.1%)
2	E	0.34	0/651	0.90	2/873 (0.2%)
2	F	0.32	0/651	0.88	2/873 (0.2%)
All	All	0.33	0/24761	0.81	34/33550 (0.1%)

There are no bond length outliers.

The worst 5 of 34 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	317	VAL	CA-C-N	8.78	129.45	120.14
1	D	317	VAL	C-N-CA	8.78	129.45	120.14
1	D	94	THR	CA-C-N	8.75	128.97	119.87
1	D	94	THR	C-N-CA	8.75	128.97	119.87
1	A	94	THR	CA-C-N	8.69	128.34	119.56

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5766	0	5686	112	0
1	B	5706	0	5560	117	0
1	C	5702	0	5556	120	0
1	D	5815	0	5737	129	0
2	E	641	0	593	15	0
2	F	641	0	593	4	0
3	A	14	0	13	0	0
3	B	14	0	13	0	0
3	C	14	0	13	1	0
3	D	14	0	13	1	0
4	A	15	0	7	5	0
4	B	15	0	7	2	0
4	C	15	0	7	1	0
4	D	15	0	7	0	0
5	B	32	0	36	0	0
5	D	32	0	36	3	0
All	All	24451	0	23877	461	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

The worst 5 of 461 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:159:ILE:HG21	1:D:176:LEU:HD11	1.63	0.79
1:C:633:ILE:HG21	1:C:639:LEU:HD12	1.66	0.76
1:C:657:GLU:HG2	1:C:660:ARG:HH21	1.50	0.76
1:C:98:PRO:HD3	1:C:110:ARG:HD2	1.71	0.73
1:B:476:ILE:HD13	1:B:734:ILE:HD12	1.70	0.72

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	737/814 (90%)	714 (97%)	23 (3%)	0	100	100
1	B	738/814 (91%)	709 (96%)	29 (4%)	0	100	100
1	C	737/814 (90%)	710 (96%)	27 (4%)	0	100	100
1	D	745/814 (92%)	714 (96%)	31 (4%)	0	100	100
2	E	83/90 (92%)	79 (95%)	4 (5%)	0	100	100
2	F	83/90 (92%)	78 (94%)	5 (6%)	0	100	100
All	All	3123/3436 (91%)	3004 (96%)	119 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	607/694 (88%)	578 (95%)	29 (5%)	23	48
1	B	592/694 (85%)	567 (96%)	25 (4%)	26	50
1	C	595/694 (86%)	575 (97%)	20 (3%)	32	54
1	D	612/694 (88%)	583 (95%)	29 (5%)	23	48
2	E	73/76 (96%)	71 (97%)	2 (3%)	39	58
2	F	73/76 (96%)	71 (97%)	2 (3%)	39	58
All	All	2552/2928 (87%)	2445 (96%)	107 (4%)	26	50

5 of 107 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	157	THR
1	C	751	LEU
1	D	612	ILE
1	C	195	ASP
1	C	400	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 24 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	242	GLN
1	C	411	ASN
1	C	340	ASN
1	C	709	ASN
1	B	333	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

10 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
4	FWD	D	902	-	14,15,15	1.99	3 (21%)	15,21,21	3.85	7 (46%)
5	FWF	D	903	-	31,33,33	0.40	0	40,48,48	2.04	5 (12%)
3	NAG	A	901	1	14,14,15	0.64	1 (7%)	17,19,21	0.76	0
3	NAG	D	901	1	14,14,15	1.02	1 (7%)	17,19,21	0.51	0
4	FWD	C	902	-	14,15,15	1.97	3 (21%)	15,21,21	3.92	7 (46%)
3	NAG	C	901	1	14,14,15	0.55	0	17,19,21	0.94	1 (5%)
4	FWD	B	902	-	14,15,15	1.97	3 (21%)	15,21,21	3.86	7 (46%)
4	FWD	A	902	-	14,15,15	1.99	3 (21%)	15,21,21	3.91	7 (46%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	NAG	B	901	1	14,14,15	0.29	0	17,19,21	0.44	0
5	FWF	B	903	-	31,33,33	0.43	0	40,48,48	1.81	5 (12%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	FWD	D	902	-	-	0/8/8/8	0/1/1/1
5	FWF	D	903	-	-	0/32/36/36	0/2/2/2
3	NAG	A	901	1	-	2/6/23/26	0/1/1/1
3	NAG	D	901	1	-	0/6/23/26	0/1/1/1
4	FWD	C	902	-	-	3/8/8/8	0/1/1/1
3	NAG	C	901	1	-	2/6/23/26	0/1/1/1
4	FWD	B	902	-	-	0/8/8/8	0/1/1/1
4	FWD	A	902	-	-	4/8/8/8	0/1/1/1
3	NAG	B	901	1	-	2/6/23/26	0/1/1/1
5	FWF	B	903	-	-	5/32/36/36	0/2/2/2

The worst 5 of 14 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	D	902	FWD	C6-C5	4.35	1.39	1.33
4	C	902	FWD	C6-C5	4.29	1.39	1.33
4	B	902	FWD	C6-C5	4.22	1.39	1.33
4	A	902	FWD	C6-C5	4.18	1.39	1.33
4	A	902	FWD	C4-C5	3.48	1.48	1.44

The worst 5 of 39 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	D	902	FWD	N3-C2-N1	7.62	121.40	114.86
4	C	902	FWD	N3-C2-N1	7.55	121.34	114.86
4	B	902	FWD	N3-C2-N1	7.46	121.27	114.86
4	A	902	FWD	N3-C2-N1	7.30	121.12	114.86
5	D	903	FWF	O2-S1-O1	-7.17	108.42	118.88

There are no chirality outliers.

5 of 18 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	A	902	FWD	C7-C8-C9-O91
4	A	902	FWD	C7-C8-C9-O92
4	C	902	FWD	C7-C8-C9-O91
5	B	903	FWF	C23-C20-S2-O4
5	B	903	FWF	C24-C20-S2-O4


There are no ring outliers.

6 monomers are involved in 13 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	D	903	FWF	3	0
3	D	901	NAG	1	0
4	C	902	FWD	1	0
3	C	901	NAG	1	0
4	B	902	FWD	2	0
4	A	902	FWD	5	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	747/814 (91%)	-0.30	8 (1%) 78 53	52, 121, 243, 357	0
1	B	746/814 (91%)	-0.28	1 (0%) 92 87	66, 123, 255, 349	0
1	C	745/814 (91%)	-0.24	1 (0%) 92 87	72, 148, 258, 374	0
1	D	753/814 (92%)	-0.35	3 (0%) 88 72	53, 122, 245, 354	0
2	E	85/90 (94%)	-0.43	0 100 100	59, 90, 127, 184	0
2	F	85/90 (94%)	-0.32	0 100 100	82, 113, 166, 227	0
All	All	3161/3436 (91%)	-0.30	13 (0%) 88 72	52, 126, 249, 374	0

The worst 5 of 13 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	792	VAL	6.1
1	A	598	ALA	5.6
1	D	601	VAL	4.3
1	A	806	ALA	3.8
1	D	602	ALA	3.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	NAG	D	901	14/15	0.58	0.19	137,181,523,523	0
3	NAG	B	901	14/15	0.75	0.12	117,222,245,247	0
3	NAG	C	901	14/15	0.80	0.17	120,137,162,180	0
4	FWD	A	902	15/15	0.89	0.10	69,102,169,175	0
4	FWD	D	902	15/15	0.89	0.10	40,76,109,148	0
4	FWD	B	902	15/15	0.92	0.09	84,107,141,176	0
4	FWD	C	902	15/15	0.93	0.11	74,106,162,180	0
3	NAG	A	901	14/15	0.94	0.12	77,91,112,129	0
5	FWF	B	903	32/32	0.95	0.11	51,110,130,159	0
5	FWF	D	903	32/32	0.96	0.10	11,61,114,151	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around FWF B 903:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around FWF D 903:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.