



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 11:33 AM UTC

PDB ID : 4U5Q / pdb_00004u5q
Title : High resolution crystal structure of reductase (R) domain of nonribosomal peptide synthetase from Mycobacterium tuberculosis
Authors : Patel, K.D.; Haque, A.S.; Priyadarshan, K.; Sankaranarayanan, R.
Deposited on : 2014-07-25
Resolution : 1.81 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.81 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1112 (1.82-1.82)
Clashscore	190562	1148 (1.82-1.82)
Ramachandran outliers	187476	1140 (1.82-1.82)
Sidechain outliers	187428	1140 (1.82-1.82)
RSRZ outliers	180081	1112 (1.82-1.82)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	478	8% 83% 7% 10%
1	B	478	10% 81% 8% 11%

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 7239 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Peptide synthetase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	429	Total	C	N	O	S	0	0	0
			3305	2092	576	625	12			
1	B	426	Total	C	N	O	S	0	0	0
			3293	2084	575	623	11			

There are 42 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	2035	MET	-	expression tag	UNP Q10896
A	2036	GLY	-	expression tag	UNP Q10896
A	2037	SER	-	expression tag	UNP Q10896
A	2038	SER	-	expression tag	UNP Q10896
A	2039	HIS	-	expression tag	UNP Q10896
A	2040	HIS	-	expression tag	UNP Q10896
A	2041	HIS	-	expression tag	UNP Q10896
A	2042	HIS	-	expression tag	UNP Q10896
A	2043	HIS	-	expression tag	UNP Q10896
A	2044	HIS	-	expression tag	UNP Q10896
A	2045	SER	-	expression tag	UNP Q10896
A	2046	SER	-	expression tag	UNP Q10896
A	2047	GLY	-	expression tag	UNP Q10896
A	2048	LEU	-	expression tag	UNP Q10896
A	2049	VAL	-	expression tag	UNP Q10896
A	2050	PRO	-	expression tag	UNP Q10896
A	2051	ARG	-	expression tag	UNP Q10896
A	2052	GLY	-	expression tag	UNP Q10896
A	2053	SER	-	expression tag	UNP Q10896
A	2054	HIS	-	expression tag	UNP Q10896
A	2055	MET	-	expression tag	UNP Q10896
B	2035	MET	-	expression tag	UNP Q10896
B	2036	GLY	-	expression tag	UNP Q10896
B	2037	SER	-	expression tag	UNP Q10896
B	2038	SER	-	expression tag	UNP Q10896

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	2039	HIS	-	expression tag	UNP Q10896
B	2040	HIS	-	expression tag	UNP Q10896
B	2041	HIS	-	expression tag	UNP Q10896
B	2042	HIS	-	expression tag	UNP Q10896
B	2043	HIS	-	expression tag	UNP Q10896
B	2044	HIS	-	expression tag	UNP Q10896
B	2045	SER	-	expression tag	UNP Q10896
B	2046	SER	-	expression tag	UNP Q10896
B	2047	GLY	-	expression tag	UNP Q10896
B	2048	LEU	-	expression tag	UNP Q10896
B	2049	VAL	-	expression tag	UNP Q10896
B	2050	PRO	-	expression tag	UNP Q10896
B	2051	ARG	-	expression tag	UNP Q10896
B	2052	GLY	-	expression tag	UNP Q10896
B	2053	SER	-	expression tag	UNP Q10896
B	2054	HIS	-	expression tag	UNP Q10896
B	2055	MET	-	expression tag	UNP Q10896

- Molecule 2 is 3,6,9,12,15,18,21,24,27-NONAOXANONACOSANE-1,29-DIOL (CCD ID: XPE) (formula: C₂₀H₄₂O₁₁).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	B	1	Total	C	O	0	0
			31	20	11		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	304	Total 304	O 304	0	0
3	B	306	Total 306	O 306	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Peptide synthetase



- Molecule 1: Peptide synthetase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	78.53Å 88.59Å 126.08Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	42.98 – 1.81 42.98 – 1.81	Depositor EDS
% Data completeness (in resolution range)	99.3 (42.98-1.81) 99.4 (42.98-1.81)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.09	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.18 (at 1.81Å)	Xtriage
Refinement program	PHENIX (1.10_2155)	Depositor
R, R_{free}	0.202 , 0.245 0.207 , 0.246	Depositor DCC
R_{free} test set	4030 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	23.9	Xtriage
Anisotropy	0.564	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 36.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	7239	wwPDB-VP
Average B, all atoms (Å ²)	28.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 47.13 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.0375e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: XPE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.31	0/3374	0.49	0/4593
1	B	0.30	0/3359	0.51	0/4571
All	All	0.30	0/6733	0.50	0/9164

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3305	0	3281	22	0
1	B	3293	0	3268	31	0
2	B	31	0	42	5	0
3	A	304	0	0	2	0
3	B	306	0	0	4	0
All	All	7239	0	6591	53	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (53) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:2151:ARG:NH2	3:A:2601:HOH:O	2.15	0.80
1:B:2187:ASP:OD1	1:B:2233:ARG:NH1	2.13	0.80
1:B:2191:ASP:OD2	3:B:2701:HOH:O	2.07	0.70
1:B:2255:ILE:HD11	1:B:2270:SER:HB2	1.72	0.69
1:B:2087:ARG:HE	2:B:2601:XPE:H142	1.57	0.68
1:B:2257:PRO:HB3	1:B:2467:SER:HB3	1.77	0.66
1:B:2187:ASP:OD2	1:B:2192:GLN:NE2	2.29	0.65
1:B:2142:ARG:NH1	3:B:2703:HOH:O	2.30	0.64
1:A:2427:GLN:HE21	1:A:2428:ARG:HG3	1.65	0.61
1:A:2411:GLU:HG2	2:B:2601:XPE:H51	1.84	0.60
1:B:2219:GLU:OE2	1:B:2224:ASN:ND2	2.33	0.60
1:B:2106:GLU:HG3	1:B:2108:ARG:NH1	2.17	0.59
1:A:2147:GLU:HG3	1:A:2151:ARG:HE	1.68	0.58
1:A:2427:GLN:HE21	1:A:2428:ARG:N	2.01	0.58
1:B:2087:ARG:HE	2:B:2601:XPE:H171	1.68	0.58
1:B:2121:ARG:HD2	1:B:2322:GLN:OE1	2.03	0.58
1:A:2256:GLU:OE2	1:A:2256:GLU:N	2.25	0.58
1:B:2106:GLU:OE1	3:B:2702:HOH:O	2.18	0.56
1:B:2464:ARG:HG2	1:B:2467:SER:N	2.21	0.55
1:B:2219:GLU:HG3	1:B:2224:ASN:HB2	1.87	0.55
1:B:2464:ARG:NH2	3:B:2706:HOH:O	2.39	0.54
1:A:2255:ILE:HD11	1:A:2270:SER:HB2	1.89	0.54
1:A:2427:GLN:NE2	1:A:2428:ARG:HG3	2.23	0.54
1:B:2217:TYR:HB3	1:B:2326:SER:HB2	1.91	0.52
1:B:2087:ARG:NE	2:B:2601:XPE:H142	2.26	0.50
1:B:2106:GLU:OE2	1:B:2106:GLU:N	2.34	0.50
1:A:2106:GLU:OE1	1:A:2108:ARG:NH2	2.46	0.49
1:A:2203:ASP:O	1:A:2242:PRO:HD2	2.16	0.46
1:A:2147:GLU:HG2	1:A:2151:ARG:HH21	1.80	0.46
1:B:2183:LYS:HG2	1:B:2227:GLY:HA3	1.98	0.46
1:B:2203:ASP:O	1:B:2242:PRO:HD2	2.16	0.46
1:B:2257:PRO:CB	1:B:2467:SER:HB3	2.46	0.46
1:A:2335:SER:HB3	1:A:2433:LEU:HD21	1.99	0.45
1:A:2117:GLY:HA2	1:A:2211:MET:HB2	1.98	0.45
1:B:2273:ARG:HD2	1:B:2288:TRP:CG	2.52	0.44
1:A:2217:TYR:HB3	1:A:2326:SER:HB2	1.99	0.44
1:B:2103:PRO:HB3	1:B:2240:LEU:O	2.18	0.44
1:B:2424:GLU:OE2	1:B:2427:GLN:NE2	2.51	0.43
1:A:2196:ARG:NH1	3:A:2620:HOH:O	2.50	0.43
1:B:2408:TRP:CE3	1:B:2496:ALA:HA	2.53	0.43
1:A:2142:ARG:HB3	1:A:2142:ARG:CZ	2.48	0.43
1:B:2087:ARG:NE	2:B:2601:XPE:H171	2.31	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:2211:MET:HE1	1:B:2326:SER:OG	2.18	0.43
1:A:2273:ARG:HD2	1:A:2288:TRP:CG	2.54	0.42
1:A:2328:TRP:CZ2	1:A:2332:MET:HG3	2.54	0.42
1:B:2177:GLU:OE1	1:B:2197:ARG:NH2	2.44	0.42
1:A:2132:ASP:HB3	1:A:2134:ASP:OD1	2.20	0.42
1:A:2241:LYS:O	1:A:2303:LEU:HD11	2.20	0.41
1:B:2106:GLU:HG3	1:B:2108:ARG:HH11	1.86	0.41
1:B:2279:TRP:CD2	1:B:2280:ALA:N	2.89	0.41
1:A:2103:PRO:HB3	1:A:2240:LEU:O	2.21	0.40
1:B:2140:LEU:HD11	1:B:2181:GLY:HA3	2.03	0.40
1:A:2343:PRO:HD3	1:A:2425:TRP:CE2	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	423/478 (88%)	415 (98%)	8 (2%)	0	100	100
1	B	416/478 (87%)	406 (98%)	10 (2%)	0	100	100
All	All	839/956 (88%)	821 (98%)	18 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	349/390 (90%)	349 (100%)	0	100	100
1	B	348/390 (89%)	348 (100%)	0	100	100
All	All	697/780 (89%)	697 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	2224	ASN
1	A	2427	GLN
1	A	2440	GLN
1	A	2458	GLN
1	B	2099	ASN
1	B	2192	GLN
1	B	2213	ASN
1	B	2298	ASN
1	B	2356	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	XPE	B	2601	-	30,30,30	0.71	0	29,29,29	0.41	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	XPE	B	2601	-	-	13/28/28/28	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (13) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	2601	XPE	C8-C9-O10-C11
2	B	2601	XPE	C11-C12-O13-C14
2	B	2601	XPE	O10-C11-C12-O13
2	B	2601	XPE	O7-C8-C9-O10
2	B	2601	XPE	O13-C14-C15-O16
2	B	2601	XPE	C9-C8-O7-C6
2	B	2601	XPE	C17-C18-O19-C20
2	B	2601	XPE	O16-C17-C18-O19
2	B	2601	XPE	C26-C27-O28-C29
2	B	2601	XPE	O28-C29-C30-O31
2	B	2601	XPE	O4-C5-C6-O7
2	B	2601	XPE	C2-C3-O4-C5
2	B	2601	XPE	C20-C21-O22-C23


There are no ring outliers.

1 monomer is involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	2601	XPE	5	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths,

bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	429/478 (89%)	0.51	36 (8%) 17 19	16, 25, 45, 61	0
1	B	426/478 (89%)	0.56	50 (11%) 9 10	16, 26, 47, 67	0
All	All	855/956 (89%)	0.53	86 (10%) 12 14	16, 25, 46, 67	0

All (86) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	2279	TRP	6.3
1	A	2075	PRO	5.4
1	B	2280	ALA	5.2
1	B	2457	LEU	5.0
1	B	2487	ASP	5.0
1	B	2211	MET	4.8
1	A	2220	LEU	4.8
1	B	2209	ALA	4.7
1	B	2258	SER	4.6
1	A	2211	MET	4.5
1	B	2488	ASN	4.5
1	B	2105	PRO	4.4
1	A	2466	CYS	4.2
1	A	2467	SER	4.1
1	B	2484	SER	4.1
1	B	2212	VAL	4.1
1	A	2279	TRP	3.9
1	A	2450	LEU	3.6
1	A	2215	PHE	3.6
1	B	2210	ALA	3.5
1	A	2256	GLU	3.4
1	A	2457	LEU	3.4
1	A	2209	ALA	3.3
1	A	2106	GLU	3.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	2257	PRO	3.3
1	B	2276	ASP	3.3
1	B	2215	PHE	3.3
1	A	2254	ALA	3.2
1	B	2463	THR	3.2
1	A	2421	ASP	3.2
1	A	2278	GLY	3.2
1	A	2217	TYR	3.1
1	A	2214	ALA	3.1
1	A	2212	VAL	3.0
1	B	2489	PRO	3.0
1	A	2484	SER	3.0
1	B	2104	SER	3.0
1	A	2258	SER	2.9
1	B	2379	VAL	2.9
1	A	2210	ALA	2.9
1	A	2276	ASP	2.9
1	B	2467	SER	2.9
1	B	2220	LEU	2.9
1	A	2216	PRO	2.9
1	B	2485	ASP	2.9
1	B	2450	LEU	2.9
1	B	2217	TYR	2.8
1	B	2214	ALA	2.8
1	B	2380	ALA	2.8
1	B	2076	THR	2.8
1	A	2219	GLU	2.7
1	A	2280	ALA	2.7
1	B	2423	ALA	2.7
1	A	2488	ASN	2.7
1	B	2464	ARG	2.6
1	B	2419	ILE	2.6
1	B	2259	ALA	2.6
1	B	2101	PRO	2.5
1	A	2427	GLN	2.5
1	B	2426	LEU	2.5
1	B	2275	VAL	2.5
1	A	2423	ALA	2.4
1	A	2277	GLY	2.4
1	A	2422	PHE	2.4
1	A	2509	LEU	2.3
1	A	2255	ILE	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	2216	PRO	2.3
1	B	2142	ARG	2.3
1	B	2182	ASP	2.3
1	B	2256	GLU	2.3
1	B	2272	THR	2.3
1	B	2449	LEU	2.3
1	B	2263	ASP	2.2
1	A	2142	ARG	2.2
1	A	2458	GLN	2.2
1	A	2485	ASP	2.1
1	B	2384	LEU	2.1
1	A	2163	GLU	2.1
1	B	2509	LEU	2.1
1	B	2281	GLY	2.1
1	B	2224	ASN	2.1
1	B	2427	GLN	2.1
1	B	2102	GLY	2.0
1	B	2254	ALA	2.0
1	B	2191	ASP	2.0
1	B	2213	ASN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.


6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	XPE	B	2601	31/31	0.81	0.14	32,39,44,49	0

The following is a graphical depiction of the model fit to experimental electron density of all

instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers ⓘ

There are no such residues in this entry.