



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2026 – 09:41 AM UTC

PDB ID : 5U5M / pdb_00005u5m
Title : CRYSTAL STRUCTURE OF I83E MEDITOPE-ENABLED
TRASTUZUMAB WITH AZIDO-MEDITOPE
Authors : Williams, J.C.; Bzymek, K.P.; Pucket, J.; Avery, K.A.; Ma, Y.; Xie, J.; Zer,
C.; Horne, D.
Deposited on : 2016-12-06
Resolution : 1.88 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.88 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1220 (1.88-1.88)
Clashscore	190562	1234 (1.88-1.88)
Ramachandran outliers	187476	1222 (1.88-1.88)
Sidechain outliers	187428	1222 (1.88-1.88)
RSRZ outliers	180081	1220 (1.88-1.88)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	214	87%13%
2	B	223	%93%7%
3	E	63	6% 94%6%
4	C	54	2% 85%15%
5	D	13	85%15%

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 5221 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called MEMAB TRASTUZUMAB, LIGHT CHAIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	214	Total	C	N	O	S	0	11	0
			1710	1068	287	347	8			

- Molecule 2 is a protein called MEMAB TRASTUZUMAB, HEAVY CHAIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	223	Total	C	N	O	S	0	8	0
			1709	1082	288	332	7			

- Molecule 3 is a protein called Protein L.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	E	63	Total	C	N	O	S	0	0	0
			479	303	77	98	1			

There are 7 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
E	19	SER	-	expression tag	UNP Q51918
E	20	GLU	-	expression tag	UNP Q51918
E	34	ILE	THR	engineered mutation	UNP Q51918
E	55	ALA	ASP	engineered mutation	UNP Q51918
E	73	ASN	TYR	engineered mutation	UNP Q51918
E	74	HIS	THR	engineered mutation	UNP Q51918
E	75	MET	ILE	engineered mutation	UNP Q51918

- Molecule 4 is a protein called Immunoglobulin G binding protein A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	C	54	Total	C	N	O	S	0	1	0
			430	261	75	93	1			


There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	1	GLY	-	expression tag	UNP Q2UW42
C	2	SER	-	expression tag	UNP Q2UW42
C	3	TYR	-	expression tag	UNP Q2UW42

- Molecule 5 is a protein called AZIDO-PEG4-MEDITOPE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
5	D	13	Total	C	N	O	S	0	1	0
			153	94	32	25	2			

- Molecule 6 is MESO-ERYTHRITOL (CCD ID: MRY) (formula: $C_4H_{10}O_4$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	B	1	Total	C	O	0	0
			8	4	4		

- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	299	Total	O	0	0
			299	299		
7	B	280	Total	O	0	0
			280	280		
7	E	57	Total	O	0	0
			57	57		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	C	75	Total 75	O 75	0	0
7	D	21	Total 21	O 21	0	0

3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: MEMAB TRASTUZUMAB, LIGHT CHAIN

Chain A: 




- Molecule 2: MEMAB TRASTUZUMAB, HEAVY CHAIN

Chain B: 



- Molecule 3: Protein L

Chain E: 



- Molecule 4: Immunoglobulin G binding protein A

Chain C: 



- Molecule 5: AZIDO-PEG4-MEDITOPE

Chain D: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	53.53Å 105.47Å 117.13Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	31.62 – 1.88 31.62 – 1.88	Depositor EDS
% Data completeness (in resolution range)	99.9 (31.62-1.88) 99.9 (31.62-1.88)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.12 (at 1.88Å)	Xtriage
Refinement program	PHENIX 1.9_1692	Depositor
R, R_{free}	0.161 , 0.197 0.163 , 0.198	Depositor DCC
R_{free} test set	2737 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	20.8	Xtriage
Anisotropy	0.312	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 48.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	5221	wwPDB-VP
Average B, all atoms (Å ²)	26.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.43% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 2GX, 56C, ACE, MRY

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.44	0/1773	0.72	0/2410
2	B	0.43	0/1775	0.71	0/2413
3	E	0.42	0/486	0.70	0/654
4	C	0.43	0/435	0.69	0/583
5	D	0.38	0/82	0.68	0/105
All	All	0.43	0/4551	0.71	0/6165

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1710	0	1671	22	0
2	B	1709	0	1695	12	0
3	E	479	0	454	3	0
4	C	430	0	408	6	0
5	D	153	0	80	1	0
6	B	8	0	10	1	0
7	A	299	0	0	6	1
7	B	280	0	0	4	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	C	75	0	0	1	0
7	D	21	0	0	0	0
7	E	57	0	0	0	1
All	All	5221	0	4318	41	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (41) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:E:68:LEU:HD12	3:E:72:GLY:HA2	1.52	0.92
1:A:21:ILE:HD11	1:A:73[A]:LEU:HD23	1.74	0.68
1:A:17:ASP:OD2	7:A:301:HOH:O	2.12	0.67
1:A:33:VAL:HG13	1:A:88[A]:CYS:SG	2.37	0.64
2:B:124:LYS:NZ	7:B:407:HOH:O	2.33	0.61
1:A:195:GLU:OE1	7:A:302:HOH:O	2.16	0.60
2:B:50:ARG:NH1	7:B:402:HOH:O	2.21	0.59
2:B:89:GLU:HB3	7:B:408:HOH:O	2.03	0.58
1:A:18[B]:ARG:HG3	1:A:76:SER:HA	1.86	0.58
1:A:88[A]:CYS:SG	1:A:89:GLN:N	2.79	0.56
1:A:142:ARG:NH2	7:A:307:HOH:O	2.37	0.56
2:B:12:VAL:HG11	2:B:86:LEU:HD13	1.88	0.55
1:A:39:ARG:NH2	7:A:308:HOH:O	2.38	0.54
2:B:83:MET:HE2	2:B:86:LEU:HD21	1.91	0.53
2:B:65[B]:LYS:HE2	4:C:41:ASN:OD1	2.09	0.52
1:A:4:MET:HE3	1:A:23:CYS:SG	2.50	0.51
2:B:130:PRO:HD3	2:B:216:LYS:HE2	1.93	0.50
1:A:134:CYS:SG	1:A:194[B]:CYS:SG	3.05	0.50
1:A:37:GLN:HB2	1:A:47:LEU:HD11	1.92	0.50
2:B:221:LYS:HG2	2:B:222:SER:H	1.77	0.50
2:B:87:ARG:NH2	7:B:405:HOH:O	2.27	0.49
1:A:123[B]:GLU:HA	1:A:126:LYS:HD2	1.95	0.48
1:A:81:GLU:OE2	7:A:303:HOH:O	2.21	0.46
4:C:3:TYR:HB3	4:C:7:GLN:HB2	1.98	0.46
4:C:43:LEU:HG	4:C:47:LYS:HE3	1.96	0.46
1:A:11:LEU:HD23	1:A:19[B]:VAL:HG13	1.99	0.45
4:C:14:ILE:HD13	4:C:28:PHE:HB3	1.99	0.45
1:A:9:ILE:HD12	5:D:1:CYS:SG	2.57	0.45
1:A:66:ARG:HG3	1:A:71:PHE:CE2	2.53	0.44
1:A:123[A]:GLU:HA	1:A:126:LYS:HD2	2.00	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:C:14:ILE:HD12	4:C:32:LEU:HD11	2.00	0.43
4:C:47:LYS:HD2	7:C:116:HOH:O	2.18	0.43
1:A:23:CYS:SG	1:A:88[A]:CYS:SG	3.16	0.43
1:A:123[B]:GLU:OE1	2:B:216:LYS:NZ	2.46	0.43
1:A:145:LYS:HE3	7:A:546:HOH:O	2.17	0.43
2:B:7:SER:OG	2:B:21[B]:SER:OG	2.38	0.42
3:E:68:LEU:HD13	3:E:75:MET:HG2	2.02	0.41
2:B:157:VAL:O	6:B:301:MRY:HAE	2.21	0.41
1:A:66:ARG:HG2	1:A:67:SER:N	2.35	0.40
1:A:175:LEU:C	1:A:175:LEU:HD23	2.46	0.40
3:E:66:ALA:HB1	3:E:75:MET:CE	2.51	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
7:A:542:HOH:O	7:E:119:HOH:O[4_555]	2.18	0.02

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	223/214 (104%)	219 (98%)	4 (2%)	0	100	100
2	B	229/223 (103%)	227 (99%)	2 (1%)	0	100	100
3	E	61/63 (97%)	61 (100%)	0	0	100	100
4	C	53/54 (98%)	53 (100%)	0	0	100	100
5	D	9/13 (69%)	9 (100%)	0	0	100	100
All	All	575/567 (101%)	569 (99%)	6 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	200/189 (106%)	200 (100%)	0	100	100
2	B	192/185 (104%)	191 (100%)	1 (0%)	81	76
3	E	46/47 (98%)	46 (100%)	0	100	100
4	C	48/47 (102%)	48 (100%)	0	100	100
5	D	10/10 (100%)	10 (100%)	0	100	100
All	All	496/478 (104%)	495 (100%)	1 (0%)	87	84

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	145	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	41	ASN
1	A	79	GLN
1	A	89	GLN
4	C	38	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

3 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The

Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	2GX	D	5	5	16,18,19	0.96	0	20,23,25	0.64	0
5	56C	D	8[A]	5	20,24,25	0.74	1 (5%)	19,25,27	1.03	2 (10%)
5	56C	D	8[B]	5	20,24,25	0.73	0	19,25,27	0.91	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	2GX	D	5	5	-	0/13/14/16	0/2/2/2
5	56C	D	8[A]	5	-	9/23/24/26	-
5	56C	D	8[B]	5	-	10/23/24/26	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	D	8[A]	56C	N04-N03	2.01	1.27	1.23

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	D	8[A]	56C	C01-C02-N01	2.37	117.47	111.82
5	D	8[B]	56C	C01-C02-N01	2.32	117.35	111.82
5	D	8[A]	56C	N01-CZ-NE	2.07	122.19	119.28

There are no chirality outliers.

All (19) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	D	8[A]	56C	O04-C07-C08-N02
5	D	8[B]	56C	O04-C07-C08-N02
5	D	8[B]	56C	O02-C05-C06-O04
5	D	8[A]	56C	O02-C05-C06-O04
5	D	8[B]	56C	O01-C03-C04-O02

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
5	D	8[A]	56C	CA-CB-CG-CD
5	D	8[A]	56C	O01-C01-C02-N01
5	D	8[B]	56C	CA-CB-CG-CD
5	D	8[B]	56C	C06-C05-O02-C04
5	D	8[A]	56C	O01-C03-C04-O02
5	D	8[B]	56C	C02-C01-O01-C03
5	D	8[B]	56C	C04-C03-O01-C01
5	D	8[A]	56C	C03-C04-O02-C05
5	D	8[A]	56C	C08-C07-O04-C06
5	D	8[B]	56C	CG-CD-NE-CZ
5	D	8[B]	56C	C08-C07-O04-C06
5	D	8[A]	56C	C05-C06-O04-C07
5	D	8[A]	56C	CG-CD-NE-CZ
5	D	8[B]	56C	O01-C01-C02-N01

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
6	MRY	B	301	-	7,7,7	1.77	2 (28%)	8,8,8	0.98	1 (12%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.

'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	MRY	B	301	-	-	2/8/8/8	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	B	301	MRY	OAD-CAC	-2.78	1.37	1.43
6	B	301	MRY	OAF-CAE	-2.56	1.38	1.43

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	B	301	MRY	CAA-CAC-CAE	-2.03	108.86	113.00

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	B	301	MRY	OAB-CAA-CAC-OAD
6	B	301	MRY	OAB-CAA-CAC-CAE

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	B	301	MRY	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	214/214 (100%)	-0.51	1 (0%) 87 90	9, 21, 41, 70	11 (5%)
2	B	223/223 (100%)	-0.52	2 (0%) 81 85	10, 21, 37, 98	8 (3%)
3	E	63/63 (100%)	0.14	4 (6%) 26 28	18, 31, 55, 72	0
4	C	54/54 (100%)	-0.34	1 (1%) 66 72	13, 21, 50, 54	1 (1%)
5	D	10/13 (76%)	-0.22	0 100 100	15, 21, 40, 41	0
All	All	564/567 (99%)	-0.42	8 (1%) 73 79	9, 21, 45, 98	20 (3%)

All (8) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	E	72	GLY	3.6
3	E	68	LEU	3.1
1	A	214	CYS	3.0
3	E	71	GLY	2.8
4	C	1	GLY	2.6
2	B	105	TYR	2.2
2	B	222	SER	2.2
3	E	70	ASP	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	56C	D	8[A]	25/26	0.85	0.15	13,29,46,47	25
5	56C	D	8[B]	25/26	0.85	0.15	14,26,48,53	25

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
5	2GX	D	5	17/18	0.97	0.05	14,16,19,22	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
6	MRY	B	301	8/8	0.90	0.09	12,19,30,32	8

6.5 Other polymers [i](#)

There are no such residues in this entry.