



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 8, 2026 – 10:07 AM UTC

PDB ID : 7U5Y / pdb_00007u5y
Title : Crystal structure of ribulose-phosphate 3-epimerase from *Pseudomonas aeruginosa*
Authors : Seattle Structural Genomics Center for Infectious Disease (SSGCID)
Deposited on : 2022-03-02
Resolution : 2.55 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.55 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1091 (2.54-2.54)
Clashscore	190562	1120 (2.54-2.54)
Ramachandran outliers	187476	1106 (2.54-2.54)
Sidechain outliers	187428	1106 (2.54-2.54)
RSRZ outliers	180081	1091 (2.54-2.54)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	232	2% 93% 5% .
1	B	232	2% 89% 6% .
1	C	232	2% 88% 7% 5%
1	D	232	2% 88% 8% .
1	E	232	2% 88% 8% .

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	232	2% 88% 7% 5%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 9982 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ribulose-phosphate 3-epimerase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	227	Total	C	N	O	S	0	0	0
			1698	1079	297	312	10			
1	B	222	Total	C	N	O	S	0	0	0
			1616	1032	276	298	10			
1	C	220	Total	C	N	O	S	0	0	0
			1639	1045	278	306	10			
1	D	223	Total	C	N	O	S	0	0	0
			1645	1049	278	308	10			
1	E	223	Total	C	N	O	S	0	0	0
			1623	1036	273	304	10			
1	F	221	Total	C	N	O	S	0	0	0
			1619	1033	274	302	10			

There are 48 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-7	MET	-	expression tag	UNP Q9I5T1
A	-6	ALA	-	expression tag	UNP Q9I5T1
A	-5	HIS	-	expression tag	UNP Q9I5T1
A	-4	HIS	-	expression tag	UNP Q9I5T1
A	-3	HIS	-	expression tag	UNP Q9I5T1
A	-2	HIS	-	expression tag	UNP Q9I5T1
A	-1	HIS	-	expression tag	UNP Q9I5T1
A	0	HIS	-	expression tag	UNP Q9I5T1
B	-7	MET	-	expression tag	UNP Q9I5T1
B	-6	ALA	-	expression tag	UNP Q9I5T1
B	-5	HIS	-	expression tag	UNP Q9I5T1
B	-4	HIS	-	expression tag	UNP Q9I5T1
B	-3	HIS	-	expression tag	UNP Q9I5T1
B	-2	HIS	-	expression tag	UNP Q9I5T1
B	-1	HIS	-	expression tag	UNP Q9I5T1
B	0	HIS	-	expression tag	UNP Q9I5T1
C	-7	MET	-	expression tag	UNP Q9I5T1

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	-6	ALA	-	expression tag	UNP Q9I5T1
C	-5	HIS	-	expression tag	UNP Q9I5T1
C	-4	HIS	-	expression tag	UNP Q9I5T1
C	-3	HIS	-	expression tag	UNP Q9I5T1
C	-2	HIS	-	expression tag	UNP Q9I5T1
C	-1	HIS	-	expression tag	UNP Q9I5T1
C	0	HIS	-	expression tag	UNP Q9I5T1
D	-7	MET	-	expression tag	UNP Q9I5T1
D	-6	ALA	-	expression tag	UNP Q9I5T1
D	-5	HIS	-	expression tag	UNP Q9I5T1
D	-4	HIS	-	expression tag	UNP Q9I5T1
D	-3	HIS	-	expression tag	UNP Q9I5T1
D	-2	HIS	-	expression tag	UNP Q9I5T1
D	-1	HIS	-	expression tag	UNP Q9I5T1
D	0	HIS	-	expression tag	UNP Q9I5T1
E	-7	MET	-	expression tag	UNP Q9I5T1
E	-6	ALA	-	expression tag	UNP Q9I5T1
E	-5	HIS	-	expression tag	UNP Q9I5T1
E	-4	HIS	-	expression tag	UNP Q9I5T1
E	-3	HIS	-	expression tag	UNP Q9I5T1
E	-2	HIS	-	expression tag	UNP Q9I5T1
E	-1	HIS	-	expression tag	UNP Q9I5T1
E	0	HIS	-	expression tag	UNP Q9I5T1
F	-7	MET	-	expression tag	UNP Q9I5T1
F	-6	ALA	-	expression tag	UNP Q9I5T1
F	-5	HIS	-	expression tag	UNP Q9I5T1
F	-4	HIS	-	expression tag	UNP Q9I5T1
F	-3	HIS	-	expression tag	UNP Q9I5T1
F	-2	HIS	-	expression tag	UNP Q9I5T1
F	-1	HIS	-	expression tag	UNP Q9I5T1
F	0	HIS	-	expression tag	UNP Q9I5T1

- Molecule 2 is ZINC ION (CCD ID: ZN) (formula: Zn) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total Zn 1 1	0	0
2	B	1	Total Zn 1 1	0	0
2	C	1	Total Zn 1 1	0	0

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	D	1	Total 1	Zn 1	0	0
2	E	1	Total 1	Zn 1	0	0
2	F	1	Total 1	Zn 1	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	40	Total 40	O 40	0	0
3	B	18	Total 18	O 18	0	0
3	C	25	Total 25	O 25	0	0
3	D	22	Total 22	O 22	0	0
3	E	16	Total 16	O 16	0	0
3	F	15	Total 15	O 15	0	0

- Molecule 1: Ribulose-phosphate 3-epimerase




-
- | Amino Acid | Count |
|------------|-------|
| MET | 10 |
| ALA | 10 |
| HIS | 10 |
| HIS | 10 |
| HIS | 10 |
| HIS | 10 |
| HIS | 10 |
| HO | 10 |
| F4 | 9 |
| A7 | 8 |
| P8 | 8 |
| F35 | 8 |
| M51 | 8 |
| V67 | 8 |
| T90 | 8 |
| R101 | 8 |
| G114 | 8 |
| Y128 | 8 |
| M139 | 8 |
| P143 | 8 |
| F144 | 8 |
| G145 | 8 |
| GLY | 8 |
| Q148 | 8 |
| I176 | 8 |
| I188 | 8 |
| V197 | 8 |
| A198 | 8 |
| H223 | 8 |
| CUN | 1 |

-
- | Category | Value (approx.) |
|----------|-----------------|
| MET | 10 |
| ALA | 15 |
| HIS | 10 |
| HIS | 10 |
| HIS | 10 |
| HIS | 10 |
| HIS | 10 |
| HIS | 10 |
| M1 | 10 |
| P8 | 15 |
| S9 | 15 |
| H34 | 15 |
| M51 | 15 |
| H68 | 15 |
| R76 | 15 |
| D80 | 15 |
| F81 | 15 |
| L82 | 15 |
| T90 | 15 |
| G109 | 15 |
| G114 | 15 |
| L115 | 15 |
| V116 | 15 |
| K127 | 15 |
| Y128 | 15 |
| S140 | 15 |
| G144 | 100 |
| F145 | 15 |
| G147 | 10 |
| G148 | 100 |
| T154 | 15 |
| A198 | 15 |
| A222 | 100 |
| HIS | 10 |
| GIN | 10 |

- MET
 ALA
 HIS
 HIS
 HIS
 HIS
 HIS
 H-1
 H-1
 M1
 P8
 S9
 I10
 L11
 R34
 F35
 D36
 T47
 M51
 V67
 M70
 T90
 G114
 Y128
 P143
 G144
 F145
 GLY
 GLY
 Q148
 E160
 A161
 R162
 A198
 F203
 H223
 GIN

- | |
|------|
| NET |
| ALA |
| HIS |
| HIS |
| HIS |
| HIS |
| HIS |
| H-1 |
| HO |
| M1 |
| P8 |
| L11 |
| F36 |
| D36 |
| T47 |
| M51 |
| V67 |
| I82 |
| T90 |
| G109 |
| G114 |
| L115 |
| L126 |
| M139 |
| P143 |
| G144 |
| F145 |
| GLY |
| GLY |
| Q148 |
| E160 |
| A161 |
| R162 |
| A198 |
| N204 |
| E210 |
| G221 |
| A222 |
| H233 |
| NEN |

- 
- WORLD WIDE
PDB
PROTEIN DATA BANK



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	99.01Å 110.13Å 126.76Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.03 – 2.55 48.03 – 2.55	Depositor EDS
% Data completeness (in resolution range)	99.7 (48.03-2.55) 99.7 (48.03-2.55)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.15 (at 2.54Å)	Xtriage
Refinement program	PHENIX 1.20.1	Depositor
R, R_{free}	0.203 , 0.258 0.203 , 0.258	Depositor DCC
R_{free} test set	1985 reflections (4.33%)	wwPDB-VP
Wilson B-factor (Å ²)	55.3	Xtriage
Anisotropy	0.605	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 48.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	9982	wwPDB-VP
Average B, all atoms (Å ²)	68.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.38% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.23	0/1734	0.43	0/2359
1	B	0.21	0/1647	0.38	0/2247
1	C	0.22	0/1670	0.41	0/2272
1	D	0.22	0/1677	0.41	0/2285
1	E	0.18	0/1655	0.38	0/2257
1	F	0.20	0/1651	0.38	0/2253
All	All	0.21	0/10034	0.40	0/13673

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1698	0	1658	9	0
1	B	1616	0	1572	10	0
1	C	1639	0	1623	11	0
1	D	1645	0	1606	13	0
1	E	1623	0	1565	11	0
1	F	1619	0	1572	10	0
2	A	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
2	E	1	0	0	0	0
2	F	1	0	0	0	0
3	A	40	0	0	1	0
3	B	18	0	0	1	0
3	C	25	0	0	1	0
3	D	22	0	0	1	0
3	E	16	0	0	0	0
3	F	15	0	0	0	0
All	All	9982	0	9596	56	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 56 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:160:GLU:OE1	1:D:128:TYR:OH	2.11	0.69
1:A:128:TYR:OH	1:F:160:GLU:OE1	2.13	0.66
1:B:51:MET:HE1	1:F:47:THR:HG22	1.82	0.62
1:A:51:MET:HE1	1:E:47:THR:HG22	1.85	0.58
1:D:90:THR:HG22	1:D:114:GLY:HA3	1.86	0.57

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	223/232 (96%)	214 (96%)	9 (4%)	0	100 100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	218/232 (94%)	209 (96%)	9 (4%)	0	100	100
1	C	216/232 (93%)	207 (96%)	9 (4%)	0	100	100
1	D	219/232 (94%)	211 (96%)	8 (4%)	0	100	100
1	E	219/232 (94%)	210 (96%)	9 (4%)	0	100	100
1	F	217/232 (94%)	209 (96%)	8 (4%)	0	100	100
All	All	1312/1392 (94%)	1260 (96%)	52 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	173/184 (94%)	172 (99%)	1 (1%)	78	87
1	B	161/184 (88%)	161 (100%)	0	100	100
1	C	169/184 (92%)	169 (100%)	0	100	100
1	D	167/184 (91%)	167 (100%)	0	100	100
1	E	161/184 (88%)	161 (100%)	0	100	100
1	F	163/184 (89%)	163 (100%)	0	100	100
All	All	994/1104 (90%)	993 (100%)	1 (0%)	88	94

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	186	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 10 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	216	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	D	223	HIS
1	E	97	GLN
1	C	216	HIS
1	C	221	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 6 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	227/232 (97%)	-0.15	2 (0%) 81 83	43, 56, 71, 110	0
1	B	222/232 (95%)	0.09	3 (1%) 73 74	45, 70, 94, 133	0
1	C	220/232 (94%)	-0.01	3 (1%) 73 74	45, 65, 95, 122	0
1	D	223/232 (96%)	0.01	4 (1%) 67 68	42, 64, 96, 132	0
1	E	223/232 (96%)	0.34	5 (2%) 62 63	46, 76, 119, 133	0
1	F	221/232 (95%)	0.14	4 (1%) 67 68	43, 72, 105, 120	0
All	All	1336/1392 (95%)	0.07	21 (1%) 70 71	42, 67, 101, 133	0

The worst 5 of 21 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	E	223	HIS	4.3
1	D	148	GLN	4.2
1	F	222	ALA	4.1
1	A	148	GLN	3.5
1	A	-4	HIS	3.3

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

6.4 Ligands




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	ZN	C	300	1/1	0.87	0.21	116,116,116,116	1
2	ZN	F	300	1/1	0.90	0.07	77,77,77,77	1
2	ZN	D	300	1/1	0.94	0.10	72,72,72,72	1
2	ZN	B	300	1/1	0.94	0.08	81,81,81,81	1
2	ZN	E	300	1/1	0.95	0.08	93,93,93,93	1
2	ZN	A	300	1/1	0.96	0.06	54,54,54,54	1

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around ZN C 300:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around ZN F 300:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around ZN D 300:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around ZN B 300:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





Electron density around ZN E 300:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around ZN A 300:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers ⓘ

There are no such residues in this entry.