



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 09:22 AM UTC

PDB ID : 8U6O / pdb_00008u6o
Title : Crystal Structure of HIV-1 Reverse Transcriptase in Complex with 5-(2-(3-oxo-3-(pyrrolidin-1-yl)propoxy)phenoxy)-2-naphthonitrile (JLJ753), a non-nucleoside inhibitor
Authors : Prucha, G.; Henry, S.; Jorgensen, W.L.; Anderson, K.S.
Deposited on : 2023-09-13
Resolution : 2.48 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.48 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7589 (2.50-2.46)
Clashscore	190562	8295 (2.50-2.46)
Ramachandran outliers	187476	8164 (2.50-2.46)
Sidechain outliers	187428	8166 (2.50-2.46)
RSRZ outliers	180081	7593 (2.50-2.46)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	556	7% 76% 22% ..
2	B	428	6% 71% 20% • 5%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 7538 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Reverse transcriptase/ribonuclease H.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	551	Total	C	N	O	S	0	0	0
			4251	2748	709	787	7			

There are 5 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	MET	-	expression tag	UNP P03366
A	0	VAL	-	expression tag	UNP P03366
A	172	ALA	LYS	engineered mutation	UNP P03366
A	173	ALA	LYS	engineered mutation	UNP P03366
A	280	SER	CYS	engineered mutation	UNP P03366


- Molecule 2 is a protein called p51 RT.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	406	Total	C	N	O	S	0	1	0
			3242	2111	536	589	6			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	280	SER	CYS	engineered mutation	UNP P03366

- Molecule 3 is 5-{2-[3-oxo-3-(pyrrolidin-1-yl)propoxy]phenoxy}naphthalene-2-carbonitrile (CCD ID: VWB) (formula: C₂₄H₂₂N₂O₃) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			29	24	2	3		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	7	Total	O	0	0
			7	7		
4	B	9	Total	O	0	0
			9	9		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Reverse transcriptase/ribonuclease H



• Molecule 2: p51 RT



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	224.19Å 68.03Å 103.80Å 90.00° 107.67° 90.00°	Depositor
Resolution (Å)	43.45 – 2.48 43.45 – 2.48	Depositor EDS
% Data completeness (in resolution range)	99.0 (43.45-2.48) 99.0 (43.45-2.48)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.25 (at 2.48Å)	Xtriage
Refinement program	PHENIX (1.20.1_4487: ???)	Depositor
R, R_{free}	0.226 , 0.269 0.230 , 0.270	Depositor DCC
R_{free} test set	2000 reflections (3.76%)	wwPDB-VP
Wilson B-factor (Å ²)	71.9	Xtriage
Anisotropy	0.061	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 53.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.35$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	7538	wwPDB-VP
Average B, all atoms (Å ²)	80.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.08% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: VWB

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.41	0/4365	0.61	1/5970 (0.0%)
2	B	0.46	2/3338 (0.1%)	0.62	2/4557 (0.0%)
All	All	0.43	2/7703 (0.0%)	0.61	3/10527 (0.0%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	354[A]	TYR	C-O	7.11	1.32	1.23
2	B	354[B]	TYR	C-O	7.11	1.32	1.23

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	354[A]	TYR	CA-C-O	6.32	128.30	121.16
2	B	354[B]	TYR	CA-C-O	6.32	128.30	121.16
1	A	3	SER	N-CA-C	5.09	122.29	107.84

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4251	0	4074	71	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	3242	0	3151	59	0
3	A	29	0	0	0	0
4	A	7	0	0	1	0
4	B	9	0	0	2	0
All	All	7538	0	7225	120	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (120) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:356:ARG:HG2	1:A:367:GLN:HE21	1.25	0.98
1:A:356:ARG:HG2	1:A:367:GLN:NE2	1.94	0.82
2:B:356:ARG:HH21	2:B:357:MET:HG3	1.41	0.82
1:A:406:TRP:CD2	2:B:420:PRO:HB3	2.16	0.80
2:B:114:ALA:HB2	2:B:214:LEU:HG	1.63	0.79
2:B:27:THR:HG22	2:B:30:LYS:H	1.47	0.79
2:B:356:ARG:HG3	2:B:362:THR:HG21	1.65	0.78
1:A:79:GLU:OE2	1:A:83:ARG:NH1	2.22	0.72
1:A:356:ARG:HD2	1:A:358:ARG:H	1.54	0.71
1:A:21:VAL:HB	1:A:59:PRO:HD3	1.77	0.67
1:A:343:GLN:HG3	1:A:349:LEU:HD11	1.76	0.66
1:A:84:THR:HG21	1:A:153:TRP:HE1	1.60	0.66
1:A:27:THR:HG23	1:A:30:LYS:HD2	1.81	0.62
2:B:254:VAL:HG13	2:B:283:LEU:HD22	1.83	0.60
1:A:406:TRP:CZ3	1:A:407:GLN:HG3	2.36	0.60
1:A:19:PRO:HB3	1:A:79:GLU:HG3	1.84	0.60
2:B:358:ARG:O	2:B:361:HIS:N	2.35	0.60
1:A:454:LYS:HG3	1:A:552:VAL:HG13	1.85	0.59
1:A:246:LEU:HD22	1:A:260:LEU:HD12	1.85	0.58
1:A:356:ARG:CG	1:A:367:GLN:HE21	2.09	0.58
2:B:319:TYR:OH	2:B:385:LYS:HD2	2.04	0.58
1:A:60:VAL:HG21	1:A:130:PHE:CD2	2.40	0.57
2:B:275:LYS:HD3	2:B:276:VAL:H	1.69	0.57
1:A:135:ILE:H	1:A:135:ILE:HD12	1.69	0.57
1:A:296:THR:O	1:A:300:GLU:HG2	2.05	0.56
1:A:405:TYR:CE2	1:A:407:GLN:HB2	2.40	0.56
1:A:266:TRP:O	1:A:269:GLN:HG2	2.06	0.55
2:B:320:ASP:OD1	2:B:322:SER:OG	2.25	0.55
2:B:282:LEU:HD11	2:B:299:ALA:HB2	1.89	0.55

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:356:ARG:HD2	1:A:358:ARG:HB2	1.90	0.54
1:A:363:ASN:HA	1:A:511:ASP:OD1	2.07	0.54
1:A:80:LEU:O	1:A:84:THR:HG23	2.08	0.54
2:B:65:LYS:HG3	2:B:72:ARG:HD2	1.90	0.54
2:B:354[B]:TYR:CE2	2:B:375:ILE:HG13	2.43	0.54
1:A:21:VAL:O	1:A:57:ASN:ND2	2.39	0.54
1:A:171:PHE:HB2	1:A:208:HIS:HD1	1.74	0.53
2:B:241:VAL:HG22	2:B:350:LYS:HA	1.91	0.53
2:B:336:GLN:HB3	2:B:353:LYS:HE2	1.91	0.52
1:A:406:TRP:CG	2:B:420:PRO:HB3	2.45	0.52
1:A:452:LEU:HD23	1:A:470:THR:HA	1.92	0.51
1:A:97:PRO:HG2	1:A:232:TYR:CD2	2.46	0.51
1:A:101:LYS:NZ	4:A:701:HOH:O	2.44	0.51
1:A:435:VAL:HG22	2:B:290:THR:HG21	1.93	0.51
2:B:183:TYR:CE2	2:B:184:MET:HG3	2.45	0.51
2:B:362:THR:HB	2:B:367:GLN:HE21	1.76	0.51
2:B:330:GLN:HG2	2:B:338:THR:OG1	2.12	0.50
2:B:58:THR:HG22	2:B:76:ASP:O	2.11	0.50
2:B:317:VAL:HG13	2:B:347:LYS:HB3	1.93	0.50
1:A:410:TRP:HD1	2:B:401:TRP:CE2	2.29	0.49
2:B:275:LYS:HG3	2:B:277:ARG:H	1.78	0.49
1:A:406:TRP:CH2	1:A:407:GLN:HG3	2.47	0.49
1:A:323:LYS:O	1:A:343:GLN:NE2	2.46	0.49
2:B:362:THR:HG22	2:B:366:LYS:HZ3	1.77	0.49
1:A:503:LEU:HD21	2:B:422:LEU:HD13	1.94	0.49
2:B:266:TRP:CZ3	2:B:426:TRP:HB3	2.48	0.48
1:A:457:TYR:CD2	1:A:484:LEU:HD13	2.49	0.48
1:A:161:GLN:NE2	2:B:140:PRO:HG3	2.28	0.48
2:B:298:GLU:H	2:B:298:GLU:CD	2.20	0.48
2:B:344:GLU:HB2	2:B:347:LYS:HG3	1.94	0.48
1:A:184:MET:HB3	1:A:185:ASP:H	1.52	0.48
1:A:27:THR:C	1:A:29:GLU:H	2.22	0.47
1:A:442:VAL:HB	1:A:481:ALA:HB1	1.95	0.47
1:A:171:PHE:HB2	1:A:208:HIS:ND1	2.29	0.47
1:A:438:GLU:HG3	1:A:461:LYS:HG3	1.96	0.47
2:B:163:SER:O	2:B:167:ILE:HG13	2.14	0.47
1:A:286:THR:O	1:A:287:LYS:C	2.57	0.47
1:A:447:ASN:HB3	1:A:450:THR:OG1	2.15	0.47
2:B:332:GLN:HG3	2:B:338:THR:HG23	1.96	0.47
1:A:203:GLU:O	1:A:207:GLN:HG2	2.15	0.47
2:B:85:GLN:HG3	2:B:88:TRP:CH2	2.50	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:202:ILE:O	1:A:206:ARG:HG3	2.14	0.46
2:B:328:GLU:HG3	2:B:342:TYR:HE2	1.79	0.46
1:A:406:TRP:CE2	2:B:420:PRO:HB3	2.49	0.46
2:B:354[B]:TYR:HE2	2:B:375:ILE:HG13	1.79	0.46
1:A:410:TRP:CD1	2:B:401:TRP:CE2	3.05	0.45
2:B:281:LYS:HA	2:B:284:ARG:HG3	1.97	0.45
1:A:89:GLU:H	1:A:89:GLU:HG2	1.47	0.45
1:A:320:ASP:OD2	1:A:323:LYS:HG3	2.17	0.45
2:B:101:LYS:O	2:B:236:PRO:HB2	2.16	0.45
1:A:110:ASP:O	1:A:217:PRO:HD3	2.17	0.44
1:A:356:ARG:O	1:A:356:ARG:HG3	2.18	0.44
2:B:247:PRO:O	2:B:307:ARG:NH2	2.48	0.44
2:B:284:ARG:O	2:B:287:LYS:NZ	2.45	0.44
1:A:498:ASP:HB2	1:A:538:ALA:HB2	1.99	0.44
1:A:532:TYR:CE2	1:A:534:ALA:HB2	2.53	0.44
2:B:65:LYS:NZ	2:B:65:LYS:H	2.15	0.44
2:B:126:LYS:HA	2:B:145:GLN:OE1	2.18	0.44
2:B:239:TRP:CD1	2:B:239:TRP:N	2.85	0.44
2:B:265:ASN:O	2:B:268:SER:OG	2.29	0.44
1:A:241:VAL:HB	1:A:266:TRP:HE1	1.81	0.43
1:A:253:THR:HG23	1:A:256:ASP:OD2	2.18	0.43
1:A:194:GLU:O	1:A:197:GLN:N	2.51	0.43
2:B:28:GLU:HB2	2:B:135:ILE:HD11	2.01	0.43
2:B:13:LYS:O	2:B:16:MET:HG3	2.18	0.43
1:A:60:VAL:HG22	1:A:75:VAL:HG22	2.01	0.43
1:A:5:ILE:HD13	1:A:167:ILE:HD11	2.00	0.42
2:B:22:LYS:O	2:B:22:LYS:HG3	2.19	0.42
1:A:171:PHE:CZ	1:A:205:LEU:HB2	2.55	0.42
1:A:357:MET:N	1:A:357:MET:SD	2.92	0.42
1:A:161:GLN:HE22	2:B:140:PRO:HG3	1.85	0.42
1:A:320:ASP:OD1	1:A:322:SER:OG	2.36	0.42
1:A:364:ASP:HB3	1:A:423:VAL:HG13	2.02	0.42
2:B:331:LYS:HB2	2:B:337:TRP:CZ3	2.55	0.42
2:B:356:ARG:HA	2:B:356:ARG:HD2	1.68	0.42
1:A:270:ILE:O	1:A:272:PRO:HD3	2.20	0.41
1:A:420:PRO:HA	1:A:421:PRO:C	2.45	0.41
2:B:103:LYS:HD3	2:B:103:LYS:HA	1.89	0.41
2:B:424:LYS:NZ	4:B:501:HOH:O	2.50	0.41
1:A:341:ILE:HG21	1:A:383:TRP:CH2	2.55	0.41
1:A:5:ILE:H	1:A:5:ILE:HG13	1.41	0.41
1:A:155:GLY:O	1:A:159:ILE:HG13	2.20	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:406:TRP:HH2	2:B:418:ASN:HA	1.85	0.41
2:B:50:ILE:HD11	2:B:143:ARG:HB3	2.02	0.41
2:B:282:LEU:H	2:B:282:LEU:HD12	1.84	0.41
1:A:303:LEU:O	1:A:307:ARG:HG3	2.21	0.41
2:B:337:TRP:O	2:B:353:LYS:HD3	2.21	0.41
1:A:246:LEU:HD22	1:A:260:LEU:CD1	2.51	0.41
1:A:429:LEU:HA	1:A:429:LEU:HD23	1.87	0.40
2:B:195:ILE:HD11	4:B:508:HOH:O	2.21	0.40
2:B:279:LEU:HD23	2:B:279:LEU:HA	1.86	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	547/556 (98%)	517 (94%)	30 (6%)	0	100	100
2	B	399/428 (93%)	384 (96%)	14 (4%)	1 (0%)	36	53
All	All	946/984 (96%)	901 (95%)	44 (5%)	1 (0%)	48	66

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	420	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	430/495 (87%)	382 (89%)	48 (11%)	6	11
2	B	339/390 (87%)	304 (90%)	35 (10%)	7	13
All	All	769/885 (87%)	686 (89%)	83 (11%)	6	11

All (83) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	20	LYS
1	A	30	LYS
1	A	50	ILE
1	A	58	THR
1	A	60	VAL
1	A	63	ILE
1	A	69	THR
1	A	70	LYS
1	A	89	GLU
1	A	90	VAL
1	A	91	GLN
1	A	101	LYS
1	A	103	LYS
1	A	111	VAL
1	A	142	ILE
1	A	178	ILE
1	A	184	MET
1	A	185	ASP
1	A	204	GLU
1	A	215	THR
1	A	216	THR
1	A	238	LYS
1	A	248	GLU
1	A	249	LYS
1	A	251	SER
1	A	253	THR
1	A	297	GLU
1	A	322	SER
1	A	324	ASP
1	A	331	LYS
1	A	338	THR
1	A	347	LYS
1	A	356	ARG
1	A	361	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	396	GLU
1	A	415	GLU
1	A	431	LYS
1	A	435	VAL
1	A	449	GLU
1	A	459	THR
1	A	473	THR
1	A	474	ASN
1	A	480	GLN
1	A	482	ILE
1	A	484	LEU
1	A	507	GLN
1	A	515	SER
1	A	516	GLU
2	B	7	THR
2	B	8	VAL
2	B	22	LYS
2	B	27	THR
2	B	36	GLU
2	B	48	SER
2	B	65	LYS
2	B	69	THR
2	B	72	ARG
2	B	163	SER
2	B	165	THR
2	B	174	GLN
2	B	175	ASN
2	B	180	ILE
2	B	249	LYS
2	B	251	SER
2	B	274	ILE
2	B	275	LYS
2	B	280	SER
2	B	286	THR
2	B	293	ILE
2	B	297	GLU
2	B	298	GLU
2	B	300	GLU
2	B	314	VAL
2	B	317	VAL
2	B	330	GLN
2	B	353	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	354[A]	TYR
2	B	354[B]	TYR
2	B	356	ARG
2	B	385	LYS
2	B	395	LYS
2	B	413	GLU
2	B	423	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	197	GLN
1	A	330	GLN
1	A	340	GLN
1	A	367	GLN
1	A	373	GLN
2	B	147	ASN
2	B	197	GLN
2	B	265	ASN
2	B	269	GLN
2	B	278	GLN
2	B	418	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	VWB	A	601	-	32,32,32	1.52	4 (12%)	42,43,43	1.06	2 (4%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	VWB	A	601	-	-	3/16/23/23	0/4/4/4

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	601	VWB	C1-N1	5.30	1.45	1.35
3	A	601	VWB	C22-C23	3.79	1.52	1.44
3	A	601	VWB	C19-C18	-2.21	1.38	1.43
3	A	601	VWB	O1-C1	-2.07	1.18	1.23

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	601	VWB	C4-C5-N1	2.65	107.91	103.44
3	A	601	VWB	C14-O3-C13	-2.55	112.09	119.10

There are no chirality outliers.


All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	601	VWB	C1-C6-C7-O2
3	A	601	VWB	C13-C8-O2-C7
3	A	601	VWB	C9-C8-O2-C7

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å ²)	Q < 0.9
1	A	551/556 (99%)	0.57	38 (6%)	23 20	54, 80, 122, 180	1 (0%)
2	B	406/428 (94%)	0.39	26 (6%)	25 22	45, 73, 116, 150	1 (0%)
All	All	957/984 (97%)	0.49	64 (6%)	24 21	45, 78, 118, 180	2 (0%)

All (64) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1	PRO	5.0
1	A	2	ILE	4.7
1	A	24	TRP	4.5
2	B	354[A]	TYR	4.3
1	A	31	ILE	4.2
2	B	422	LEU	3.5
2	B	6	GLU	3.5
1	A	135	ILE	3.4
1	A	60	VAL	3.4
1	A	398	TRP	3.4
2	B	5	ILE	3.4
2	B	419	THR	3.3
2	B	240	THR	3.2
1	A	543	GLY	3.1
2	B	232	TYR	3.0
1	A	137	ASN	3.0
2	B	87	PHE	2.9
2	B	241	VAL	2.9
2	B	423	VAL	2.8
1	A	357	MET	2.8
2	B	361	HIS	2.8
1	A	252	TRP	2.7
1	A	34	LEU	2.7
1	A	244	ILE	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	28	GLU	2.6
1	A	411	ILE	2.6
1	A	25	PRO	2.5
1	A	59	PRO	2.5
2	B	95	PRO	2.5
2	B	295	LEU	2.5
2	B	24	TRP	2.5
1	A	89	GLU	2.4
2	B	90	VAL	2.4
1	A	554	ALA	2.4
1	A	333	GLY	2.4
1	A	332	GLN	2.4
1	A	131	THR	2.4
2	B	94	ILE	2.4
1	A	295	LEU	2.4
2	B	299	ALA	2.3
2	B	421	PRO	2.3
1	A	58	THR	2.3
2	B	68	SER	2.3
2	B	278	GLN	2.2
2	B	428	GLN	2.2
1	A	413	GLU	2.2
2	B	276	VAL	2.2
1	A	402	TRP	2.2
1	A	410	TRP	2.2
1	A	552	VAL	2.1
1	A	91	GLN	2.1
1	A	286	THR	2.1
1	A	553	SER	2.1
1	A	289	LEU	2.1
2	B	273	GLY	2.1
2	B	274	ILE	2.1
1	A	298	GLU	2.1
1	A	132	ILE	2.1
2	B	88	TRP	2.1
1	A	130	PHE	2.0
1	A	23	GLN	2.0
1	A	124	PHE	2.0
1	A	548	VAL	2.0
2	B	97	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	VWB	A	601	29/29	0.91	0.12	64,71,77,78	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around VWB A 601:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.