



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 9, 2026 – 05:16 AM UTC

PDB ID : 3U7U / pdb_00003u7u
Title : Crystal structure of extracellular region of human epidermal growth factor receptor 4 in complex with neuregulin-1 beta
Authors : Liu, P.; Cleveland IV, T.E.; Bouyain, S.; Longo, P.A.; Leahy, D.J.
Deposited on : 2011-10-14
Resolution : 3.03 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.03 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3685 (3.08-3.00)
Clashscore	190562	4007 (3.08-3.00)
Ramachandran outliers	187476	3834 (3.08-3.00)
Sidechain outliers	187428	3836 (3.08-3.00)
RSRZ outliers	180081	3684 (3.08-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	615	
1	B	615	
1	C	615	
1	D	615	
1	E	615	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	615	79%15%
2	G	55	2% 82%9%9%
2	H	55	2% 73%20%7%
2	I	55	4% 69%22%9%
2	J	55	2% 71%22%7%
2	K	55	2% 75%15%11%
2	L	55	82%11%7%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 30527 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Receptor tyrosine-protein kinase erbB-4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	550	Total	C	N	O	S	0	0	0
			4284	2658	746	829	51			
1	B	601	Total	C	N	O	S	0	0	0
			4685	2908	815	904	58			
1	C	598	Total	C	N	O	S	0	0	0
			4649	2879	812	900	58			
1	D	600	Total	C	N	O	S	0	0	0
			4683	2906	815	904	58			
1	E	593	Total	C	N	O	S	0	0	0
			4615	2856	807	894	58			
1	F	591	Total	C	N	O	S	0	0	0
			4595	2843	804	890	58			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	524	ASP	GLY	conflict	UNP Q15303
B	524	ASP	GLY	conflict	UNP Q15303
C	524	ASP	GLY	conflict	UNP Q15303
D	524	ASP	GLY	conflict	UNP Q15303
E	524	ASP	GLY	conflict	UNP Q15303
F	524	ASP	GLY	conflict	UNP Q15303

- Molecule 2 is a protein called Neuregulin 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	G	50	Total	C	N	O	S	0	0	0
			391	243	67	73	8			
2	H	51	Total	C	N	O	S	0	0	0
			398	247	68	75	8			
2	I	50	Total	C	N	O	S	0	0	0
			391	243	67	73	8			

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	J	51	Total	C	N	O	S	0	0	0
			398	247	68	75	8			
2	K	49	Total	C	N	O	S	0	0	0
			385	240	66	71	8			
2	L	51	Total	C	N	O	S	0	0	0
			398	247	68	75	8			

- Molecule 3 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	B	1	Total	C	N	O	0	0
			14	8	1	5		
3	B	1	Total	C	N	O	0	0
			14	8	1	5		
3	B	1	Total	C	N	O	0	0
			14	8	1	5		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	B	1	Total	C	N	O	0	0
			14	8	1	5		
3	B	1	Total	C	N	O	0	0
			14	8	1	5		
3	C	1	Total	C	N	O	0	0
			14	8	1	5		
3	C	1	Total	C	N	O	0	0
			14	8	1	5		
3	D	1	Total	C	N	O	0	0
			14	8	1	5		
3	D	1	Total	C	N	O	0	0
			14	8	1	5		
3	D	1	Total	C	N	O	0	0
			14	8	1	5		
3	D	1	Total	C	N	O	0	0
			14	8	1	5		
3	D	1	Total	C	N	O	0	0
			14	8	1	5		
3	E	1	Total	C	N	O	0	0
			14	8	1	5		
3	E	1	Total	C	N	O	0	0
			14	8	1	5		
3	E	1	Total	C	N	O	0	0
			14	8	1	5		
3	F	1	Total	C	N	O	0	0
			14	8	1	5		
3	F	1	Total	C	N	O	0	0
			14	8	1	5		
3	F	1	Total	C	N	O	0	0
			14	8	1	5		
3	F	1	Total	C	N	O	0	0
			14	8	1	5		
3	F	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	30	Total	O	0	0
			30	30		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	63	Total 63	O 63	0	0
4	C	37	Total 37	O 37	0	0
4	D	55	Total 55	O 55	0	0
4	E	33	Total 33	O 33	0	0
4	F	52	Total 52	O 52	0	0
4	G	1	Total 1	O 1	0	0
4	H	9	Total 9	O 9	0	0
4	I	2	Total 2	O 2	0	0
4	J	3	Total 3	O 3	0	0
4	K	2	Total 2	O 2	0	0
4	L	4	Total 4	O 4	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Receptor tyrosine-protein kinase erbB-4




- Molecule 1: Receptor tyrosine-protein kinase erbB-4




- Molecule 1: Receptor tyrosine-protein kinase erbB-4




• Molecule 1: Receptor tyrosine-protein kinase erbB-4




• Molecule 1: Receptor tyrosine-protein kinase erbB-4




● Molecule 1: Receptor tyrosine-protein kinase erbB-4

Chain F:  79% 15% . .


● Molecule 2: Neuregulin 1

Chain G:  2% 82% 9% 9%


● Molecule 2: Neuregulin 1

Chain H:  2% 73% 20% 7%

● Molecule 2: Neuregulin 1


Chain I:  4% 69% 22% 9%

● Molecule 2: Neuregulin 1



Chain J:  2% 71% 22% 7%

● Molecule 2: Neuregulin 1

Chain K:  2% 75% 15% 11%



● Molecule 2: Neuregulin 1



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	85.73Å 223.51Å 146.92Å 90.00° 99.72° 90.00°	Depositor
Resolution (Å)	49.63 – 3.03 49.63 – 3.03	Depositor EDS
% Data completeness (in resolution range)	(Not available) (49.63-3.03) 98.7 (49.63-3.03)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.91 (at 3.01Å)	Xtriage
Refinement program	BUSTER-TNT BUSTER 2.8.0, BUSTER 2.8.0	Depositor
R, R_{free}	0.190 , 0.227 (Not available) , 0.237	Depositor DCC
R_{free} test set	5198 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	73.3	Xtriage
Anisotropy	0.361	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 85.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	30527	wwPDB-VP
Average B, all atoms (Å ²)	90.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.60% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.84	0/4366	1.32	26/5908 (0.4%)
1	B	0.88	2/4783 (0.0%)	1.35	43/6476 (0.7%)
1	C	0.84	1/4743 (0.0%)	1.32	33/6420 (0.5%)
1	D	0.85	1/4782 (0.0%)	1.33	36/6477 (0.6%)
1	E	0.85	2/4709 (0.0%)	1.34	26/6374 (0.4%)
1	F	0.88	1/4688 (0.0%)	1.35	40/6346 (0.6%)
2	G	0.84	0/397	1.37	2/528 (0.4%)
2	H	0.79	0/404	1.34	4/538 (0.7%)
2	I	0.75	0/397	1.32	4/528 (0.8%)
2	J	0.84	0/404	1.31	2/538 (0.4%)
2	K	0.75	0/391	1.38	6/520 (1.2%)
2	L	0.82	0/404	1.39	2/538 (0.4%)
All	All	0.85	7/30468 (0.0%)	1.34	224/41191 (0.5%)

The worst 5 of 7 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	F	35	MET	SD-CE	8.17	2.00	1.79
1	E	35	MET	CA-C	-5.56	1.47	1.53
1	B	226	ASN	CA-C	-5.50	1.49	1.52
1	C	244	ASN	CA-C	5.33	1.58	1.52
1	B	318	MET	SD-CE	5.24	1.92	1.79

The worst 5 of 224 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	F	47	ARG	N-CA-C	7.76	121.22	110.35
1	E	446	THR	CA-C-N	7.70	131.11	120.49
1	E	446	THR	C-N-CA	7.70	131.11	120.49
2	K	3	SER	CA-C-N	7.67	136.19	121.54
2	K	3	SER	C-N-CA	7.67	136.19	121.54

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4284	0	4089	40	0
1	B	4685	0	4432	51	0
1	C	4649	0	4407	47	0
1	D	4683	0	4427	54	0
1	E	4615	0	4365	49	0
1	F	4595	0	4348	49	0
2	G	391	0	373	1	0
2	H	398	0	380	4	0
2	I	391	0	373	4	0
2	J	398	0	380	5	0
2	K	385	0	368	1	0
2	L	398	0	380	2	0
3	A	56	0	51	0	0
3	B	84	0	78	2	0
3	C	28	0	26	0	0
3	D	70	0	65	7	0
3	E	42	0	39	0	0
3	F	84	0	78	2	0
4	A	30	0	0	0	0
4	B	63	0	0	2	0
4	C	37	0	0	1	0
4	D	55	0	0	2	0
4	E	33	0	0	2	0
4	F	52	0	0	2	0
4	G	1	0	0	0	0
4	H	9	0	0	0	0
4	I	2	0	0	0	0
4	J	3	0	0	0	0
4	K	2	0	0	0	0
4	L	4	0	0	0	0
All	All	30527	0	28659	285	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 285 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:143:ARG:O	1:D:145:PRO:HD3	1.69	0.90
1:C:145:PRO:HB2	1:F:254:PHE:HB3	1.52	0.88
1:E:3:VAL:HG13	1:E:35:MET:HG3	1.57	0.87
1:D:3:VAL:HG13	1:D:35:MET:HG3	1.56	0.87
1:D:206:HIS:HD2	1:D:208:GLU:H	1.20	0.86

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	544/615 (88%)	508 (93%)	31 (6%)	5 (1%)	14	44
1	B	593/615 (96%)	558 (94%)	33 (6%)	2 (0%)	36	67
1	C	590/615 (96%)	548 (93%)	37 (6%)	5 (1%)	16	46
1	D	592/615 (96%)	553 (93%)	35 (6%)	4 (1%)	18	49
1	E	585/615 (95%)	544 (93%)	38 (6%)	3 (0%)	24	57
1	F	583/615 (95%)	546 (94%)	32 (6%)	5 (1%)	14	44
2	G	46/55 (84%)	43 (94%)	2 (4%)	1 (2%)	5	23
2	H	47/55 (86%)	43 (92%)	3 (6%)	1 (2%)	5	24
2	I	46/55 (84%)	43 (94%)	2 (4%)	1 (2%)	5	23
2	J	47/55 (86%)	44 (94%)	2 (4%)	1 (2%)	5	24
2	K	45/55 (82%)	41 (91%)	3 (7%)	1 (2%)	5	23
2	L	47/55 (86%)	44 (94%)	2 (4%)	1 (2%)	5	24
All	All	3765/4020 (94%)	3515 (93%)	220 (6%)	30 (1%)	16	46

5 of 30 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	149	ASN
1	B	149	ASN
1	C	149	ASN
1	C	159	SER
1	D	128	LYS

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	488/544 (90%)	459 (94%)	29 (6%)	18	47
1	B	534/544 (98%)	506 (95%)	28 (5%)	21	51
1	C	531/544 (98%)	504 (95%)	27 (5%)	21	52
1	D	534/544 (98%)	506 (95%)	28 (5%)	21	51
1	E	527/544 (97%)	498 (94%)	29 (6%)	19	50
1	F	525/544 (96%)	498 (95%)	27 (5%)	21	52
2	G	45/49 (92%)	44 (98%)	1 (2%)	45	71
2	H	46/49 (94%)	45 (98%)	1 (2%)	45	71
2	I	45/49 (92%)	44 (98%)	1 (2%)	45	71
2	J	46/49 (94%)	44 (96%)	2 (4%)	26	57
2	K	44/49 (90%)	43 (98%)	1 (2%)	44	70
2	L	46/49 (94%)	45 (98%)	1 (2%)	45	71
All	All	3411/3558 (96%)	3236 (95%)	175 (5%)	21	52

5 of 175 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	43	ILE
1	F	43	ILE
1	E	201	VAL
1	E	429	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	F	203	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 85 such sidechains are listed below:

Mol	Chain	Res	Type
1	E	206	HIS
1	F	144	ASN
1	E	421	GLN
1	E	606	HIS
1	F	354	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

26 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	NAG	D	1001	1	14,14,15	1.50	3 (21%)	17,19,21	2.93	7 (41%)
3	NAG	E	1003	1	14,14,15	2.34	4 (28%)	17,19,21	2.10	6 (35%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	NAG	D	1003	1	14,14,15	1.26	1 (7%)	17,19,21	2.15	5 (29%)
3	NAG	F	1005	1	14,14,15	1.80	3 (21%)	17,19,21	2.38	7 (41%)
3	NAG	B	1003	1	14,14,15	1.69	2 (14%)	17,19,21	2.23	8 (47%)
3	NAG	A	1001	1	14,14,15	1.27	2 (14%)	17,19,21	3.21	9 (52%)
3	NAG	B	1001	1	14,14,15	1.35	1 (7%)	17,19,21	3.15	8 (47%)
3	NAG	D	1004	1	14,14,15	1.71	3 (21%)	17,19,21	2.18	5 (29%)
3	NAG	E	1002	1	14,14,15	1.29	1 (7%)	17,19,21	2.04	6 (35%)
3	NAG	F	1004	1	14,14,15	2.33	4 (28%)	17,19,21	2.72	8 (47%)
3	NAG	F	1002	1	14,14,15	1.27	1 (7%)	17,19,21	2.28	6 (35%)
3	NAG	D	1005	1	14,14,15	1.48	2 (14%)	17,19,21	3.00	9 (52%)
3	NAG	B	1002	1	14,14,15	1.58	2 (14%)	17,19,21	2.32	6 (35%)
3	NAG	B	1004	1	14,14,15	3.68	8 (57%)	17,19,21	4.44	10 (58%)
3	NAG	F	1001	1	14,14,15	1.64	3 (21%)	17,19,21	3.69	10 (58%)
3	NAG	A	1004	1	14,14,15	2.28	3 (21%)	17,19,21	4.09	11 (64%)
3	NAG	C	1002	1	14,14,15	1.62	3 (21%)	17,19,21	2.00	6 (35%)
3	NAG	B	1006	1	14,14,15	1.90	5 (35%)	17,19,21	2.58	8 (47%)
3	NAG	B	1005	1	14,14,15	2.72	5 (35%)	17,19,21	2.80	8 (47%)
3	NAG	E	1001	1	14,14,15	1.31	2 (14%)	17,19,21	3.33	9 (52%)
3	NAG	A	1003	1	14,14,15	1.25	1 (7%)	17,19,21	1.78	5 (29%)
3	NAG	F	1003	1	14,14,15	1.09	1 (7%)	17,19,21	2.17	7 (41%)
3	NAG	D	1002	1	14,14,15	1.73	4 (28%)	17,19,21	2.07	5 (29%)
3	NAG	A	1002	1	14,14,15	1.64	3 (21%)	17,19,21	2.53	9 (52%)
3	NAG	C	1001	1	14,14,15	1.18	0	17,19,21	3.54	8 (47%)
3	NAG	F	1006	1	14,14,15	2.25	4 (28%)	17,19,21	1.24	2 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	D	1001	1	-	2/6/23/26	0/1/1/1
3	NAG	E	1003	1	-	2/6/23/26	0/1/1/1
3	NAG	D	1003	1	-	1/6/23/26	0/1/1/1
3	NAG	F	1005	1	-	1/6/23/26	0/1/1/1
3	NAG	B	1003	1	-	4/6/23/26	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	A	1001	1	-	2/6/23/26	0/1/1/1
3	NAG	B	1001	1	-	2/6/23/26	0/1/1/1
3	NAG	D	1004	1	-	2/6/23/26	0/1/1/1
3	NAG	E	1002	1	-	0/6/23/26	0/1/1/1
3	NAG	F	1004	1	-	1/6/23/26	0/1/1/1
3	NAG	F	1002	1	-	0/6/23/26	0/1/1/1
3	NAG	D	1005	1	-	0/6/23/26	0/1/1/1
3	NAG	B	1002	1	-	0/6/23/26	0/1/1/1
3	NAG	B	1004	1	-	0/6/23/26	0/1/1/1
3	NAG	F	1001	1	-	2/6/23/26	0/1/1/1
3	NAG	A	1004	1	-	2/6/23/26	0/1/1/1
3	NAG	C	1002	1	-	0/6/23/26	0/1/1/1
3	NAG	B	1006	1	-	0/6/23/26	0/1/1/1
3	NAG	B	1005	1	-	1/6/23/26	0/1/1/1
3	NAG	E	1001	1	-	2/6/23/26	0/1/1/1
3	NAG	A	1003	1	-	1/6/23/26	0/1/1/1
3	NAG	F	1003	1	-	2/6/23/26	0/1/1/1
3	NAG	D	1002	1	-	0/6/23/26	0/1/1/1
3	NAG	A	1002	1	-	0/6/23/26	0/1/1/1
3	NAG	C	1001	1	-	2/6/23/26	0/1/1/1
3	NAG	F	1006	1	-	1/6/23/26	0/1/1/1

The worst 5 of 71 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	1004	NAG	C1-C2	9.54	1.65	1.52
3	A	1004	NAG	C1-C2	6.61	1.61	1.52
3	B	1005	NAG	C1-C2	6.41	1.61	1.52
3	F	1004	NAG	C1-C2	6.20	1.60	1.52
3	E	1003	NAG	C1-C2	5.54	1.59	1.52

The worst 5 of 188 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	1004	NAG	C2-N2-C7	11.60	138.45	122.90
3	A	1004	NAG	C1-O5-C5	-9.20	99.85	112.19
3	C	1001	NAG	C1-C2-N2	-8.89	96.43	110.43
3	A	1001	NAG	C1-C2-N2	-8.79	96.58	110.43
3	F	1001	NAG	C1-C2-N2	-8.65	96.81	110.43

There are no chirality outliers.

5 of 30 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	F	1004	NAG	C3-C2-N2-C7
3	A	1001	NAG	O5-C5-C6-O6
3	B	1001	NAG	O5-C5-C6-O6
3	C	1001	NAG	O5-C5-C6-O6
3	D	1001	NAG	O5-C5-C6-O6

There are no ring outliers.

4 monomers are involved in 11 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	D	1004	NAG	2	0
3	F	1004	NAG	2	0
3	D	1005	NAG	5	0
3	B	1005	NAG	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2	OWAB(Å ²)	Q < 0.9
1	A	550/615 (89%)	-0.21	1 (0%) 91 83	47, 85, 249, 280	0
1	B	601/615 (97%)	-0.33	2 (0%) 90 79	45, 69, 115, 151	0
1	C	598/615 (97%)	-0.13	7 (1%) 76 54	50, 93, 173, 229	0
1	D	600/615 (97%)	-0.36	3 (0%) 87 72	45, 77, 117, 172	0
1	E	593/615 (96%)	-0.21	2 (0%) 90 79	45, 93, 164, 228	0
1	F	591/615 (96%)	-0.33	3 (0%) 87 72	43, 72, 155, 278	0
2	G	50/55 (90%)	0.15	1 (2%) 65 41	68, 98, 142, 151	0
2	H	51/55 (92%)	-0.16	1 (1%) 65 41	60, 85, 130, 140	0
2	I	50/55 (90%)	0.01	2 (4%) 42 23	85, 104, 143, 147	0
2	J	51/55 (92%)	-0.03	1 (1%) 65 41	75, 100, 136, 140	0
2	K	49/55 (89%)	0.13	1 (2%) 65 41	87, 110, 143, 158	0
2	L	51/55 (92%)	-0.12	0 100 100	62, 86, 126, 143	0
All	All	3835/4020 (95%)	-0.24	24 (0%) 85 69	43, 83, 156, 280	0

The worst 5 of 24 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	G	28	LEU	3.3
1	F	201	VAL	3.1
1	C	524	ASP	3.0
1	D	201	VAL	3.0
1	C	201	VAL	2.9

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	NAG	B	1003	14/15	0.43	0.11	140,145,150,152	0
3	NAG	F	1005	14/15	0.66	0.14	126,130,135,136	0
3	NAG	D	1003	14/15	0.67	0.10	126,131,134,137	0
3	NAG	A	1003	14/15	0.67	0.10	145,151,158,160	0
3	NAG	D	1004	14/15	0.74	0.15	125,129,134,135	0
3	NAG	B	1004	14/15	0.76	0.16	89,93,97,97	0
3	NAG	D	1005	14/15	0.79	0.15	108,111,116,118	0
3	NAG	A	1004	14/15	0.80	0.13	117,121,125,127	0
3	NAG	F	1006	14/15	0.81	0.14	127,129,135,136	0
3	NAG	B	1005	14/15	0.82	0.10	100,105,108,111	0
3	NAG	E	1003	14/15	0.82	0.14	147,149,154,158	0
3	NAG	B	1006	14/15	0.84	0.12	118,121,125,125	0
3	NAG	C	1001	14/15	0.84	0.10	72,75,80,81	0
3	NAG	D	1001	14/15	0.84	0.12	66,70,74,74	0
3	NAG	F	1003	14/15	0.85	0.09	118,123,128,130	0
3	NAG	E	1001	14/15	0.87	0.12	66,70,74,74	0
3	NAG	A	1001	14/15	0.88	0.09	69,72,77,78	0
3	NAG	A	1002	14/15	0.90	0.10	71,76,78,80	0
3	NAG	B	1002	14/15	0.90	0.10	81,86,88,89	0
3	NAG	D	1002	14/15	0.92	0.10	73,77,80,81	0
3	NAG	F	1004	14/15	0.92	0.09	77,81,84,85	0
3	NAG	C	1002	14/15	0.92	0.09	72,77,81,82	0
3	NAG	F	1001	14/15	0.92	0.07	55,59,63,63	0
3	NAG	E	1002	14/15	0.93	0.09	77,82,85,87	0
3	NAG	F	1002	14/15	0.93	0.09	70,74,77,78	0
3	NAG	B	1001	14/15	0.93	0.07	60,64,67,68	0

6.5 Other polymers [i](#)

There are no such residues in this entry.