



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 1, 2026 – 07:00 PM UTC

PDB ID : 4U7U / pdb_00004u7u
Title : Crystal structure of RNA-guided immune Cascade complex from E.coli
Authors : Zhao, H.; Sheng, G.; Wang, J.; Wang, M.; Bunkoczi, G.; Gong, W.; Wei, Z.; Wang, Y.
Deposited on : 2014-07-31
Resolution : 3.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.






Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2672 (3.00-3.00)
Clashscore	190562	2977 (3.00-3.00)
Ramachandran outliers	187476	2877 (3.00-3.00)
Sidechain outliers	187428	2880 (3.00-3.00)
RSRZ outliers	180081	2671 (3.00-3.00)
RNA backbone	3983	1109 (3.20-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	502	74% 21% . .
1	M	502	80% 15% . .
2	B	160	81% 17% .
2	C	160	84% 11% . .

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	N	160	
2	O	160	
3	D	201	
3	P	201	
4	E	363	
4	F	363	
4	G	363	
4	H	363	
4	I	363	
4	J	363	
4	Q	363	
4	R	363	
4	S	363	
4	T	363	
4	U	363	
4	V	363	
5	K	224	
5	W	224	
6	L	61	
6	X	61	

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 54555 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CRISPR system Cascade subunit CasA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	480	Total	C	N	O	S	0	0	0
			3739	2386	659	674	20			
1	M	489	Total	C	N	O	S	0	0	0
			3812	2428	673	691	20			

- Molecule 2 is a protein called CRISPR system Cascade subunit CasB.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	157	Total	C	N	O	S	0	0	0
			1291	808	249	227	7			
2	C	157	Total	C	N	O	S	0	0	0
			1292	809	249	227	7			
2	N	152	Total	C	N	O	S	0	0	0
			1246	784	240	215	7			
2	O	156	Total	C	N	O	S	0	0	0
			1247	785	234	221	7			

- Molecule 3 is a protein called CRISPR system Cascade subunit CasE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	D	191	Total	C	N	O	S	0	0	0
			1451	931	255	258	7			
3	P	191	Total	C	N	O	S	0	0	0
			1459	936	258	258	7			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
D	-1	ALA	-	expression tag	UNP Q46897
D	0	TRP	-	expression tag	UNP Q46897
P	-1	ALA	-	expression tag	UNP Q46897
P	0	TRP	-	expression tag	UNP Q46897

- Molecule 4 is a protein called CRISPR system Cascade subunit CasC.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	E	357	Total	C	N	O	S	0	0	0
			2752	1721	490	525	16			
4	F	359	Total	C	N	O	S	0	0	0
			2760	1727	489	528	16			
4	G	356	Total	C	N	O	S	0	0	0
			2753	1720	490	527	16			
4	H	363	Total	C	N	O	S	0	0	0
			2774	1736	493	529	16			
4	I	355	Total	C	N	O	S	0	0	0
			2731	1711	485	519	16			
4	J	351	Total	C	N	O	S	0	0	0
			2680	1678	473	513	16			
4	Q	362	Total	C	N	O	S	0	0	0
			2790	1744	495	535	16			
4	R	359	Total	C	N	O	S	0	0	0
			2736	1712	484	524	16			
4	S	361	Total	C	N	O	S	0	0	0
			2773	1734	494	529	16			
4	T	360	Total	C	N	O	S	0	0	0
			2777	1738	494	529	16			
4	U	354	Total	C	N	O	S	0	0	0
			2723	1704	482	521	16			
4	V	355	Total	C	N	O	S	0	0	0
			2725	1704	482	523	16			

- Molecule 5 is a protein called CRISPR system Cascade subunit CasD.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
5	K	218	Total	C	N	O	S	0	0	0
			1722	1090	305	318	9			
5	W	218	Total	C	N	O	S	0	0	0
			1726	1093	306	318	9			


- Molecule 6 is a RNA chain called crRNA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
6	L	61	Total	C	N	O	P	0	0	0
			1298	580	230	428	60			
6	X	61	Total	C	N	O	P	0	0	0
			1298	580	230	428	60			


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: CRISPR system Cascade subunit CasA




• Molecule 1: CRISPR system Cascade subunit CasA




• Molecule 2: CRISPR system Cascade subunit CasB





- Molecule 2: CRISPR system Cascade subunit CasB




- Molecule 2: CRISPR system Cascade subunit CasB




- Molecule 2: CRISPR system Cascade subunit CasB




- Molecule 3: CRISPR system Cascade subunit CasE




- Molecule 3: CRISPR system Cascade subunit CasE




- Molecule 4: CRISPR system Cascade subunit CasC




- Molecule 4: CRISPR system Cascade subunit CasC




- Molecule 4: CRISPR system Cascade subunit CasC




- Molecule 4: CRISPR system Cascade subunit CasC





- Molecule 4: CRISPR system Cascade subunit CasC








• Molecule 4: CRISPR system Cascade subunit CasC




• Molecule 4: CRISPR system Cascade subunit CasC




• Molecule 4: CRISPR system Cascade subunit CasC




• Molecule 4: CRISPR system Cascade subunit CasC




• Molecule 4: CRISPR system Cascade subunit CasC



• Molecule 4: CRISPR system Cascade subunit CasC





• Molecule 4: CRISPR system Cascade subunit CasC







- Molecule 5: CRISPR system Cascade subunit CasD



- Molecule 5: CRISPR system Cascade subunit CasD



- Molecule 6: crRNA



- Molecule 6: crRNA



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	111.41Å 118.14Å 225.87Å 92.23° 93.55° 106.06°	Depositor
Resolution (Å)	43.80 – 3.00 43.80 – 3.00	Depositor EDS
% Data completeness (in resolution range)	88.8 (43.80-3.00) 88.7 (43.80-3.00)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.22 (at 3.01Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8.4_1496)	Depositor
R, R_{free}	0.179 , 0.218 0.186 , 0.222	Depositor DCC
R_{free} test set	9794 reflections (4.45%)	wwPDB-VP
Wilson B-factor (Å ²)	55.6	Xtriage
Anisotropy	0.128	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 39.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	54555	wwPDB-VP
Average B, all atoms (Å ²)	57.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.83% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.58	0/3825	0.88	10/5197 (0.2%)
1	M	0.84	1/3901 (0.0%)	0.88	9/5303 (0.2%)
2	B	0.82	0/1316	0.83	1/1777 (0.1%)
2	C	0.81	1/1317 (0.1%)	1.03	6/1778 (0.3%)
2	N	0.68	0/1270	0.83	2/1714 (0.1%)
2	O	0.62	0/1272	0.96	9/1724 (0.5%)
3	D	0.64	0/1480	1.08	7/2011 (0.3%)
3	P	0.54	0/1488	0.97	3/2019 (0.1%)
4	E	0.83	1/2795 (0.0%)	0.82	1/3768 (0.0%)
4	F	0.82	0/2803	0.83	4/3785 (0.1%)
4	G	0.87	0/2796	0.81	0/3771
4	H	0.87	0/2819	0.85	0/3809
4	I	0.84	0/2774	0.84	4/3744 (0.1%)
4	J	0.72	0/2722	0.84	4/3681 (0.1%)
4	Q	0.97	0/2834	0.84	2/3823 (0.1%)
4	R	0.81	0/2779	0.88	7/3758 (0.2%)
4	S	0.81	0/2817	0.85	1/3802 (0.0%)
4	T	0.82	0/2821	0.84	2/3805 (0.1%)
4	U	0.70	0/2766	0.91	7/3737 (0.2%)
4	V	0.97	0/2768	0.93	6/3741 (0.2%)
5	K	0.63	0/1764	0.88	2/2398 (0.1%)
5	W	0.90	0/1768	0.93	5/2402 (0.2%)
6	L	0.60	0/1450	0.58	0/2259
6	X	0.62	0/1450	0.56	0/2259
All	All	0.79	3/55795 (0.0%)	0.86	92/76065 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
4	R	0	2

Continued on next page...

Continued from previous page...

Mol	Chain	#Chirality outliers	#Planarity outliers
4	U	0	2
All	All	0	4

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	112	ALA	N-CA	-5.65	1.39	1.46
4	E	19	ASN	C-O	-5.24	1.17	1.24
1	M	170	THR	C-O	-5.04	1.21	1.23

The worst 5 of 92 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	112	ALA	N-CA-C	-17.31	92.42	111.28
3	D	140	HIS	CA-C-N	12.75	135.78	119.84
3	D	140	HIS	C-N-CA	12.75	135.78	119.84
2	C	108	ALA	N-CA-C	9.94	124.72	112.59
2	C	110	ARG	N-CA-C	-9.57	97.90	110.43

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
4	R	132	ASP	Peptide
4	R	338	ASP	Peptide
4	U	335	SER	Peptide
4	U	94	LEU	Peptide

5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3739	0	3705	67	0
1	M	3812	0	3775	50	0
2	B	1291	0	1295	17	0
2	C	1292	0	1300	26	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	N	1246	0	1253	16	0
2	O	1247	0	1226	19	0
3	D	1451	0	1447	38	0
3	P	1459	0	1466	34	0
4	E	2752	0	2729	36	0
4	F	2760	0	2730	42	0
4	G	2753	0	2727	36	0
4	H	2774	0	2731	36	0
4	I	2731	0	2714	38	0
4	J	2680	0	2624	41	0
4	Q	2790	0	2764	44	0
4	R	2736	0	2672	44	0
4	S	2773	0	2734	49	0
4	T	2777	0	2761	38	0
4	U	2723	0	2690	90	0
4	V	2725	0	2682	41	0
5	K	1722	0	1700	33	0
5	W	1726	0	1711	33	0
6	L	1298	0	658	13	0
6	X	1298	0	658	15	0
All	All	54555	0	52752	813	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

The worst 5 of 813 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:U:82:ARG:CG	4:U:83:PHE:CE2	1.79	1.60
4:U:82:ARG:CG	4:U:83:PHE:HE2	1.03	1.58
4:U:82:ARG:C	4:U:83:PHE:CD2	1.88	1.51
4:U:82:ARG:CB	4:U:83:PHE:CE2	1.92	1.50
4:U:82:ARG:C	4:U:83:PHE:HD2	1.23	1.44

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	470/502 (94%)	432 (92%)	36 (8%)	2 (0%)	30	65
1	M	485/502 (97%)	457 (94%)	24 (5%)	4 (1%)	16	50
2	B	155/160 (97%)	149 (96%)	6 (4%)	0	100	100
2	C	155/160 (97%)	147 (95%)	8 (5%)	0	100	100
2	N	148/160 (92%)	140 (95%)	7 (5%)	1 (1%)	18	53
2	O	154/160 (96%)	147 (96%)	7 (4%)	0	100	100
3	D	185/201 (92%)	178 (96%)	7 (4%)	0	100	100
3	P	185/201 (92%)	176 (95%)	9 (5%)	0	100	100
4	E	353/363 (97%)	333 (94%)	20 (6%)	0	100	100
4	F	355/363 (98%)	335 (94%)	20 (6%)	0	100	100
4	G	352/363 (97%)	335 (95%)	17 (5%)	0	100	100
4	H	361/363 (99%)	348 (96%)	12 (3%)	1 (0%)	36	70
4	I	351/363 (97%)	335 (95%)	16 (5%)	0	100	100
4	J	345/363 (95%)	324 (94%)	21 (6%)	0	100	100
4	Q	358/363 (99%)	347 (97%)	10 (3%)	1 (0%)	36	70
4	R	355/363 (98%)	335 (94%)	20 (6%)	0	100	100
4	S	357/363 (98%)	343 (96%)	13 (4%)	1 (0%)	36	70
4	T	356/363 (98%)	343 (96%)	13 (4%)	0	100	100
4	U	350/363 (96%)	325 (93%)	23 (7%)	2 (1%)	21	56
4	V	351/363 (97%)	331 (94%)	20 (6%)	0	100	100
5	K	216/224 (96%)	211 (98%)	5 (2%)	0	100	100
5	W	216/224 (96%)	210 (97%)	5 (2%)	1 (0%)	24	60
All	All	6613/6850 (96%)	6281 (95%)	319 (5%)	13 (0%)	43	76

5 of 13 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
4	U	336	ASP
5	W	45	GLY
1	M	160	GLY
1	M	321	ASN
1	M	488	LYS

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	401/426 (94%)	385 (96%)	16 (4%)	28	62
1	M	409/426 (96%)	395 (97%)	14 (3%)	32	66
2	B	136/138 (99%)	135 (99%)	1 (1%)	76	86
2	C	136/138 (99%)	132 (97%)	4 (3%)	37	70
2	N	130/138 (94%)	125 (96%)	5 (4%)	29	63
2	O	128/138 (93%)	123 (96%)	5 (4%)	28	62
3	D	151/171 (88%)	141 (93%)	10 (7%)	15	46
3	P	152/171 (89%)	143 (94%)	9 (6%)	18	50
4	E	289/298 (97%)	281 (97%)	8 (3%)	38	70
4	F	290/298 (97%)	280 (97%)	10 (3%)	32	66
4	G	291/298 (98%)	286 (98%)	5 (2%)	53	78
4	H	288/298 (97%)	282 (98%)	6 (2%)	47	75
4	I	287/298 (96%)	283 (99%)	4 (1%)	59	80
4	J	279/298 (94%)	273 (98%)	6 (2%)	45	74
4	Q	294/298 (99%)	286 (97%)	8 (3%)	39	71
4	R	282/298 (95%)	277 (98%)	5 (2%)	51	77
4	S	290/298 (97%)	283 (98%)	7 (2%)	43	73
4	T	293/298 (98%)	290 (99%)	3 (1%)	68	84
4	U	287/298 (96%)	267 (93%)	20 (7%)	14	44
4	V	286/298 (96%)	274 (96%)	12 (4%)	26	61

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
5	K	185/192 (96%)	179 (97%)	6 (3%)	34	67
5	W	186/192 (97%)	178 (96%)	8 (4%)	26	60
All	All	5470/5706 (96%)	5298 (97%)	172 (3%)	35	68

5 of 172 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
4	Q	220	SER
4	U	126	VAL
4	R	183	SER
4	T	269	THR
4	U	335	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 72 such sidechains are listed below:

Mol	Chain	Res	Type
4	S	359	ASN
5	W	163	GLN
4	T	67	HIS
4	V	13	HIS
4	G	274	ASN

5.3.3 RNA ⓘ

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
6	L	60/61 (98%)	20 (33%)	2 (3%)
6	X	60/61 (98%)	20 (33%)	2 (3%)
All	All	120/122 (98%)	40 (33%)	4 (3%)

5 of 40 RNA backbone outliers are listed below:

Mol	Chain	Res	Type
6	L	2	U
6	L	4	A
6	L	9	G
6	L	10	G
6	L	15	C

All (4) RNA pucker outliers are listed below:

Mol	Chain	Res	Type
6	L	27	A
6	L	45	U
6	X	15	C
6	X	45	U

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	480/502 (95%)	0.02	4 (0%) 82 64	37, 80, 111, 135	0
1	M	489/502 (97%)	-0.39	1 (0%) 91 83	26, 42, 68, 122	0
2	B	157/160 (98%)	-0.36	2 (1%) 75 53	30, 44, 84, 120	0
2	C	157/160 (98%)	-0.17	2 (1%) 75 53	32, 52, 85, 115	0
2	N	152/160 (95%)	-0.20	0 100 100	40, 56, 85, 114	0
2	O	156/160 (97%)	0.21	4 (2%) 57 34	53, 84, 124, 146	0
3	D	191/201 (95%)	0.31	9 (4%) 36 19	41, 70, 112, 122	0
3	P	191/201 (95%)	0.37	9 (4%) 36 19	52, 78, 111, 127	0
4	E	357/363 (98%)	-0.29	6 (1%) 69 45	30, 48, 78, 104	0
4	F	359/363 (98%)	-0.29	3 (0%) 82 64	28, 48, 86, 121	0
4	G	356/363 (98%)	-0.40	2 (0%) 85 69	24, 44, 76, 101	0
4	H	363/363 (100%)	-0.21	5 (1%) 73 51	25, 47, 93, 116	0
4	I	355/363 (97%)	-0.33	0 100 100	26, 47, 79, 104	0
4	J	351/363 (96%)	-0.08	4 (1%) 78 57	35, 62, 110, 129	0
4	Q	362/363 (99%)	-0.37	5 (1%) 73 51	23, 39, 72, 114	0
4	R	359/363 (98%)	-0.11	10 (2%) 55 32	27, 50, 119, 155	0
4	S	361/363 (99%)	-0.24	4 (1%) 78 57	30, 52, 99, 118	0
4	T	360/363 (99%)	-0.32	3 (0%) 82 64	32, 50, 75, 121	0
4	U	354/363 (97%)	-0.01	3 (0%) 82 64	41, 68, 129, 146	0
4	V	355/363 (97%)	-0.40	4 (1%) 78 57	21, 38, 85, 120	0
5	K	218/224 (97%)	-0.07	2 (0%) 81 61	44, 67, 90, 106	0
5	W	218/224 (97%)	-0.49	0 100 100	22, 35, 53, 81	0
6	L	61/61 (100%)	-0.68	0 100 100	31, 47, 101, 118	0
6	X	61/61 (100%)	-0.55	0 100 100	25, 50, 120, 150	0

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
All	All	6823/6972 (97%)	-0.21	82 (1%)	76 55	21, 52, 103, 155	0

The worst 5 of 82 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
4	T	336	ASP	5.4
4	F	213	HIS	5.1
4	R	209	GLN	4.9
4	T	209	GLN	4.4
1	A	200	PRO	4.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.