



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 4, 2026 – 10:32 PM UTC

PDB ID : 5U7W / pdb_00005u7w
Title : Crystal structure of a nucleoside triphosphate diphosphohydrolase (NTPDase) from the legume *Trifolium repens* in complex with adenine and phosphate
Authors : Cumming, M.H.; Summers, E.L.; Oulavallickal, T.; Roberts, N.; Arcus, V.L.
Deposited on : 2016-12-12
Resolution : 1.76 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.76 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3183 (1.76-1.76)
Clashscore	190562	3299 (1.76-1.76)
Ramachandran outliers	187476	3274 (1.76-1.76)
Sidechain outliers	187428	3274 (1.76-1.76)
RSRZ outliers	180081	3183 (1.76-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	426	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	ADE	A	501	-	X	-	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	PO4	A	502	-	-	X	-

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 3520 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Apyrase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	405	3108	1997	509	592	10	0	6	0

There are 9 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MET	-	initiating methionine	UNP B9U139
A	419	VAL	-	expression tag	UNP B9U139
A	420	GLU	-	expression tag	UNP B9U139
A	421	HIS	-	expression tag	UNP B9U139
A	422	HIS	-	expression tag	UNP B9U139
A	423	HIS	-	expression tag	UNP B9U139
A	424	HIS	-	expression tag	UNP B9U139
A	425	HIS	-	expression tag	UNP B9U139
A	426	HIS	-	expression tag	UNP B9U139

- Molecule 2 is ADENINE (CCD ID: ADE) (formula: C₅H₅N₅).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	N	0	0
			10	5	5		

- Molecule 3 is PHOSPHATE ION (CCD ID: PO4) (formula: O_4P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	P	0	0
			5	4	1		
3	A	1	Total	O	P	0	0
			5	4	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	392	Total	O	0	0
			392	392		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Apyrase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	53.91Å 52.81Å 72.18Å 90.00° 94.96° 90.00°	Depositor
Resolution (Å)	53.71 – 1.76 53.71 – 1.76	Depositor EDS
% Data completeness (in resolution range)	93.2 (53.71-1.76) 93.2 (53.71-1.76)	Depositor EDS
R_{merge}	0.14	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.64 (at 1.76Å)	Xtriage
Refinement program	REFMAC 5.7.0032	Depositor
R, R_{free}	0.181 , 0.224 0.190 , 0.231	Depositor DCC
R_{free} test set	1887 reflections (4.68%)	wwPDB-VP
Wilson B-factor (Å ²)	17.8	Xtriage
Anisotropy	0.122	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 37.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	3520	wwPDB-VP
Average B, all atoms (Å ²)	20.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.22% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ADE, PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.31	11/3194 (0.3%)	1.20	11/4338 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	159	LEU	CA-CB	-6.34	1.44	1.53
1	A	304	LEU	C-O	-5.99	1.18	1.24
1	A	292	GLN	C-O	-5.89	1.15	1.24
1	A	224	GLY	N-CA	5.83	1.53	1.45
1	A	215	ARG	CZ-NH1	5.59	1.40	1.32
1	A	398	LEU	C-O	-5.45	1.17	1.24
1	A	125	ILE	C-O	-5.22	1.18	1.24
1	A	139	ILE	N-CA	-5.17	1.40	1.46
1	A	353	VAL	CA-CB	-5.16	1.51	1.54
1	A	38	VAL	C-O	-5.12	1.18	1.24
1	A	124	SER	CA-C	-5.12	1.46	1.52

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	125	ILE	CB-CA-C	-6.30	102.96	111.15
1	A	293	LYS	N-CA-C	6.23	118.16	111.36

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	126	ILE	CB-CA-C	6.10	119.64	112.68
1	A	63	ILE	CB-CA-C	-6.09	107.89	114.35
1	A	16	THR	CB-CA-C	-5.96	99.63	109.99
1	A	312	ASP	CA-C-N	-5.85	113.66	119.92
1	A	312	ASP	C-N-CA	-5.85	113.66	119.92
1	A	271	TYR	CA-C-N	-5.80	114.00	119.85
1	A	271	TYR	C-N-CA	-5.80	114.00	119.85
1	A	189	ILE	N-CA-C	5.57	115.91	108.11
1	A	129	THR	N-CA-CB	5.02	117.59	110.16

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	127	ASP	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3108	0	3044	64	0
2	A	10	0	4	0	0
3	A	10	0	0	2	0
4	A	392	0	0	23	1
All	All	3520	0	3048	65	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

All (65) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:140:ASN:HD21	1:A:154:VAL:H	1.21	0.88
1:A:409:PRO:O	1:A:410:LYS:CB	2.22	0.87
3:A:502:PO4:O1	4:A:601:HOH:O	1.92	0.86

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:275:TYR:HB2	4:A:750:HOH:O	1.77	0.85
1:A:243:GLU:HG3	4:A:826:HOH:O	1.77	0.82
1:A:91:LEU:HD22	1:A:125:ILE:HD11	1.64	0.80
1:A:175:ALA:HA	1:A:189:ILE:HD13	1.64	0.78
1:A:412:GLU:O	4:A:602:HOH:O	2.00	0.78
1:A:24:HIS:HD2	4:A:765:HOH:O	1.71	0.74
1:A:257[A]:ASP:OD1	4:A:603:HOH:O	2.06	0.73
1:A:91:LEU:C	1:A:91:LEU:HD23	2.17	0.70
1:A:33:HIS:HE1	1:A:389:GLU:OE2	1.76	0.68
1:A:275:TYR:CB	4:A:750:HOH:O	2.40	0.65
1:A:386:GLU:OE2	4:A:606:HOH:O	2.16	0.63
1:A:312:ASP:HB3	4:A:614:HOH:O	1.98	0.62
1:A:275:TYR:CD2	1:A:284:TRP:HB2	2.33	0.62
1:A:411:PHE:O	1:A:412:GLU:HB2	2.00	0.62
1:A:275:TYR:CD2	4:A:750:HOH:O	2.52	0.60
1:A:5:THR:OG1	1:A:24:HIS:HE1	1.84	0.59
1:A:177[A]:ASN:OD1	1:A:178:ALA:N	2.35	0.59
1:A:88:THR:HG22	1:A:90:GLY:H	1.67	0.58
1:A:89:ALA:N	1:A:89:ALA:C	2.61	0.58
1:A:127:ASP:HA	4:A:714:HOH:O	2.03	0.58
1:A:4:ILE:HG23	1:A:5:THR:HG23	1.88	0.56
1:A:140:ASN:HD21	1:A:154:VAL:N	1.99	0.55
1:A:275:TYR:HB3	4:A:837:HOH:O	2.06	0.54
1:A:208:HIS:HD2	4:A:957:HOH:O	1.91	0.54
1:A:25:PHE:CD1	1:A:29:LEU:HD22	2.44	0.53
1:A:308:VAL:HB	1:A:328:LYS:HE2	1.91	0.52
1:A:79[A]:SER:OG	4:A:607:HOH:O	2.18	0.52
1:A:91:LEU:HD22	1:A:125:ILE:CD1	2.39	0.51
1:A:175:ALA:HA	1:A:189:ILE:CD1	2.39	0.50
1:A:314:ASN:N	4:A:612:HOH:O	2.25	0.50
1:A:192:ILE:HG22	4:A:818:HOH:O	2.13	0.49
1:A:91:LEU:HD23	1:A:91:LEU:O	2.12	0.49
1:A:97:ASP:HA	4:A:883:HOH:O	2.13	0.47
1:A:304:LEU:HB3	1:A:305:PRO:HD3	1.96	0.47
1:A:202:TYR:CD1	1:A:286:GLY:HA3	2.51	0.46
1:A:88:THR:CG2	1:A:90:GLY:H	2.29	0.45
1:A:115:THR:CG2	4:A:605:HOH:O	2.64	0.45
1:A:6:SER:HB2	1:A:411:PHE:CE2	2.51	0.45
1:A:192:ILE:CD1	1:A:199:TYR:HB2	2.47	0.44
1:A:178:ALA:HA	1:A:179:PRO:HD2	1.78	0.44
1:A:192:ILE:CG2	4:A:818:HOH:O	2.65	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:149:ARG:NH1	4:A:628:HOH:O	2.45	0.43
1:A:100:GLU:HG2	4:A:883:HOH:O	2.18	0.43
1:A:159:LEU:HB2	1:A:301:PHE:CE2	2.54	0.43
1:A:91:LEU:CD2	1:A:99:SER:OG	2.67	0.43
1:A:54:ASN:C	1:A:54:ASN:HD22	2.26	0.42
1:A:33:HIS:CE1	1:A:389:GLU:OE2	2.65	0.42
1:A:163[B]:SER:OG	3:A:502:PO4:O3	2.37	0.42
1:A:68:GLN:NE2	4:A:639:HOH:O	2.51	0.42
1:A:136:TRP:CE2	1:A:201:LEU:HD11	2.54	0.42
1:A:91:LEU:HD21	1:A:99:SER:OG	2.19	0.42
1:A:91:LEU:C	1:A:91:LEU:CD2	2.91	0.41
1:A:285:ASN:ND2	1:A:287:GLY:H	2.18	0.41
1:A:179:PRO:HB3	1:A:275:TYR:CE1	2.55	0.41
1:A:85:LEU:C	1:A:85:LEU:HD23	2.46	0.41
1:A:91:LEU:HD21	1:A:99:SER:HA	2.02	0.41
1:A:202:TYR:CD2	1:A:202:TYR:C	2.99	0.41
1:A:52:ALA:HA	1:A:102:ILE:HD11	2.03	0.40
1:A:86:GLY:HA3	1:A:126:ILE:HD12	2.04	0.40
1:A:353:VAL:N	1:A:354:PRO:CD	2.85	0.40
1:A:268:LYS:HD2	4:A:652:HOH:O	2.21	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:877:HOH:O	4:A:894:HOH:O[2_546]	1.64	0.56

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	406/426 (95%)	390 (96%)	14 (3%)	2 (0%)	24 11

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	128	GLY
1	A	410	LYS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	328/354 (93%)	317 (97%)	11 (3%)	32 13

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	29	LEU
1	A	88	THR
1	A	91	LEU
1	A	94	LEU
1	A	115	THR
1	A	127	ASP
1	A	177[A]	ASN
1	A	177[B]	ASN
1	A	192	ILE
1	A	285	ASN
1	A	412	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (15) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	24	HIS
1	A	33	HIS
1	A	42	ASN
1	A	54	ASN
1	A	68	GLN
1	A	119	GLN
1	A	140	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	145	ASN
1	A	165	GLN
1	A	208	HIS
1	A	276	GLN
1	A	277	ASN
1	A	285	ASN
1	A	327	ASN
1	A	363	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	ADE	A	501	-	11,11,11	2.12	3 (27%)	15,15,15	2.70	9 (60%)
3	PO4	A	503	-	4,4,4	0.74	0	6,6,6	1.23	1 (16%)
3	PO4	A	502	-	4,4,4	0.94	0	6,6,6	0.91	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ADE	A	501	-	-	-	0/2/2/2

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	501	ADE	C5-C4	4.85	1.46	1.39
2	A	501	ADE	C5-C6	3.13	1.49	1.41
2	A	501	ADE	C4-N9	-2.28	1.33	1.36

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	501	ADE	N9-C4-N3	4.66	134.24	126.69
2	A	501	ADE	C5-N7-C8	3.98	107.97	103.48
2	A	501	ADE	N3-C2-N1	-3.83	122.78	128.58
2	A	501	ADE	C5-C4-N3	-3.77	119.88	126.75
2	A	501	ADE	C4-N9-C8	3.65	109.83	106.43
2	A	501	ADE	N9-C8-N7	-2.95	109.38	113.87
2	A	501	ADE	C4-C5-N7	-2.69	106.94	110.67
2	A	501	ADE	C6-C5-N7	2.55	137.00	132.09
3	A	503	PO4	O3-P-O2	2.34	115.19	107.91
2	A	501	ADE	C2-N1-C6	2.04	122.08	118.73

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	502	PO4	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	405/426 (95%)	0.18	23 (5%)	29 33	7, 17, 35, 47	6 (1%)

All (23) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	410	LYS	3.4
1	A	88	THR	3.3
1	A	411	PHE	3.0
1	A	189	ILE	2.8
1	A	95	ASP	2.7
1	A	192	ILE	2.7
1	A	97	ASP	2.6
1	A	89	ALA	2.6
1	A	312	ASP	2.5
1	A	178	ALA	2.5
1	A	152	ASN	2.4
1	A	177[A]	ASN	2.4
1	A	127	ASP	2.3
1	A	288	GLY	2.3
1	A	93	LEU	2.2
1	A	275	TYR	2.2
1	A	412	GLU	2.2
1	A	251	THR	2.1
1	A	313	PRO	2.1
1	A	4	ILE	2.1
1	A	128	GLY	2.0
1	A	180	LYS	2.0
1	A	76	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	PO4	A	503	5/5	0.75	0.14	52,57,66,68	0
3	PO4	A	502	5/5	0.83	0.12	45,49,51,53	0
2	ADE	A	501	10/10	0.87	0.12	31,34,37,38	0

6.5 Other polymers [i](#)

There are no such residues in this entry.