



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 03:33 PM UTC

PDB ID : 6U74 / pdb_00006u74
Title : BRD4-BD1 in complex with the cyclic peptide 3.1_2
Authors : Patel, K.; Walshe, J.L.; Walport, L.J.; Mackay, J.P.
Deposited on : 2019-08-31
Resolution : 1.85 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49

1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 1.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R _{free}	180053	3428 (1.86-1.86)
Clashscore	190562	3579 (1.86-1.86)
Ramachandran outliers	187476	3553 (1.86-1.86)
Sidechain outliers	187428	3553 (1.86-1.86)
RSRZ outliers	180081	3429 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	146	34% 77%10%12%
1	B	146	6% 79%8%13%
1	C	146	28% 62%10%28%
1	D	146	25% 63%9%28%
2	E	16	19% 94%6%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	F	16	 A horizontal bar chart showing the quality of chain F. The bar is divided into three segments: a red segment on the left labeled '12%', a green segment in the middle labeled '88%', and a yellow segment on the right labeled '12%'.

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 4626 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Bromodomain-containing protein 4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	128	Total	C	N	O	S	0	1	0
			1067	691	176	194	6			
1	B	127	Total	C	N	O	S	5	0	0
			1052	681	174	191	6			
1	C	105	Total	C	N	O	S	0	0	0
			889	584	143	156	6			
1	D	105	Total	C	N	O	S	0	0	0
			889	584	143	156	6			

There are 76 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	36	GLN	-	expression tag	UNP O60885
A	37	GLY	-	expression tag	UNP O60885
A	38	PRO	-	expression tag	UNP O60885
A	39	LEU	-	expression tag	UNP O60885
A	40	GLY	-	expression tag	UNP O60885
A	41	SER	-	expression tag	UNP O60885
A	169	PRO	-	expression tag	UNP O60885
A	170	GLU	-	expression tag	UNP O60885
A	171	PHE	-	expression tag	UNP O60885
A	172	PRO	-	expression tag	UNP O60885
A	173	GLY	-	expression tag	UNP O60885
A	174	ARG	-	expression tag	UNP O60885
A	175	LEU	-	expression tag	UNP O60885
A	176	GLU	-	expression tag	UNP O60885
A	177	ARG	-	expression tag	UNP O60885
A	178	PRO	-	expression tag	UNP O60885
A	179	HIS	-	expression tag	UNP O60885
A	180	ARG	-	expression tag	UNP O60885
A	181	ASP	-	expression tag	UNP O60885
B	36	GLN	-	expression tag	UNP O60885
B	37	GLY	-	expression tag	UNP O60885

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	38	PRO	-	expression tag	UNP O60885
B	39	LEU	-	expression tag	UNP O60885
B	40	GLY	-	expression tag	UNP O60885
B	41	SER	-	expression tag	UNP O60885
B	169	PRO	-	expression tag	UNP O60885
B	170	GLU	-	expression tag	UNP O60885
B	171	PHE	-	expression tag	UNP O60885
B	172	PRO	-	expression tag	UNP O60885
B	173	GLY	-	expression tag	UNP O60885
B	174	ARG	-	expression tag	UNP O60885
B	175	LEU	-	expression tag	UNP O60885
B	176	GLU	-	expression tag	UNP O60885
B	177	ARG	-	expression tag	UNP O60885
B	178	PRO	-	expression tag	UNP O60885
B	179	HIS	-	expression tag	UNP O60885
B	180	ARG	-	expression tag	UNP O60885
B	181	ASP	-	expression tag	UNP O60885
C	36	GLN	-	expression tag	UNP O60885
C	37	GLY	-	expression tag	UNP O60885
C	38	PRO	-	expression tag	UNP O60885
C	39	LEU	-	expression tag	UNP O60885
C	40	GLY	-	expression tag	UNP O60885
C	41	SER	-	expression tag	UNP O60885
C	169	PRO	-	expression tag	UNP O60885
C	170	GLU	-	expression tag	UNP O60885
C	171	PHE	-	expression tag	UNP O60885
C	172	PRO	-	expression tag	UNP O60885
C	173	GLY	-	expression tag	UNP O60885
C	174	ARG	-	expression tag	UNP O60885
C	175	LEU	-	expression tag	UNP O60885
C	176	GLU	-	expression tag	UNP O60885
C	177	ARG	-	expression tag	UNP O60885
C	178	PRO	-	expression tag	UNP O60885
C	179	HIS	-	expression tag	UNP O60885
C	180	ARG	-	expression tag	UNP O60885
C	181	ASP	-	expression tag	UNP O60885
D	36	GLN	-	expression tag	UNP O60885
D	37	GLY	-	expression tag	UNP O60885
D	38	PRO	-	expression tag	UNP O60885
D	39	LEU	-	expression tag	UNP O60885
D	40	GLY	-	expression tag	UNP O60885
D	41	SER	-	expression tag	UNP O60885

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	169	PRO	-	expression tag	UNP O60885
D	170	GLU	-	expression tag	UNP O60885
D	171	PHE	-	expression tag	UNP O60885
D	172	PRO	-	expression tag	UNP O60885
D	173	GLY	-	expression tag	UNP O60885
D	174	ARG	-	expression tag	UNP O60885
D	175	LEU	-	expression tag	UNP O60885
D	176	GLU	-	expression tag	UNP O60885
D	177	ARG	-	expression tag	UNP O60885
D	178	PRO	-	expression tag	UNP O60885
D	179	HIS	-	expression tag	UNP O60885
D	180	ARG	-	expression tag	UNP O60885
D	181	ASP	-	expression tag	UNP O60885

- Molecule 2 is a protein called cyclic peptide 3.1_2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	E	16	Total	C	N	O	S	0	0	1
			136	88	25	22	1			
2	F	16	Total	C	N	O	S	0	0	1
			136	88	25	22	1			


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	96	Total	O	0	0
			96	96		
3	B	142	Total	O	0	0
			142	142		
3	C	98	Total	O	0	0
			98	98		
3	D	84	Total	O	0	0
			84	84		
3	E	16	Total	O	0	0
			16	16		
3	F	21	Total	O	0	0
			21	21		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Bromodomain-containing protein 4




- Molecule 1: Bromodomain-containing protein 4




- Molecule 1: Bromodomain-containing protein 4





- Molecule 1: Bromodomain-containing protein 4







● Molecule 2: cyclic peptide 3.1_2



● Molecule 2: cyclic peptide 3.1_2



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	58.68Å 49.23Å 115.94Å 90.00° 102.58° 90.00°	Depositor
Resolution (Å)	45.51 – 1.85 45.51 – 1.85	Depositor EDS
% Data completeness (in resolution range)	98.3 (45.51-1.85) 98.3 (45.51-1.85)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.17 (at 1.86Å)	Xtriage
Refinement program	PHENIX 1.14 _3260	Depositor
R, R_{free}	0.284 , 0.323 0.285 , 0.324	Depositor DCC
R_{free} test set	2813 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å ²)	10.3	Xtriage
Anisotropy	0.295	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 44.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.000 for h,-k,-h-l	Xtriage
F_o, F_c correlation	0.88	EDS
Total number of atoms	4626	wwPDB-VP
Average B, all atoms (Å ²)	27.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 50.39 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 6.5162e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ACE, NH2, ALY

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.10	0/1097	0.27	0/1494
1	B	0.12	0/1082	0.30	0/1473
1	C	0.09	0/913	0.25	0/1239
1	D	0.10	0/913	0.27	0/1239
2	E	0.12	0/97	0.27	0/127
2	F	0.15	0/97	0.34	0/127
All	All	0.10	0/4199	0.28	0/5699

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1067	0	1067	8	0
1	B	1052	0	1050	5	0
1	C	889	0	891	7	0
1	D	889	0	891	7	0
2	E	136	0	136	1	0
2	F	136	0	136	2	0
3	A	96	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	142	0	0	0	0
3	C	98	0	0	0	0
3	D	84	0	0	0	0
3	E	16	0	0	0	0
3	F	21	0	0	0	0
All	All	4626	0	4171	27	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (27) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:73:THR:HG22	1:A:156:LEU:HD21	1.79	0.64
1:D:82:PRO:HB3	2:F:5:ALY:HG2	1.90	0.53
1:A:68:ARG:O	1:A:72:LYS:HD3	2.10	0.52
1:C:79:PHE:HB3	1:C:149:MET:HB3	1.92	0.52
1:B:79:PHE:HB3	1:B:149:MET:HB3	1.95	0.48
1:D:145:ASP:OD1	1:D:146:ILE:N	2.46	0.48
1:A:97:TYR:HA	1:A:139:TYR:CE1	2.49	0.48
1:B:73:THR:HG22	1:B:156:LEU:HD21	1.96	0.47
1:C:87:VAL:O	1:C:98:TYR:OH	2.30	0.46
1:C:145:ASP:OD1	1:C:146:ILE:N	2.49	0.46
1:A:79:PHE:HB3	1:A:149:MET:HB3	1.97	0.46
1:D:80:ALA:HA	1:D:153:LEU:HD21	1.98	0.46
1:D:164:LEU:HD12	1:D:165:PRO:HD2	1.98	0.45
1:B:74:LEU:HD21	1:B:157:PHE:HB2	1.99	0.45
1:D:81:TRP:CD2	2:F:7:ALY:HE2	2.52	0.45
1:A:141:LYS:NZ	1:A:141:LYS:HB3	2.32	0.44
1:D:79:PHE:HB3	1:D:149:MET:HB3	2.00	0.44
1:B:81:TRP:CG	1:B:82:PRO:HD3	2.52	0.44
1:A:81:TRP:CG	1:A:82:PRO:HD3	2.52	0.44
1:C:164:LEU:HD12	1:C:165:PRO:HD2	2.00	0.43
1:A:158:LEU:HD23	1:A:161:ILE:HD11	2.01	0.42
1:A:94:LEU:HD11	2:E:12:ALY:HG3	2.02	0.42
1:C:66:LEU:HD23	1:C:70:VAL:HG21	2.02	0.42
1:B:158:LEU:HD23	1:B:161:ILE:HD11	2.03	0.41
1:C:81:TRP:CG	1:C:82:PRO:HD3	2.56	0.41
1:D:74:LEU:HD11	1:D:157:PHE:HB2	2.03	0.41
1:C:74:LEU:HD11	1:C:157:PHE:HB2	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	127/146 (87%)	125 (98%)	2 (2%)	0	100	100
1	B	125/146 (86%)	123 (98%)	2 (2%)	0	100	100
1	C	103/146 (70%)	103 (100%)	0	0	100	100
1	D	103/146 (70%)	103 (100%)	0	0	100	100
2	E	11/16 (69%)	10 (91%)	1 (9%)	0	100	100
2	F	11/16 (69%)	10 (91%)	1 (9%)	0	100	100
All	All	480/616 (78%)	474 (99%)	6 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	121/136 (89%)	120 (99%)	1 (1%)	73	67
1	B	119/136 (88%)	118 (99%)	1 (1%)	73	67
1	C	98/136 (72%)	98 (100%)	0	100	100
1	D	98/136 (72%)	97 (99%)	1 (1%)	68	60
2	E	10/10 (100%)	10 (100%)	0	100	100
2	F	10/10 (100%)	10 (100%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
All	All	456/564 (81%)	453 (99%)	3 (1%)	76	70

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	39	LEU
1	B	162	ASN
1	D	103	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	93	ASN
1	A	127	GLN
1	B	62	GLN
1	B	93	ASN
1	C	78	GLN
1	C	116	ASN
1	D	62	GLN
1	D	64	GLN
1	D	78	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

6 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	ALY	F	12	2	10,11,12	0.88	0	7,12,14	0.57	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	ALY	E	12	2	10,11,12	0.86	0	7,12,14	0.52	0
2	ALY	F	7	2	10,11,12	0.86	0	7,12,14	0.68	0
2	ALY	E	5	2	10,11,12	0.85	0	7,12,14	0.63	0
2	ALY	E	7	2	10,11,12	0.87	0	7,12,14	0.79	0
2	ALY	F	5	2	10,11,12	0.84	0	7,12,14	0.63	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ALY	F	12	2	-	0/9/10/12	-
2	ALY	E	12	2	-	0/9/10/12	-
2	ALY	F	7	2	-	4/9/10/12	-
2	ALY	E	5	2	-	0/9/10/12	-
2	ALY	E	7	2	-	2/9/10/12	-
2	ALY	F	5	2	-	0/9/10/12	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	F	7	ALY	N-CA-CB-CG
2	F	7	ALY	C-CA-CB-CG
2	E	7	ALY	CG-CD-CE-NZ
2	F	7	ALY	CG-CD-CE-NZ
2	F	7	ALY	CE-CD-CG-CB
2	E	7	ALY	CE-CD-CG-CB

There are no ring outliers.

3 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	E	12	ALY	1	0
2	F	7	ALY	1	0
2	F	5	ALY	1	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	128/146 (87%)	1.92	50 (39%) 1 0	12, 30, 46, 71	1 (0%)
1	B	127/146 (86%)	0.52	9 (7%) 22 24	5, 15, 29, 42	1 (0%)
1	C	105/146 (71%)	1.99	41 (39%) 1 0	20, 32, 58, 71	0
1	D	105/146 (71%)	1.56	37 (35%) 1 1	11, 26, 52, 64	0
2	E	11/16 (68%)	1.58	3 (27%) 1 1	23, 26, 38, 39	0
2	F	11/16 (68%)	0.84	2 (18%) 3 3	13, 16, 26, 31	0
All	All	487/616 (79%)	1.46	142 (29%) 1 1	5, 26, 46, 71	2 (0%)

All (142) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	164	LEU	7.1
1	C	165	PRO	6.9
1	C	162	ASN	5.7
1	C	163	GLU	5.1
1	C	68	ARG	5.1
1	D	161	ILE	4.9
1	A	41	SER	4.8
1	A	42	SER	4.8
1	A	39	LEU	4.7
1	C	64	GLN	4.6
1	D	165	PRO	4.6
2	E	3	THR	4.5
1	D	68	ARG	4.4
1	B	166	THR	4.3
1	D	69	VAL	4.3
1	C	161	ILE	4.3
1	C	98	TYR	4.2
1	A	40	GLY	4.2
1	C	61	ASN	4.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	164	LEU	4.1
1	C	72	LYS	4.0
1	A	64	GLN	4.0
1	D	61	ASN	4.0
1	C	90	VAL	4.0
1	D	65	TYR	3.9
1	D	63	LEU	3.8
1	A	68	ARG	3.8
1	D	163	GLU	3.8
1	A	57	LYS	3.8
1	D	116	ASN	3.7
1	A	59	GLN	3.7
1	D	164	LEU	3.7
1	A	156	LEU	3.6
1	A	85	GLN	3.6
1	D	64	GLN	3.5
1	B	41	SER	3.5
1	D	162	ASN	3.5
1	D	158	LEU	3.5
1	D	159	GLN	3.4
1	A	72	LYS	3.4
1	C	159	GLN	3.4
1	D	95	PRO	3.4
1	D	153	LEU	3.4
1	D	120	TRP	3.3
1	A	96	ASP	3.3
1	C	69	VAL	3.3
1	D	67	LEU	3.3
1	A	165	PRO	3.2
1	A	67	LEU	3.2
1	D	62	GLN	3.2
1	A	166	THR	3.1
1	C	122	ALA	3.1
1	A	161	ILE	3.0
1	C	65	TYR	3.0
1	D	72	LYS	3.0
2	F	6	GLY	3.0
1	C	157	PHE	3.0
1	D	112	LYS	3.0
1	A	117	ASN	3.0
1	A	61	ASN	3.0
1	B	85	GLN	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	96	ASP	2.9
2	E	6	GLY	2.9
1	A	89	ALA	2.9
1	D	70	VAL	2.9
1	A	95	PRO	2.9
1	B	42	SER	2.9
1	C	66	LEU	2.9
1	A	65	TYR	2.8
2	F	3	THR	2.8
1	D	155	LYS	2.8
1	C	116	ASN	2.8
1	C	74	LEU	2.8
1	D	160	LYS	2.8
1	C	123	GLN	2.8
1	C	160	LYS	2.8
1	D	118	TYR	2.7
1	C	79	PHE	2.7
1	B	40	GLY	2.7
1	D	74	LEU	2.7
1	C	117	ASN	2.7
1	D	117	ASN	2.7
1	C	155	LYS	2.7
1	A	53	PRO	2.7
1	A	92	LEU	2.7
1	A	127	GLN	2.6
1	A	142	PRO	2.6
1	C	95	PRO	2.6
1	A	73	THR	2.6
1	C	103	THR	2.6
1	D	66	LEU	2.6
1	A	100	ILE	2.6
1	C	127	GLN	2.6
1	A	99	LYS	2.6
1	A	91	LYS	2.5
1	D	127	GLN	2.5
1	A	63	LEU	2.5
1	C	63	LEU	2.5
1	D	111	LYS	2.5
1	C	118	TYR	2.4
1	A	112	LYS	2.4
1	A	134[A]	THR	2.4
1	B	91	LYS	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	73	THR	2.4
1	A	75	TRP	2.4
1	C	93	ASN	2.4
1	C	153	LEU	2.4
1	D	98	TYR	2.4
1	C	112	LYS	2.4
1	A	70	VAL	2.4
1	C	81	TRP	2.4
1	D	102	LYS	2.4
1	A	79	PHE	2.4
1	A	90	VAL	2.4
1	A	81	TRP	2.4
1	C	67	LEU	2.3
1	A	69	VAL	2.3
1	A	93	ASN	2.3
1	A	111	LYS	2.3
1	A	157	PHE	2.3
1	D	157	PHE	2.3
1	B	59	GLN	2.3
1	D	73	THR	2.3
1	A	129	PHE	2.2
1	A	158	LEU	2.2
1	C	126	ILE	2.2
1	A	163	GLU	2.2
1	A	103	THR	2.2
1	B	93	ASN	2.1
1	B	90	VAL	2.1
1	C	71	LEU	2.1
1	D	122	ALA	2.1
1	A	97	TYR	2.1
1	C	94	LEU	2.1
1	D	96	ASP	2.1
2	E	4	ILE	2.1
1	D	114	LEU	2.1
1	C	138	ILE	2.0
1	C	92	LEU	2.0
1	A	101	ILE	2.0
1	A	60	THR	2.0
1	A	123	GLN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	ALY	E	7	12/13	0.80	0.21	26,32,48,59	0
2	ALY	E	5	12/13	0.81	0.17	29,31,41,47	0
2	ALY	F	7	12/13	0.84	0.20	18,24,55,56	0
2	ALY	E	12	12/13	0.89	0.12	20,22,26,27	0
2	ALY	F	5	12/13	0.90	0.11	12,20,31,33	0
2	ALY	F	12	12/13	0.92	0.09	11,11,19,22	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.