



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 02:08 PM UTC

PDB ID : 6U7C / pdb_00006u7c
Title : Human GRK2 in complex with Gbetagamma subunits and CCG258747
Authors : Bouley, R.; Tesmer, J.J.G.
Deposited on : 2019-09-02
Resolution : 2.44 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.44 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2340 (2.46-2.42)
Clashscore	190562	2400 (2.46-2.42)
Ramachandran outliers	187476	2379 (2.46-2.42)
Sidechain outliers	187428	2379 (2.46-2.42)
RSRZ outliers	180081	2340 (2.46-2.42)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	689	9% 76% 15% 8%
2	B	340	0% 89% 10%
3	G	71	7% 79% 18%

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 8327 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Beta-adrenergic receptor kinase 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	631	Total	C	N	O	S	0	0	0
			5159	3288	903	933	35			

- Molecule 2 is a protein called Guanine nucleotide-binding protein G(I)/G(S)/G(T) subunit beta-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	339	Total	C	N	O	S	0	0	0
			2607	1607	468	511	21			


- Molecule 3 is a protein called Guanine nucleotide-binding protein G(I)/G(S)/G(O) subunit gamma-2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	G	58	Total	C	N	O	S	0	0	0
			447	280	79	85	3			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
G	68	SER	CYS	engineered mutation	UNP P63212

- Molecule 4 is 5-[(3S,4R)-3-{[(2H-1,3-benzodioxol-5-yl)oxy]methyl}piperidin-4-yl]-2-fluoro-N-[(2H-indazol-3-yl)methyl]benzamide (CCD ID: Q1Y) (formula: C₂₈H₂₇FN₄O₄) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	F	N	O	
			37	28	1	4	4	0

- Molecule 5 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	Mg		
			1	1	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	40	Total	O		
			40	40	0	0
6	B	33	Total	O		
			33	33	0	0
6	G	3	Total	O		
			3	3	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Beta-adrenergic receptor kinase 1




- Molecule 2: Guanine nucleotide-binding protein G(I)/G(S)/G(T) subunit beta-1



- Molecule 3: Guanine nucleotide-binding protein G(I)/G(S)/G(O) subunit gamma-2





4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	60.63Å 240.61Å 214.58Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.90 – 2.44 29.90 – 2.44	Depositor EDS
% Data completeness (in resolution range)	97.0 (29.90-2.44) 79.5 (29.90-2.44)	Depositor EDS
R_{merge}	0.24	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.27 (at 2.45Å)	Xtriage
Refinement program	PHENIX 1.13_2998	Depositor
R, R_{free}	0.237 , 0.262 0.233 , 0.259	Depositor DCC
R_{free} test set	1999 reflections (3.39%)	wwPDB-VP
Wilson B-factor (Å ²)	34.7	Xtriage
Anisotropy	0.571	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 55.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	8327	wwPDB-VP
Average B, all atoms (Å ²)	93.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.11% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: Q1Y, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.16	0/5274	0.37	1/7086 (0.0%)
2	B	0.16	0/2654	0.40	0/3597
3	G	0.13	0/453	0.30	0/610
All	All	0.16	0/8381	0.38	1/11293 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	369	ASP	CB-CA-C	-5.03	109.20	116.54

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	476	GLU	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5159	0	5152	59	0
2	B	2607	0	2510	23	0
3	G	447	0	460	1	0
4	A	37	0	0	0	0
5	A	1	0	0	0	0
6	A	40	0	0	3	0
6	B	33	0	0	0	0
6	G	3	0	0	0	0
All	All	8327	0	8122	81	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (81) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:605:GLU:HA	1:A:625:ARG:HD3	1.62	0.79
2:B:45:MET:HE3	2:B:308:LEU:HD21	1.70	0.74
1:A:169:ASP:OD1	1:A:169:ASP:N	2.17	0.68
1:A:442:ARG:HB2	1:A:446:GLU:HG3	1.75	0.67
1:A:315:TYR:OH	1:A:334:SER:O	2.06	0.66
1:A:644:LYS:NZ	6:A:803:HOH:O	2.31	0.62
1:A:173:ARG:NH2	1:A:534:ASP:OD1	2.33	0.62
1:A:587:ARG:HD2	1:A:600:LEU:HD22	1.81	0.61
1:A:664:MET:HE3	2:B:101:MET:HG3	1.83	0.60
2:B:51:LEU:HB2	2:B:336:LEU:HB2	1.85	0.58
2:B:215:GLU:HB3	2:B:217:MET:HE3	1.84	0.58
2:B:146:LEU:HD11	2:B:159:THR:HB	1.85	0.57
1:A:71:PHE:HB2	1:A:170:LYS:HG3	1.87	0.57
1:A:565:MET:HE3	1:A:634:CYS:SG	2.45	0.56
1:A:346:LYS:HE3	1:A:367:ALA:HB1	1.89	0.55
2:B:79:LEU:HG	2:B:95:LEU:HD21	1.89	0.55
1:A:358:ALA:HB3	1:A:361:VAL:HG23	1.89	0.55
1:A:377:LEU:O	1:A:381:LEU:HG	2.09	0.52
1:A:324:LEU:O	1:A:331:VAL:HA	2.09	0.52
1:A:565:MET:HE2	1:A:632:LEU:HB3	1.92	0.52
1:A:609:VAL:HG22	1:A:622:LEU:HD22	1.90	0.51
1:A:417:PHE:O	1:A:422:ARG:NH2	2.43	0.51
1:A:414:PRO:O	1:A:422:ARG:NH1	2.43	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:225:LYS:NZ	1:A:496:ILE:O	2.36	0.50
1:A:410:ALA:HB2	1:A:431:ARG:HH22	1.77	0.50
1:A:357:MET:HE3	1:A:362:LEU:HD21	1.93	0.49
1:A:239:GLU:HB2	1:A:337:GLY:HA2	1.95	0.48
1:A:220:LYS:HE2	1:A:222:LEU:HD21	1.94	0.48
2:B:166:CYS:HB2	2:B:180:PHE:HB2	1.94	0.48
2:B:235:PHE:CG	2:B:236:PRO:HD2	2.48	0.48
2:B:283:ARG:HB3	3:G:51:LEU:HD11	1.95	0.48
2:B:30:LEU:HD23	2:B:262:MET:HE2	1.96	0.48
1:A:35:PRO:O	1:A:176:GLN:NE2	2.34	0.47
2:B:286:LEU:HD22	2:B:327:VAL:HG21	1.95	0.47
1:A:39:ILE:HG13	1:A:43:MET:HG2	1.97	0.47
1:A:318:LEU:HG	1:A:380:MET:HE3	1.97	0.46
1:A:279:LEU:HD23	1:A:279:LEU:HA	1.80	0.46
1:A:347:PRO:HD2	1:A:368:TYR:O	2.15	0.46
1:A:314:VAL:HG12	1:A:370:SER:HA	1.97	0.46
2:B:327:VAL:HG13	2:B:339:TRP:HB2	1.97	0.46
1:A:413:LEU:HD12	1:A:422:ARG:HG2	1.98	0.45
1:A:361:VAL:HA	1:A:368:TYR:CE2	2.52	0.45
1:A:398:ASP:HB3	1:A:401:GLU:HB2	1.97	0.45
1:A:316:ARG:CZ	1:A:351:VAL:HG11	2.47	0.44
1:A:620:LEU:HD21	1:A:644:LYS:HB2	2.00	0.44
1:A:294:MET:HE3	1:A:452:PHE:CZ	2.53	0.44
1:A:614:ILE:HD11	1:A:631:ILE:HD13	1.99	0.44
2:B:250:CYS:HB2	2:B:264:TYR:HB2	1.98	0.44
1:A:178:LYS:NZ	6:A:809:HOH:O	2.51	0.44
1:A:319:LYS:HE3	1:A:321:ALA:HB3	2.00	0.44
2:B:139:LEU:HB3	2:B:169:TRP:CE3	2.53	0.44
2:B:211:TRP:CZ3	2:B:218:CYS:HB2	2.53	0.43
2:B:235:PHE:CD1	2:B:236:PRO:HD2	2.54	0.43
1:A:603:MET:HB3	1:A:651:TYR:HA	2.00	0.43
1:A:411:VAL:HG22	1:A:413:LEU:HD23	2.01	0.42
2:B:180:PHE:CE1	2:B:216:GLY:HA2	2.54	0.42
1:A:162:PHE:O	1:A:166:ILE:HG12	2.19	0.42
1:A:314:VAL:HG13	1:A:342:PHE:CE2	2.53	0.42
2:B:262:MET:SD	2:B:302:ALA:HB2	2.59	0.42
1:A:90:LYS:HA	1:A:90:LYS:HD3	1.85	0.42
1:A:410:ALA:HB2	1:A:431:ARG:NH2	2.34	0.42
2:B:124:TYR:CE2	2:B:135:VAL:HG22	2.54	0.42
1:A:260:ALA:O	1:A:510:PRO:HA	2.19	0.42
2:B:65:THR:OG1	2:B:107:PRO:O	2.30	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:392:ARG:HH11	1:A:395:LYS:HA	1.84	0.42
2:B:61:MET:HE2	2:B:61:MET:HB2	1.90	0.42
2:B:112:VAL:HG23	2:B:124:TYR:HB2	2.00	0.42
1:A:354:HIS:HA	1:A:357:MET:HE2	2.02	0.41
1:A:359:PRO:HD2	1:A:436:ARG:HH22	1.85	0.41
1:A:121:SER:O	1:A:123:PRO:HD3	2.20	0.41
1:A:604:GLU:HG3	6:A:827:HOH:O	2.20	0.41
1:A:217:TYR:OH	1:A:520:GLU:OE1	2.38	0.41
1:A:496:ILE:HG22	1:A:497:LYS:N	2.35	0.41
1:A:581:PHE:CD2	1:A:590:TRP:HB3	2.55	0.41
1:A:648:ARG:HA	1:A:651:TYR:CD2	2.55	0.41
1:A:246:VAL:HG21	1:A:336:LEU:HD11	2.02	0.41
1:A:581:PHE:HB3	1:A:588:LEU:HD11	2.03	0.41
1:A:32:ILE:HG13	1:A:212:ASP:HB2	2.04	0.40
1:A:420:GLU:H	1:A:420:GLU:HG2	1.58	0.40
1:A:426:GLU:O	1:A:430:GLN:HG2	2.21	0.40
2:B:124:TYR:CZ	2:B:135:VAL:HG22	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	625/689 (91%)	607 (97%)	18 (3%)	0	100	100
2	B	337/340 (99%)	322 (96%)	15 (4%)	0	100	100
3	G	56/71 (79%)	55 (98%)	1 (2%)	0	100	100
All	All	1018/1100 (92%)	984 (97%)	34 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	562/609 (92%)	548 (98%)	14 (2%)	42	54
2	B	282/283 (100%)	282 (100%)	0	100	100
3	G	47/58 (81%)	46 (98%)	1 (2%)	47	60
All	All	891/950 (94%)	876 (98%)	15 (2%)	53	66

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	169	ASP
1	A	228	LYS
1	A	294	MET
1	A	350	SER
1	A	351	VAL
1	A	409	MET
1	A	411	VAL
1	A	412	GLU
1	A	420	GLU
1	A	477	VAL
1	A	499	LEU
1	A	546	GLN
1	A	612	THR
1	A	625	ARG
3	G	54	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	135	HIS
1	A	546	GLN
1	A	598	GLN
1	A	633	GLN
2	B	13	GLN
2	B	156	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 2 ligands modelled in this entry, 1 is monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	Q1Y	A	701	-	39,42,42	1.92	9 (23%)	47,59,59	1.56	10 (21%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	Q1Y	A	701	-	-	5/18/35/35	0/6/6/6

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	701	Q1Y	C25-N26	5.55	1.46	1.33
4	A	701	Q1Y	C05-C06	4.60	1.58	1.51
4	A	701	Q1Y	C28-N36	-4.01	1.31	1.35

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	701	Q1Y	C34-N35	3.69	1.38	1.33
4	A	701	Q1Y	C22-C06	-2.94	1.50	1.53
4	A	701	Q1Y	C34-C29	-2.89	1.37	1.43
4	A	701	Q1Y	O14-C13	2.28	1.41	1.38
4	A	701	Q1Y	O16-C17	2.14	1.41	1.38
4	A	701	Q1Y	C08-C07	2.02	1.55	1.51

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	701	Q1Y	C22-C06-C07	3.51	114.18	109.75
4	A	701	Q1Y	C05-C06-C07	-3.22	107.45	113.67
4	A	701	Q1Y	O16-C17-C18	3.06	131.93	127.86
4	A	701	Q1Y	C23-C24-C02	2.85	120.37	116.83
4	A	701	Q1Y	C19-C07-C06	2.72	114.04	109.88
4	A	701	Q1Y	C03-C02-C24	-2.60	120.26	123.07
4	A	701	Q1Y	O14-C13-C12	2.37	132.29	127.72
4	A	701	Q1Y	C08-C07-C06	-2.11	108.32	112.09
4	A	701	Q1Y	C22-C06-C05	-2.09	108.69	112.54
4	A	701	Q1Y	C02-C24-C25	-2.01	118.86	124.32

There are no chirality outliers.

All (5) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	A	701	Q1Y	N26-C27-C28-C29
4	A	701	Q1Y	C06-C07-C08-O09
4	A	701	Q1Y	C19-C07-C08-O09
4	A	701	Q1Y	C11-C10-O09-C08
4	A	701	Q1Y	C18-C10-O09-C08

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the

average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	631/689 (91%)	0.89	65 (10%) 12 9	49, 96, 157, 200	0
2	B	339/340 (99%)	0.26	4 (1%) 76 78	53, 73, 107, 169	0
3	G	58/71 (81%)	0.61	5 (8%) 16 13	62, 86, 150, 172	0
All	All	1028/1100 (93%)	0.67	74 (7%) 21 18	49, 87, 150, 200	0

All (74) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	496	ILE	4.9
1	A	495	GLY	4.9
3	G	7	ALA	4.2
1	A	483	PHE	4.0
1	A	342	PHE	3.9
1	A	548	GLY	3.7
2	B	127	LYS	3.6
1	A	359	PRO	3.5
1	A	547	LEU	3.5
1	A	367	ALA	3.4
1	A	362	LEU	3.4
1	A	391	PHE	3.4
1	A	572	PHE	3.4
1	A	411	VAL	3.4
1	A	477	VAL	3.4
1	A	354	HIS	3.2
1	A	398	ASP	3.1
1	A	408	THR	3.1
1	A	573	LEU	3.0
1	A	53	VAL	3.0
1	A	221	CYS	3.0
1	A	137	GLY	3.0
1	A	499	LEU	3.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	542	ALA	2.8
1	A	361	VAL	2.8
1	A	433	VAL	2.8
1	A	396	THR	2.7
1	A	335	ASP	2.7
1	A	394	HIS	2.7
1	A	407	LEU	2.7
1	A	121	SER	2.7
1	A	596	ALA	2.6
1	A	368	TYR	2.6
1	A	154	CYS	2.5
1	A	371	SER	2.5
3	G	52	THR	2.5
1	A	410	ALA	2.5
1	A	400	HIS	2.4
1	A	628	LYS	2.4
1	A	493	THR	2.4
1	A	571	PRO	2.4
1	A	392	ARG	2.4
1	A	393	GLN	2.4
1	A	409	MET	2.4
1	A	545	LYS	2.4
2	B	47	THR	2.3
1	A	348	HIS	2.3
1	A	124	PHE	2.3
1	A	370	SER	2.3
1	A	75	HIS	2.3
1	A	402	ILE	2.3
1	A	76	LEU	2.3
1	A	669	ARG	2.3
1	A	485	ILE	2.3
3	G	64	LYS	2.3
1	A	350	SER	2.2
3	G	54	VAL	2.2
1	A	30	LYS	2.2
1	A	235	LEU	2.2
1	A	404	ARG	2.2
1	A	146	PHE	2.2
3	G	53	PRO	2.2
1	A	106	ARG	2.2
1	A	138	LYS	2.1
1	A	546	GLN	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	130	GLU	2.1
1	A	479	ALA	2.1
2	B	271	CYS	2.1
1	A	412	GLU	2.1
1	A	444	ALA	2.0
2	B	131	GLY	2.0
1	A	439	CYS	2.0
1	A	413	LEU	2.0
1	A	357	MET	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	MG	A	702	1/1	0.87	0.08	154,154,154,154	0
4	Q1Y	A	701	37/37	0.94	0.11	52,81,99,101	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.