



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 25, 2026 – 02:27 AM EDT

PDB ID : 1U8X / pdb_00001u8x
Title : CRYSTAL STRUCTURE OF GLVA FROM BACILLUS SUBTILIS, A METAL-REQUIRING, NAD-DEPENDENT 6-PHOSPHO-ALPHA-GLUCOSIDASE
Authors : Rajan, S.S.; Yang, X.; Collart, F.; Anderson, W.F.; Midwest Center for Structural Genomics (MCSG)
Deposited on : 2004-08-09
Resolution : 2.05 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.05 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2260 (2.04-2.04)
Clashscore	190562	2333 (2.04-2.04)
Ramachandran outliers	187476	2318 (2.04-2.04)
Sidechain outliers	187428	2318 (2.04-2.04)
RSRZ outliers	180081	2260 (2.04-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	X	472	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	G6P	X	902	X	-	-	-

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 3672 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Maltose-6'-phosphate glucosidase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
1	X	436	3449	2188	581	659	5	16	0	0	0

There are 40 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
X	-22	MSE	-	expression tag	UNP P54716
X	-21	HIS	-	expression tag	UNP P54716
X	-20	HIS	-	expression tag	UNP P54716
X	-19	HIS	-	expression tag	UNP P54716
X	-18	HIS	-	expression tag	UNP P54716
X	-17	HIS	-	expression tag	UNP P54716
X	-16	HIS	-	expression tag	UNP P54716
X	-15	SER	-	cloning artifact	UNP P54716
X	-14	SER	-	cloning artifact	UNP P54716
X	-13	GLY	-	cloning artifact	UNP P54716
X	-12	VAL	-	cloning artifact	UNP P54716
X	-11	ASP	-	cloning artifact	UNP P54716
X	-10	LEU	-	cloning artifact	UNP P54716
X	-9	GLY	-	cloning artifact	UNP P54716
X	-8	THR	-	cloning artifact	UNP P54716
X	-7	GLU	-	cloning artifact	UNP P54716
X	-6	ASN	-	cloning artifact	UNP P54716
X	-5	LEU	-	cloning artifact	UNP P54716
X	-4	TYR	-	cloning artifact	UNP P54716
X	-3	PHE	-	cloning artifact	UNP P54716
X	-2	GLN	-	cloning artifact	UNP P54716
X	-1	SER	-	cloning artifact	UNP P54716
X	0	ASN	-	cloning artifact	UNP P54716
X	1	MSE	MET	modified residue	UNP P54716
X	24	MSE	MET	modified residue	UNP P54716
X	84	MSE	MET	modified residue	UNP P54716
X	94	MSE	MET	modified residue	UNP P54716

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
X	122	MSE	MET	modified residue	UNP P54716
X	135	MSE	MET	modified residue	UNP P54716
X	144	MSE	MET	modified residue	UNP P54716
X	173	MSE	MET	modified residue	UNP P54716
X	181	MSE	MET	modified residue	UNP P54716
X	193	MSE	MET	modified residue	UNP P54716
X	218	MSE	MET	modified residue	UNP P54716
X	275	MSE	MET	modified residue	UNP P54716
X	290	MSE	MET	modified residue	UNP P54716
X	303	MSE	MET	modified residue	UNP P54716
X	339	MSE	MET	modified residue	UNP P54716
X	357	MSE	MET	modified residue	UNP P54716
X	385	MSE	MET	modified residue	UNP P54716

- Molecule 2 is 6-O-phosphono-alpha-D-glucopyranose (CCD ID: G6P) (formula: $C_6H_{13}O_9P$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	X	1	Total	C	O	P	0	0
			16	6	9	1		

- Molecule 3 is MANGANESE (II) ION (CCD ID: MN) (formula: Mn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	X	1	Total	Mn	0	0
			1	1		

- Molecule 4 is NICOTINAMIDE-ADENINE-DINUCLEOTIDE (CCD ID: NAD) (formula: $\text{C}_{21}\text{H}_{27}\text{N}_7\text{O}_{14}\text{P}_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	X	1	Total 44	C 21	N 7	O 14	P 2	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	X	162	Total O 162 162	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Maltose-6'-phosphate glucosidase



4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	82.60Å 102.02Å 144.75Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.02 – 2.05 30.02 – 2.05	Depositor EDS
% Data completeness (in resolution range)	95.0 (30.02-2.05) 95.0 (30.02-2.05)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.51 (at 2.05Å)	Xtriage
Refinement program	REFMAC 5.1.24	Depositor
R, R_{free}	0.195 , 0.221 0.208 , 0.216	Depositor DCC
R_{free} test set	3834 reflections (9.95%)	wwPDB-VP
Wilson B-factor (Å ²)	33.4	Xtriage
Anisotropy	0.541	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 38.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	3672	wwPDB-VP
Average B, all atoms (Å ²)	31.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.45% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAD, MN, G6P

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	X	0.75	1/3507 (0.0%)	0.90	0/4726

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	X	0	2

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	X	24	MSE	SE-CE	-5.14	1.80	1.95

There are no bond angle outliers.

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	X	298	PHE	Peptide
1	X	311	GLU	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	X	3449	0	3397	62	0
2	X	16	0	9	0	0
3	X	1	0	0	0	0
4	X	44	0	26	7	0
5	X	162	0	0	5	1
All	All	3672	0	3432	62	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (62) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:X:24:MSE:HE3	1:X:28:HIS:ND1	1.75	1.02
1:X:3:LYS:N	1:X:4:LYS:HA	2.01	0.75
1:X:24:MSE:CE	1:X:28:HIS:ND1	2.49	0.75
1:X:172:ASP:HB3	1:X:323:TYR:CE2	2.22	0.75
1:X:177:ILE:HG22	1:X:181:MSE:HE3	1.71	0.71
1:X:244:ASN:O	5:X:1044:HOH:O	2.08	0.71
1:X:147:TYR:O	4:X:900:NAD:H2N	1.91	0.70
1:X:24:MSE:HE1	1:X:325:VAL:HG21	1.74	0.69
1:X:359:GLU:OE2	5:X:937:HOH:O	2.14	0.66
1:X:298:PHE:O	1:X:298:PHE:CG	2.50	0.65
1:X:24:MSE:HE1	1:X:325:VAL:CG2	2.28	0.63
1:X:15:SER:HB3	1:X:18:THR:HG22	1.82	0.62
1:X:86:HIS:HB3	1:X:147:TYR:CE1	2.33	0.62
1:X:293:ARG:O	1:X:294:GLU:C	2.43	0.62
1:X:296:PHE:O	1:X:300:GLN:HG2	2.02	0.59
1:X:146:ASN:HD21	1:X:148:SER:HB2	1.68	0.59
1:X:321:ALA:O	1:X:324:ILE:HG12	2.03	0.59
1:X:86:HIS:HB3	1:X:147:TYR:CD1	2.41	0.56
1:X:218:MSE:HE3	1:X:222:LYS:NZ	2.21	0.55
1:X:296:PHE:CD2	1:X:297:ILE:HG23	2.42	0.54
1:X:17:PHE:O	1:X:21:ILE:HG23	2.07	0.53
1:X:321:ALA:HB1	1:X:324:ILE:HD11	1.91	0.53
1:X:85:ALA:O	1:X:147:TYR:CD1	2.62	0.53
1:X:24:MSE:HE2	1:X:24:MSE:O	2.09	0.52
1:X:311:GLU:HB2	5:X:1029:HOH:O	2.08	0.52
1:X:158:THR:HG21	1:X:167:ILE:HD13	1.92	0.51
1:X:26:LEU:HD21	1:X:58:ILE:HD13	1.92	0.51
1:X:15:SER:HB2	4:X:900:NAD:H4B	1.93	0.50
1:X:24:MSE:HE2	1:X:25:LEU:HA	1.93	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:X:173:MSE:HB3	1:X:174:PRO:HD3	1.95	0.49
1:X:14:GLY:C	4:X:900:NAD:O3B	2.56	0.48
1:X:22:VAL:O	1:X:26:LEU:HD22	2.13	0.48
1:X:132:LEU:C	1:X:132:LEU:HD23	2.39	0.48
1:X:148:SER:C	5:X:1058:HOH:O	2.55	0.48
1:X:147:TYR:CE2	1:X:324:ILE:HD12	2.47	0.48
1:X:171:CYS:HA	4:X:900:NAD:O7N	2.15	0.47
1:X:24:MSE:HE2	1:X:24:MSE:C	2.40	0.46
1:X:16:THR:HB	1:X:297:ILE:HD11	1.98	0.46
1:X:341:LEU:HD12	1:X:369:PRO:HB3	1.97	0.46
1:X:300:GLN:HE21	1:X:303:MSE:HE2	1.81	0.46
1:X:297:ILE:C	1:X:299:SER:H	2.25	0.45
1:X:324:ILE:H	1:X:324:ILE:HD13	1.82	0.45
1:X:311:GLU:O	1:X:312:ASN:HB2	2.17	0.44
1:X:14:GLY:C	4:X:900:NAD:HO3A	2.25	0.44
1:X:181:MSE:HE1	1:X:206:TRP:CD2	2.53	0.44
1:X:13:GLY:O	1:X:18:THR:HG21	2.18	0.43
1:X:177:ILE:CG2	1:X:181:MSE:HE3	2.44	0.43
1:X:297:ILE:C	1:X:299:SER:N	2.76	0.43
1:X:188:SER:HB2	1:X:192:GLU:OE2	2.19	0.43
1:X:218:MSE:HE3	1:X:222:LYS:HZ1	1.83	0.43
1:X:85:ALA:O	1:X:147:TYR:HD1	2.01	0.43
1:X:330:ALA:HB2	1:X:339:MSE:CE	2.49	0.43
1:X:15:SER:OG	4:X:900:NAD:O5B	2.37	0.42
1:X:443:GLN:N	1:X:444:SER:HA	2.34	0.42
1:X:18:THR:N	1:X:19:PRO:CD	2.83	0.42
1:X:147:TYR:HE2	1:X:324:ILE:HD12	1.84	0.42
1:X:15:SER:OG	4:X:900:NAD:O1N	2.38	0.42
1:X:321:ALA:CB	1:X:324:ILE:HD11	2.49	0.42
1:X:244:ASN:HB3	5:X:1016:HOH:O	2.19	0.41
1:X:3:LYS:N	1:X:4:LYS:CA	2.78	0.41
1:X:249:LYS:HE2	1:X:275:MSE:HE3	2.02	0.41
1:X:293:ARG:O	1:X:296:PHE:HB3	2.19	0.41

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:X:931:HOH:O	5:X:931:HOH:O[4_575]	1.18	1.02

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	X	432/472 (92%)	410 (95%)	16 (4%)	6 (1%)	9 3

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	X	294	GLU
1	X	203	PHE
1	X	298	PHE
1	X	299	SER
1	X	443	GLN
1	X	296	PHE

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	X	373/386 (97%)	340 (91%)	33 (9%)	9 4

All (33) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	X	5	SER
1	X	17	PHE
1	X	18	THR
1	X	21	ILE
1	X	23	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	X	24	MSE
1	X	26	LEU
1	X	29	LEU
1	X	39	LEU
1	X	45	GLU
1	X	46	ARG
1	X	49	ARG
1	X	103	LEU
1	X	147	TYR
1	X	233	LYS
1	X	234	THR
1	X	242	SER
1	X	251	ARG
1	X	278	LYS
1	X	279	SER
1	X	294	GLU
1	X	306	ARG
1	X	307	GLU
1	X	308	GLN
1	X	311	GLU
1	X	314	GLU
1	X	319	ASP
1	X	324	ILE
1	X	325	VAL
1	X	413	LEU
1	X	442	ASP
1	X	443	GLN
1	X	444	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	X	146	ASN
1	X	169	ASN
1	X	210	GLN
1	X	227	GLN
1	X	300	GLN
1	X	320	HIS
1	X	409	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 3 ligands modelled in this entry, 1 is monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	NAD	X	900	-	46,48,48	1.32	6 (13%)	64,73,73	1.71	9 (14%)
2	G6P	X	902	3	16,16,16	1.12	1 (6%)	23,24,24	1.04	2 (8%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAD	X	900	-	-	7/30/62/62	0/5/5/5
2	G6P	X	902	3	1/1/6/6	1/6/26/26	0/1/1/1

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	X	900	NAD	O7N-C7N	5.26	1.33	1.24

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	X	902	G6P	P-O3P	3.52	1.61	1.50
4	X	900	NAD	C5A-N7A	-3.34	1.33	1.39
4	X	900	NAD	C2N-N1N	2.62	1.37	1.35
4	X	900	NAD	C8A-N9A	-2.25	1.33	1.37
4	X	900	NAD	C6N-N1N	2.15	1.40	1.35
4	X	900	NAD	PN-O3	2.10	1.61	1.59

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	X	900	NAD	C5A-C4A-N3A	-5.81	118.72	126.72
4	X	900	NAD	C4D-O4D-C1D	-4.84	105.49	109.92
4	X	900	NAD	N3A-C2A-N1A	-4.60	121.62	128.58
4	X	900	NAD	N3A-C4A-N9A	4.18	134.28	127.17
4	X	900	NAD	C2A-N3A-C4A	3.77	121.04	111.83
4	X	900	NAD	N9A-C8A-N7A	-2.97	109.72	113.94
4	X	900	NAD	C5A-N7A-C8A	2.67	107.64	103.45
2	X	902	G6P	O1P-P-O6	2.63	113.54	106.67
4	X	900	NAD	C4A-C5A-N7A	-2.37	107.87	110.58
4	X	900	NAD	C2B-C1B-N9A	-2.32	107.55	113.30
2	X	902	G6P	C1-O5-C5	-2.15	109.49	113.65

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
2	X	902	G6P	C1

All (8) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	X	900	NAD	C5B-O5B-PA-O2A
4	X	900	NAD	O4D-C1D-N1N-C6N
4	X	900	NAD	C3B-C4B-C5B-O5B
4	X	900	NAD	O4B-C4B-C5B-O5B
4	X	900	NAD	C5B-O5B-PA-O3
4	X	900	NAD	O4D-C1D-N1N-C2N
2	X	902	G6P	C6-O6-P-O1P
4	X	900	NAD	PA-O3-PN-O2N


There are no ring outliers.

1 monomer is involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	X	900	NAD	7	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	X	420/472 (88%)	0.64	46 (10%)	10 10	13, 29, 51, 76	1 (0%)

All (46) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	X	147	TYR	5.6
1	X	445	PRO	5.6
1	X	298	PHE	5.2
1	X	295	ALA	5.2
1	X	297	ILE	4.9
1	X	17	PHE	4.2
1	X	313	SER	3.7
1	X	316	LYS	3.4
1	X	293	ARG	3.4
1	X	286	ALA	3.3
1	X	296	PHE	3.3
1	X	317	ILE	3.3
1	X	243	TRP	3.2
1	X	283	HIS	3.1
1	X	442	ASP	3.0
1	X	312	ASN	3.0
1	X	101	ILE	2.8
1	X	213	GLU	2.8
1	X	276	VAL	2.8
1	X	292	GLY	2.7
1	X	320	HIS	2.7
1	X	287	ASN	2.7
1	X	54	CYS	2.7
1	X	314	GLU	2.6
1	X	319	ASP	2.6
1	X	301	CYS	2.5
1	X	234	THR	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	X	305	THR	2.5
1	X	294	GLU	2.5
1	X	300	GLN	2.5
1	X	304	ILE	2.4
1	X	272	PRO	2.4
1	X	444	SER	2.4
1	X	5	SER	2.3
1	X	97	LEU	2.3
1	X	148	SER	2.3
1	X	277	LYS	2.2
1	X	60	GLU	2.2
1	X	242	SER	2.2
1	X	315	ILE	2.2
1	X	321	ALA	2.2
1	X	289	VAL	2.1
1	X	48	ASP	2.1
1	X	102	PRO	2.1
1	X	57	PHE	2.1
1	X	307	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	NAD	X	900	44/44	0.80	0.36	39,45,48,48	44
2	G6P	X	902	16/16	0.86	0.27	44,52,53,53	16
3	MN	X	700	1/1	0.95	0.09	33,33,33,33	1

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



Electron density around G6P X 902:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.