



wwPDB X-ray Structure Validation Summary Report ⓘ

Jun 23, 2026 – 12:16 PM EDT

PDB ID : 4U8T / pdb_00004u8t
Title : Crystal structure of YTH domain of Zygosaccharomyces rouxii MRB1 protein
in complex with N6-Methyladenosine RNA
Authors : Luo, S.; Tong, L.
Deposited on : 2014-08-04
Resolution : 2.70 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3538 (2.70-2.70)
Clashscore	190562	3843 (2.70-2.70)
Ramachandran outliers	187476	3778 (2.70-2.70)
Sidechain outliers	187428	3778 (2.70-2.70)
RSRZ outliers	180081	3538 (2.70-2.70)
RNA backbone	3983	1044 (2.90-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	177	2% 75% 13% 11%
1	B	177	2% 75% 11% 13%
1	C	177	2% 72% 15% 12%
1	D	177	76% 12% 11%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	E	177	3% 71% 15% • 14%
1	F	177	2% 70% 15% • 14%
2	O	7	14% 43% 29% 14%
2	P	7	43% 43% 14%
2	Q	7	43% 29% 14% 14%
2	R	7	14% 14% 43% 29% 14%
2	S	7	43% 43% 14%
2	T	7	57% 29% 14%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 8572 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ZYRO0G01672p.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	157	Total	C	N	O	S	Se	0	0	0
			1278	823	223	229	1	2			
1	B	154	Total	C	N	O	S	Se	0	0	0
			1253	810	217	223	1	2			
1	C	155	Total	C	N	O	S	Se	0	0	0
			1262	815	221	223	1	2			
1	D	157	Total	C	N	O	S	Se	0	0	0
			1278	824	221	230	1	2			
1	E	153	Total	C	N	O	S	Se	0	0	0
			1245	806	216	220	1	2			
1	F	153	Total	C	N	O	S	Se	0	0	0
			1245	806	216	220	1	2			

There are 78 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	153	MSE	-	initiating methionine	UNP C5E1V0
A	154	GLY	-	expression tag	UNP C5E1V0
A	155	SER	-	expression tag	UNP C5E1V0
A	156	SER	-	expression tag	UNP C5E1V0
A	157	HIS	-	expression tag	UNP C5E1V0
A	158	HIS	-	expression tag	UNP C5E1V0
A	159	HIS	-	expression tag	UNP C5E1V0
A	160	HIS	-	expression tag	UNP C5E1V0
A	161	HIS	-	expression tag	UNP C5E1V0
A	162	HIS	-	expression tag	UNP C5E1V0
A	163	SER	-	expression tag	UNP C5E1V0
A	164	SER	-	expression tag	UNP C5E1V0
A	165	GLY	-	expression tag	UNP C5E1V0
B	153	MSE	-	initiating methionine	UNP C5E1V0
B	154	GLY	-	expression tag	UNP C5E1V0
B	155	SER	-	expression tag	UNP C5E1V0
B	156	SER	-	expression tag	UNP C5E1V0

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	157	HIS	-	expression tag	UNP C5E1V0
B	158	HIS	-	expression tag	UNP C5E1V0
B	159	HIS	-	expression tag	UNP C5E1V0
B	160	HIS	-	expression tag	UNP C5E1V0
B	161	HIS	-	expression tag	UNP C5E1V0
B	162	HIS	-	expression tag	UNP C5E1V0
B	163	SER	-	expression tag	UNP C5E1V0
B	164	SER	-	expression tag	UNP C5E1V0
B	165	GLY	-	expression tag	UNP C5E1V0
C	153	MSE	-	initiating methionine	UNP C5E1V0
C	154	GLY	-	expression tag	UNP C5E1V0
C	155	SER	-	expression tag	UNP C5E1V0
C	156	SER	-	expression tag	UNP C5E1V0
C	157	HIS	-	expression tag	UNP C5E1V0
C	158	HIS	-	expression tag	UNP C5E1V0
C	159	HIS	-	expression tag	UNP C5E1V0
C	160	HIS	-	expression tag	UNP C5E1V0
C	161	HIS	-	expression tag	UNP C5E1V0
C	162	HIS	-	expression tag	UNP C5E1V0
C	163	SER	-	expression tag	UNP C5E1V0
C	164	SER	-	expression tag	UNP C5E1V0
C	165	GLY	-	expression tag	UNP C5E1V0
D	153	MSE	-	initiating methionine	UNP C5E1V0
D	154	GLY	-	expression tag	UNP C5E1V0
D	155	SER	-	expression tag	UNP C5E1V0
D	156	SER	-	expression tag	UNP C5E1V0
D	157	HIS	-	expression tag	UNP C5E1V0
D	158	HIS	-	expression tag	UNP C5E1V0
D	159	HIS	-	expression tag	UNP C5E1V0
D	160	HIS	-	expression tag	UNP C5E1V0
D	161	HIS	-	expression tag	UNP C5E1V0
D	162	HIS	-	expression tag	UNP C5E1V0
D	163	SER	-	expression tag	UNP C5E1V0
D	164	SER	-	expression tag	UNP C5E1V0
D	165	GLY	-	expression tag	UNP C5E1V0
E	153	MSE	-	initiating methionine	UNP C5E1V0
E	154	GLY	-	expression tag	UNP C5E1V0
E	155	SER	-	expression tag	UNP C5E1V0
E	156	SER	-	expression tag	UNP C5E1V0
E	157	HIS	-	expression tag	UNP C5E1V0
E	158	HIS	-	expression tag	UNP C5E1V0
E	159	HIS	-	expression tag	UNP C5E1V0

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
E	160	HIS	-	expression tag	UNP C5E1V0
E	161	HIS	-	expression tag	UNP C5E1V0
E	162	HIS	-	expression tag	UNP C5E1V0
E	163	SER	-	expression tag	UNP C5E1V0
E	164	SER	-	expression tag	UNP C5E1V0
E	165	GLY	-	expression tag	UNP C5E1V0
F	153	MSE	-	initiating methionine	UNP C5E1V0
F	154	GLY	-	expression tag	UNP C5E1V0
F	155	SER	-	expression tag	UNP C5E1V0
F	156	SER	-	expression tag	UNP C5E1V0
F	157	HIS	-	expression tag	UNP C5E1V0
F	158	HIS	-	expression tag	UNP C5E1V0
F	159	HIS	-	expression tag	UNP C5E1V0
F	160	HIS	-	expression tag	UNP C5E1V0
F	161	HIS	-	expression tag	UNP C5E1V0
F	162	HIS	-	expression tag	UNP C5E1V0
F	163	SER	-	expression tag	UNP C5E1V0
F	164	SER	-	expression tag	UNP C5E1V0
F	165	GLY	-	expression tag	UNP C5E1V0

- Molecule 2 is a RNA chain called RNA (5'-R(*AP*GP*GP*(6MZ)P*CP*AP*U)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	O	7	Total	C	N	O	P	0	0	0
			150	69	30	45	6			
2	P	7	Total	C	N	O	P	0	0	0
			150	69	30	45	6			
2	Q	7	Total	C	N	O	P	0	0	0
			150	69	30	45	6			
2	R	7	Total	C	N	O	P	0	0	0
			150	69	30	45	6			
2	S	7	Total	C	N	O	P	0	0	0
			150	69	30	45	6			
2	T	7	Total	C	N	O	P	0	0	0
			150	69	30	45	6			

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	21	Total	O	0	0
			21	21		
3	B	10	Total	O	0	0
			10	10		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	C	10	Total 10	O 10	0	0
3	D	16	Total 16	O 16	0	0
3	E	10	Total 10	O 10	0	0
3	F	15	Total 15	O 15	0	0
3	O	5	Total 5	O 5	0	0
3	P	4	Total 4	O 4	0	0
3	Q	6	Total 6	O 6	0	0
3	R	2	Total 2	O 2	0	0
3	S	5	Total 5	O 5	0	0
3	T	7	Total 7	O 7	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: ZYRO0G01672p

Chain A: 




• Molecule 1: ZYRO0G01672p

Chain B: 



• Molecule 1: ZYRO0G01672p

Chain C: 





• Molecule 1: ZYRO0G01672p

Chain D: 






- Molecule 1: ZYRO0G01672p




- Molecule 1: ZYRO0G01672p



- Molecule 2: RNA (5'-R(*AP*GP*GP*(6MZ)P*CP*AP*U)-3')



- Molecule 2: RNA (5'-R(*AP*GP*GP*(6MZ)P*CP*AP*U)-3')



- Molecule 2: RNA (5'-R(*AP*GP*GP*(6MZ)P*CP*AP*U)-3')




- Molecule 2: RNA (5'-R(*AP*GP*GP*(6MZ)P*CP*AP*U)-3')



- Molecule 2: RNA (5'-R(*AP*GP*GP*(6MZ)P*CP*AP*U)-3')

Chain S:  43% 43% 14%



- Molecule 2: RNA (5'-R(*AP*GP*GP*(6MZ)P*CP*AP*U)-3')

Chain T:  57% 29% 14%



4 Data and refinement statistics

Property	Value	Source
Space group	P 61 2 2	Depositor
Cell constants a, b, c, α , β , γ	117.39Å 117.39Å 380.93Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	24.92 – 2.70 24.92 – 2.70	Depositor EDS
% Data completeness (in resolution range)	94.2 (24.92-2.70) 88.1 (24.92-2.70)	Depositor EDS
R_{merge}	0.14	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.07 (at 2.72Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.9_1675)	Depositor
R, R_{free}	0.199 , 0.245 0.203 , 0.248	Depositor DCC
R_{free} test set	1849 reflections (4.19%)	wwPDB-VP
Wilson B-factor (Å ²)	39.7	Xtriage
Anisotropy	0.094	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 38.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	8572	wwPDB-VP
Average B, all atoms (Å ²)	34.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.52% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: 6MZ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.98	0/1309	0.96	3/1764 (0.2%)
1	B	0.94	1/1284 (0.1%)	0.94	4/1731 (0.2%)
1	C	0.90	0/1293	0.89	2/1742 (0.1%)
1	D	0.94	0/1310	0.88	1/1767 (0.1%)
1	E	0.92	0/1276	0.91	4/1720 (0.2%)
1	F	0.83	0/1276	0.89	5/1720 (0.3%)
2	O	0.85	0/141	0.79	0/216
2	P	0.74	0/141	0.80	0/216
2	Q	0.69	0/141	0.88	0/216
2	R	0.70	0/141	0.78	0/216
2	S	0.67	0/141	0.80	0/216
2	T	0.76	0/141	0.76	0/216
All	All	0.90	1/8594 (0.0%)	0.90	19/11740 (0.2%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	169	PRO	CA-C	5.83	1.55	1.51

The worst 5 of 19 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	219	GLN	N-CA-C	7.42	119.37	111.28
1	A	269	ILE	CB-CA-C	-6.89	104.90	111.44
1	C	168	ILE	CA-C-N	6.68	127.26	120.38
1	C	168	ILE	C-N-CA	6.68	127.26	120.38
1	A	219	GLN	N-CA-C	6.62	118.16	111.07

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1278	0	1266	18	1
1	B	1253	0	1244	12	1
1	C	1262	0	1258	29	0
1	D	1278	0	1266	17	0
1	E	1245	0	1240	20	0
1	F	1245	0	1240	23	0
2	O	150	0	80	8	0
2	P	150	0	80	2	0
2	Q	150	0	80	4	0
2	R	150	0	80	6	0
2	S	150	0	80	3	0
2	T	150	0	80	3	0
3	A	21	0	0	1	0
3	B	10	0	0	0	0
3	C	10	0	0	0	0
3	D	16	0	0	0	0
3	E	10	0	0	0	0
3	F	15	0	0	0	0
3	O	5	0	0	0	0
3	P	4	0	0	0	0
3	Q	6	0	0	0	0
3	R	2	0	0	0	0
3	S	5	0	0	0	0
3	T	7	0	0	0	0
All	All	8572	0	7994	120	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 120 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:259:ARG:NH2	2:O:0:6MZ:OP2	2.05	0.90
1:D:217:LEU:HD11	1:D:223:ILE:HG13	1.58	0.86
1:B:210:LEU:HB3	1:B:241:MSE:HE1	1.64	0.80

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:225:LEU:HD21	1:E:241:MSE:HE3	1.62	0.79
1:C:182:VAL:HG23	1:C:301:ILE:HD11	1.64	0.79

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:189:GLU:OE2	1:B:196:TYR:OH[5_554]	2.01	0.19

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	151/177 (85%)	145 (96%)	6 (4%)	0	100	100
1	B	148/177 (84%)	143 (97%)	5 (3%)	0	100	100
1	C	149/177 (84%)	144 (97%)	5 (3%)	0	100	100
1	D	153/177 (86%)	148 (97%)	5 (3%)	0	100	100
1	E	147/177 (83%)	142 (97%)	5 (3%)	0	100	100
1	F	147/177 (83%)	142 (97%)	5 (3%)	0	100	100
All	All	895/1062 (84%)	864 (96%)	31 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	142/158 (90%)	142 (100%)	0	100	100
1	B	139/158 (88%)	134 (96%)	5 (4%)	31	60
1	C	140/158 (89%)	138 (99%)	2 (1%)	59	82
1	D	142/158 (90%)	141 (99%)	1 (1%)	76	90
1	E	138/158 (87%)	137 (99%)	1 (1%)	76	90
1	F	138/158 (87%)	135 (98%)	3 (2%)	45	74
All	All	839/948 (88%)	827 (99%)	12 (1%)	59	82

5 of 12 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	289	MSE
1	E	305	ILE
1	F	289	MSE
1	F	215	GLU
1	B	281	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 14 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	299	GLN
1	D	230	ASN
1	F	299	GLN
1	E	299	GLN
1	F	230	ASN

5.3.3 RNA ⓘ

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
2	O	6/7 (85%)	4 (66%)	0
2	P	6/7 (85%)	3 (50%)	0
2	Q	6/7 (85%)	4 (66%)	0
2	R	6/7 (85%)	4 (66%)	0
2	S	6/7 (85%)	3 (50%)	0
2	T	6/7 (85%)	2 (33%)	0
All	All	36/42 (85%)	20 (55%)	0

5 of 20 RNA backbone outliers are listed below:

Mol	Chain	Res	Type
2	O	-2	G
2	O	-1	G
2	O	0	6MZ
2	O	1	C
2	P	-2	G

There are no RNA pucker outliers to report.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

6 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	6MZ	Q	0	2	22,25,26	1.55	6 (27%)	29,36,39	2.21	10 (34%)
2	6MZ	T	0	2	22,25,26	1.48	4 (18%)	29,36,39	2.41	11 (37%)
2	6MZ	O	0	2	22,25,26	1.40	5 (22%)	29,36,39	2.37	11 (37%)
2	6MZ	R	0	2	22,25,26	1.47	5 (22%)	29,36,39	2.25	10 (34%)
2	6MZ	S	0	2	22,25,26	1.46	4 (18%)	29,36,39	2.50	12 (41%)
2	6MZ	P	0	2	22,25,26	1.53	5 (22%)	29,36,39	2.15	9 (31%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	6MZ	Q	0	2	-	2/9/27/28	0/3/3/3
2	6MZ	T	0	2	-	0/9/27/28	0/3/3/3
2	6MZ	O	0	2	-	0/9/27/28	0/3/3/3
2	6MZ	R	0	2	-	0/9/27/28	0/3/3/3
2	6MZ	S	0	2	-	0/9/27/28	0/3/3/3

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	6MZ	P	0	2	-	0/9/27/28	0/3/3/3

The worst 5 of 29 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	R	0	6MZ	C5-C4	4.12	1.46	1.39
2	P	0	6MZ	C5-C4	4.03	1.46	1.39
2	S	0	6MZ	C5-C4	3.96	1.46	1.39
2	Q	0	6MZ	C5-C4	3.81	1.45	1.39
2	O	0	6MZ	C5-C4	3.62	1.45	1.39

The worst 5 of 63 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	S	0	6MZ	C9-N6-C6	-6.37	116.94	122.85
2	O	0	6MZ	C5-C4-N3	-5.94	118.54	126.72
2	R	0	6MZ	C5-C4-N3	-5.75	118.80	126.72
2	P	0	6MZ	C5-C4-N3	-5.34	119.37	126.72
2	Q	0	6MZ	C6-C5-N7	5.22	138.12	132.43

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	Q	0	6MZ	C5-C6-N6-C9
2	Q	0	6MZ	N1-C6-N6-C9

There are no ring outliers.

4 monomers are involved in 11 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	Q	0	6MZ	1	0
2	O	0	6MZ	5	0
2	R	0	6MZ	2	0
2	S	0	6MZ	3	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	155/177 (87%)	-0.27	4 (2%) 57 54	13, 25, 53, 67	0
1	B	152/177 (85%)	-0.16	4 (2%) 57 54	19, 30, 55, 84	0
1	C	153/177 (86%)	0.00	3 (1%) 65 62	20, 32, 58, 86	0
1	D	155/177 (87%)	-0.32	0 100 100	17, 31, 48, 69	0
1	E	151/177 (85%)	-0.12	5 (3%) 49 45	17, 32, 59, 76	0
1	F	151/177 (85%)	0.12	3 (1%) 65 62	23, 39, 58, 73	0
2	O	6/7 (85%)	-0.36	0 100 100	22, 26, 38, 56	0
2	P	6/7 (85%)	-0.28	0 100 100	23, 35, 47, 66	0
2	Q	6/7 (85%)	-0.22	0 100 100	24, 37, 43, 63	0
2	R	6/7 (85%)	0.19	1 (16%) 4 3	27, 38, 54, 73	0
2	S	6/7 (85%)	0.09	0 100 100	22, 34, 51, 69	0
2	T	6/7 (85%)	-0.29	0 100 100	22, 31, 43, 56	0
All	All	953/1104 (86%)	-0.13	20 (2%) 63 61	13, 32, 56, 86	0

The worst 5 of 20 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	171	TRP	3.8
1	F	171	TRP	3.7
1	E	320	VAL	3.6
1	B	171	TRP	3.5
1	E	165	GLY	3.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	6MZ	O	0	23/24	0.97	0.06	17,20,23,31	0
2	6MZ	R	0	23/24	0.97	0.07	26,31,35,36	0
2	6MZ	S	0	23/24	0.97	0.07	17,26,31,32	0
2	6MZ	P	0	23/24	0.98	0.06	18,21,28,34	0
2	6MZ	Q	0	23/24	0.98	0.06	21,25,27,36	0
2	6MZ	T	0	23/24	0.98	0.05	17,20,23,30	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.