



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 10:15 PM UTC

PDB ID : 5U89 / pdb_00005u89
Title : Crystal structure of a cross-module fragment from the dimodular NRPS Dhbf
Authors : Tarry, M.J.; Schmeing, T.M.
Deposited on : 2016-12-14
Resolution : 3.08 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.08 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2010 (3.10-3.06)
Clashscore	190562	2102 (3.10-3.06)
Ramachandran outliers	187476	1982 (3.10-3.06)
Sidechain outliers	187428	1981 (3.10-3.06)
RSRZ outliers	180081	2010 (3.10-3.06)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	1080	6% 87% 9% .
2	B	83	5% 60% 17% 23%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 8107 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Amino acid adenylation domain protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	1039	7567	4895	1300	1345	27	0	0	0

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	434	GLY	-	expression tag	UNP A0A0F6BHX2
A	435	ALA	-	expression tag	UNP A0A0F6BHX2
A	436	MET	-	expression tag	UNP A0A0F6BHX2
A	1505	ALA	-	expression tag	UNP A0A0F6BHX2
A	1506	ALA	-	expression tag	UNP A0A0F6BHX2
A	1507	ALA	-	expression tag	UNP A0A0F6BHX2
A	1508	GLU	-	expression tag	UNP A0A0F6BHX2
A	1509	ASN	-	expression tag	UNP A0A0F6BHX2
A	1510	LEU	-	expression tag	UNP A0A0F6BHX2
A	1511	TYR	-	expression tag	UNP A0A0F6BHX2
A	1512	PHE	-	expression tag	UNP A0A0F6BHX2
A	1513	GLN	-	expression tag	UNP A0A0F6BHX2


- Molecule 2 is a protein called MbtH domain protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	64	493	318	84	89	2	0	0	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	1	GLY	-	expression tag	UNP A0A0F6BHX3
B	2	ALA	-	expression tag	UNP A0A0F6BHX3
B	3	MET	-	expression tag	UNP A0A0F6BHX3
B	4	GLY	-	expression tag	UNP A0A0F6BHX3

- Molecule 3 is 5'-({[(2R)-3-amino-2-{[2-({N-[(2R)-2-hydroxy-3,3-dimethyl-4-(phosphonoox y)butanoyl]-beta-alanyl}amino)ethyl]sulfanyl}propyl]sulfonyl}amino)-5'-deoxyadenosine (CCD ID: MJ8) (formula: C₂₄H₄₂N₉O₁₂PS₂).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
			Total	C	N	O	P	S		
3	A	1	47	24	9	11	1	2	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Amino acid adenylation domain protein



- Molecule 2: MbtH domain protein



4 Data and refinement statistics

Property	Value	Source
Space group	P 65 2 2	Depositor
Cell constants a, b, c, α , β , γ	119.51Å 119.51Å 589.65Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	58.56 – 3.08 58.56 – 3.08	Depositor EDS
% Data completeness (in resolution range)	98.5 (58.56-3.08) 98.5 (58.56-3.08)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.57 (at 3.07Å)	Xtriage
Refinement program	PHENIX 1.9_1692	Depositor
R, R_{free}	0.216 , 0.259 0.224 , 0.258	Depositor DCC
R_{free} test set	2296 reflections (4.80%)	wwPDB-VP
Wilson B-factor (Å ²)	69.1	Xtriage
Anisotropy	0.373	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 75.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.42$, $\langle L^2 \rangle = 0.24$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.90	EDS
Total number of atoms	8107	wwPDB-VP
Average B, all atoms (Å ²)	78.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.67% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MJ8

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.29	0/7749	0.74	5/10641 (0.0%)
2	B	0.28	0/509	0.72	0/700
All	All	0.28	0/8258	0.74	5/11341 (0.0%)

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	901	ILE	CA-C-N	6.14	126.11	120.03
1	A	901	ILE	C-N-CA	6.14	126.11	120.03
1	A	1396	HIS	CA-C-N	5.14	126.27	119.84
1	A	1396	HIS	C-N-CA	5.14	126.27	119.84
1	A	479	GLU	CB-CA-C	-5.02	110.40	117.23

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	7567	0	7153	50	0
2	B	493	0	452	9	0
3	A	47	0	0	0	0
All	All	8107	0	7605	57	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (57) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:759:SER:HB3	1:A:772:GLY:HA3	1.71	0.71
1:A:1184:ILE:HG13	1:A:1185:VAL:HG23	1.74	0.69
1:A:1082:VAL:HG11	1:A:1413:LEU:HD12	1.74	0.67
1:A:840:ARG:NH2	1:A:848:ASP:OD2	2.32	0.62
1:A:549:MET:HE2	1:A:610:TYR:HB2	1.83	0.59
1:A:1330:ARG:NH1	1:A:1335:LEU:O	2.35	0.59
1:A:965:PRO:HB2	1:A:971:GLU:HG2	1.86	0.57
1:A:631:PHE:HZ	1:A:660:LEU:HD12	1.71	0.56
2:B:18:VAL:HB	2:B:43:PHE:HB3	1.87	0.55
1:A:489:ASN:OD1	1:A:529:LYS:NZ	2.40	0.55
1:A:647:LEU:HD21	1:A:661:GLU:HA	1.88	0.54
1:A:1064:TRP:HB2	1:A:1183:MET:HE1	1.90	0.54
1:A:459:PHE:HB2	1:A:667:ILE:HG21	1.91	0.53
2:B:21:ASN:OD1	2:B:22:ASP:N	2.41	0.53
1:A:552:ASP:OD2	1:A:610:TYR:OH	2.21	0.53
1:A:584:THR:O	1:A:588:ILE:HG12	2.10	0.51
1:A:603:LEU:HB2	2:B:67:LEU:HD21	1.92	0.51
1:A:1305:GLN:HB3	1:A:1350:MET:HE1	1.92	0.51
1:A:998:HIS:CE1	1:A:1000:LEU:HB3	2.45	0.50
1:A:716:ILE:HG22	1:A:740:ALA:HB2	1.93	0.49
1:A:1142:LEU:HD22	1:A:1169:LEU:HD11	1.95	0.48
1:A:1230:LEU:HD11	1:A:1379:PRO:HG2	1.95	0.48
1:A:866:ILE:O	1:A:870:GLU:HG3	2.14	0.48
1:A:1058:PHE:CE1	1:A:1230:LEU:HD12	2.49	0.48
2:B:46:ASN:OD1	2:B:47:THR:N	2.44	0.48
1:A:814:ASP:OD1	1:A:815:LEU:N	2.48	0.47
1:A:1281:MET:HB2	1:A:1451:LEU:HB2	1.95	0.47
1:A:1457:TYR:CE1	1:A:1467:VAL:HG21	2.49	0.47
1:A:752:THR:HG21	1:A:800:ILE:HG21	1.96	0.46
1:A:798:LEU:HD22	1:A:839:VAL:HG21	1.98	0.45
1:A:631:PHE:HE1	1:A:755:TRP:HB2	1.81	0.45
1:A:527:VAL:O	1:A:530:THR:HG22	2.15	0.45
1:A:705:TRP:CZ2	1:A:721:VAL:HG13	2.53	0.45
1:A:870:GLU:OE2	1:A:883:LEU:HA	2.18	0.44
1:A:881:ARG:HH22	1:A:937:GLU:CD	2.25	0.44
2:B:30:PRO:HB2	2:B:33:ILE:HD13	2.00	0.44
1:A:799:TYR:CE2	1:A:834:ARG:HB3	2.53	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1286:HIS:CE1	1:A:1440:GLU:HG3	2.54	0.43
1:A:1045:VAL:HG22	1:A:1225:TRP:CD2	2.54	0.43
2:B:21:ASN:HB3	2:B:25:GLN:H	1.83	0.43
1:A:631:PHE:CZ	1:A:660:LEU:HD12	2.52	0.43
2:B:62:MET:HE3	2:B:62:MET:HB3	1.83	0.43
1:A:785:GLU:O	2:B:8:PRO:HG3	2.19	0.43
1:A:789:PRO:HD3	1:A:825:TYR:OH	2.19	0.43
1:A:824:PRO:HG2	1:A:825:TYR:CE2	2.53	0.43
1:A:1480:GLU:HG2	1:A:1492:ILE:HG23	2.02	0.42
1:A:1359:ARG:HH21	1:A:1486:ASP:CG	2.26	0.42
1:A:527:VAL:HG11	1:A:534:TYR:HB3	2.02	0.42
2:B:35:ILE:HA	2:B:36:PRO:HD3	1.91	0.42
1:A:656:ASP:HB2	1:A:754:ILE:HG12	2.02	0.41
1:A:839:VAL:HG11	1:A:847:LEU:HD23	2.01	0.41
1:A:809:TYR:HB2	1:A:812:ARG:HB2	2.02	0.41
1:A:538:ASP:HB3	1:A:541:TYR:CD1	2.56	0.41
1:A:746:LEU:HD13	1:A:755:TRP:NE1	2.36	0.41
1:A:866:ILE:HG23	1:A:884:VAL:HB	2.02	0.41
1:A:1436:LEU:HD13	1:A:1455:PHE:CZ	2.56	0.40
1:A:928:VAL:HA	1:A:929:PRO:HD3	1.88	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	1025/1080 (95%)	996 (97%)	27 (3%)	2 (0%)	43	71
2	B	62/83 (75%)	60 (97%)	2 (3%)	0	100	100
All	All	1087/1163 (94%)	1056 (97%)	29 (3%)	2 (0%)	43	71

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	754	ILE
1	A	753	THR

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	721/928 (78%)	715 (99%)	6 (1%)	73	79
2	B	49/69 (71%)	49 (100%)	0	100	100
All	All	770/997 (77%)	764 (99%)	6 (1%)	73	79

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	440	ARG
1	A	545	ARG
1	A	586	GLN
1	A	883	LEU
1	A	1000	LEU
1	A	1229	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	745	ASN
1	A	788	GLN
1	A	855	HIS
1	A	1243	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	MJ8	A	1601	1	44,49,50	1.13	5 (11%)	57,70,73	2.19	12 (21%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	MJ8	A	1601	1	-	14/42/62/63	0/3/3/3

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	1601	MJ8	C17-S16	-3.63	1.66	1.77
3	A	1601	MJ8	C22-S21	-3.28	1.76	1.81
3	A	1601	MJ8	C18-S21	-3.04	1.77	1.83
3	A	1601	MJ8	C09-N08	2.10	1.35	1.31
3	A	1601	MJ8	C07-N08	-2.07	1.35	1.39

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	1601	MJ8	O42-S16-O43	-7.12	109.38	119.34
3	A	1601	MJ8	C22-S21-C18	6.11	110.75	101.67
3	A	1601	MJ8	C14-N15-S16	-5.76	109.02	120.79
3	A	1601	MJ8	C07-C06-N05	-5.24	119.50	126.72
3	A	1601	MJ8	N03-C04-N05	-4.86	121.23	128.58
3	A	1601	MJ8	N10-C09-N08	-3.55	108.91	113.94
3	A	1601	MJ8	C04-N05-C06	3.46	120.28	111.83
3	A	1601	MJ8	N05-C06-N10	3.39	132.93	127.17
3	A	1601	MJ8	C06-C07-N08	-2.75	107.44	110.58
3	A	1601	MJ8	C07-N08-C09	2.64	107.59	103.45
3	A	1601	MJ8	C06-N10-C09	2.62	108.49	105.74
3	A	1601	MJ8	C27-C26-C25	-2.24	108.66	112.39

There are no chirality outliers.

All (14) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	A	1601	MJ8	C14-N15-S16-C17
3	A	1601	MJ8	C18-C17-S16-O43
3	A	1601	MJ8	C18-C17-S16-O42
3	A	1601	MJ8	C18-C17-S16-N15
3	A	1601	MJ8	O12-C13-C14-N15
3	A	1601	MJ8	C25-C26-C27-N28
3	A	1601	MJ8	O40-C29-N28-C27
3	A	1601	MJ8	C30-C29-N28-C27
3	A	1601	MJ8	C30-C32-C35-O36
3	A	1601	MJ8	C33-C32-C35-O36
3	A	1601	MJ8	C34-C32-C35-O36
3	A	1601	MJ8	S21-C22-C23-N24
3	A	1601	MJ8	S16-C17-C18-C19
3	A	1601	MJ8	C17-C18-S21-C22

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the

average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	1039/1080 (96%)	0.44	70 (6%) 24 12	43, 76, 118, 168	0
2	B	64/83 (77%)	0.28	4 (6%) 26 13	51, 70, 112, 147	0
All	All	1103/1163 (94%)	0.43	74 (6%) 24 12	43, 75, 117, 168	0

All (74) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	770	SER	5.3
1	A	769	PRO	4.9
1	A	1042	LYS	4.7
2	B	6	THR	4.3
1	A	616	GLY	4.1
1	A	457	GLU	4.0
1	A	599	THR	3.9
1	A	1114	HIS	3.9
1	A	1206	ARG	3.8
1	A	1397	PRO	3.8
2	B	22	ASP	3.7
1	A	614	SER	3.7
1	A	1126	HIS	3.6
1	A	737	HIS	3.6
1	A	843	MET	3.5
1	A	1387	LEU	3.4
1	A	889	ASP	3.4
1	A	848	ASP	3.4
2	B	47	THR	3.3
1	A	1052	GLU	3.2
1	A	956	PHE	3.1
1	A	1205	CYS	3.1
1	A	1043	PRO	3.0
1	A	734	SER	3.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	1118	SER	3.0
1	A	793	GLY	2.9
1	A	1445	ASP	2.9
1	A	1388	ASN	2.9
1	A	913	ALA	2.9
1	A	904	GLU	2.8
1	A	730	SER	2.8
1	A	1396	HIS	2.7
1	A	1493	ASP	2.7
1	A	894	GLN	2.7
1	A	1269	GLU	2.6
1	A	1050	CYS	2.6
1	A	989	ASP	2.6
1	A	644	HIS	2.6
1	A	580	ASP	2.6
1	A	1446	GLY	2.5
1	A	1202	ASN	2.5
1	A	955	ASP	2.5
1	A	1377	ASP	2.5
1	A	1229	ILE	2.5
2	B	69	ASP	2.5
1	A	452	HIS	2.5
1	A	846	SER	2.4
1	A	946	ILE	2.4
1	A	1291	ASP	2.4
1	A	453	PRO	2.4
1	A	1033	LYS	2.4
1	A	440	ARG	2.3
1	A	1237	ASP	2.3
1	A	1212	TRP	2.3
1	A	1354	GLY	2.3
1	A	1414	GLU	2.3
1	A	656	ASP	2.3
1	A	1261	PRO	2.2
1	A	1234	THR	2.2
1	A	1408	THR	2.2
1	A	461	GLN	2.2
1	A	617	LYS	2.2
1	A	1273	GLU	2.2
1	A	1211	GLU	2.2
1	A	1385	GLU	2.2
1	A	615	THR	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	855	HIS	2.1
1	A	957	THR	2.1
1	A	571	PHE	2.1
1	A	900	VAL	2.1
1	A	458	ASN	2.1
1	A	570	HIS	2.0
1	A	472	GLU	2.0
1	A	863	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	MJ8	A	1601	47/48	0.87	0.19	67,104,222,290	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around MJ8 A 1601:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.