



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 25, 2026 – 02:24 AM EDT

PDB ID : 1U95 / pdb_00001u95
Title : Crystal structure of the HIV-1 Cross Neutralizing Monoclonal Antibody 2F5
in complex with gp41 Peptide ELDHWAS
Authors : Bryson, S.; Julien, J.-P.; Hynes, R.C.; Pai, E.F.
Deposited on : 2004-08-09
Resolution : 2.24 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.24 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3416 (2.26-2.22)
Clashscore	190562	3556 (2.26-2.22)
Ramachandran outliers	187476	3500 (2.26-2.22)
Sidechain outliers	187428	3501 (2.26-2.22)
RSRZ outliers	180081	3415 (2.26-2.22)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	214	4% 67% 29% .
2	B	235	7% 66% 26% ...
3	C	7	14% 100%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 3591 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ANTIBODY 2F5 (LIGHT CHAIN).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	214	Total	C	N	O	S	0	0	0
			1644	1025	281	333	5			

- Molecule 2 is a protein called ANTIBODY 2F5 (HEAVY CHAIN).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	227	Total	C	N	O	S	0	0	0
			1705	1084	289	325	7			

- Molecule 3 is a protein called GP41 PEPTIDE.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	C	7	Total	C	N	O	0	0	0
			61	38	10	13			


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	86	Total	O	0	0
			86	86		
4	B	91	Total	O	0	0
			91	91		
4	C	4	Total	O	0	0
			4	4		

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: ANTIBODY 2F5 (LIGHT CHAIN)



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	57.70Å 64.90Å 175.90Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	80.00 – 2.24 80.00 – 2.24	Depositor EDS
% Data completeness (in resolution range)	95.7 (80.00-2.24) 95.6 (80.00-2.24)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.06	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.39 (at 2.22Å)	Xtriage
Refinement program	CNS 1.0	Depositor
R, R_{free}	0.210 , 0.238 0.212 , 0.240	Depositor DCC
R_{free} test set	1579 reflections (4.80%)	wwPDB-VP
Wilson B-factor (Å ²)	38.9	Xtriage
Anisotropy	0.109	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 35.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	3591	wwPDB-VP
Average B, all atoms (Å ²)	37.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.59% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.06	1/1681 (0.1%)	1.24	12/2284 (0.5%)
2	B	1.10	3/1746 (0.2%)	1.30	13/2385 (0.5%)
3	C	0.78	0/63	1.09	0/84
All	All	1.08	4/3490 (0.1%)	1.27	25/4753 (0.5%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	40	PRO	CA-C	6.87	1.55	1.51
2	B	216	CYS	CB-SG	5.47	1.99	1.81
2	B	87	THR	CA-CB	5.30	1.61	1.53
1	A	193	GLU	CG-CD	5.10	1.64	1.52

All (25) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	97	THR	N-CA-C	10.36	123.71	110.24
2	B	143	LYS	N-CA-C	8.23	122.84	109.59
1	A	114	SER	N-CA-C	-7.54	96.95	109.46
2	B	101	ASP	N-CA-C	7.25	121.36	112.23
1	A	142	ARG	N-CA-C	7.21	119.14	111.28
2	B	100(N)	MET	N-CA-C	-7.05	95.50	107.99
1	A	75	ILE	N-CA-C	-7.04	97.58	107.80
2	B	120	SER	N-CA-C	-6.70	98.77	109.76
1	A	83	PHE	N-CA-C	-6.60	98.43	108.67

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	30	SER	N-CA-C	-6.40	104.73	113.30
2	B	114	THR	N-CA-C	6.35	119.45	110.50
2	B	56	ASP	N-CA-C	-6.13	101.21	110.28
2	B	144	ASP	N-CA-C	5.93	119.29	111.28
2	B	38	ARG	CB-CG-CD	5.92	124.91	111.30
2	B	38	ARG	NE-CZ-NH2	-5.91	113.88	119.20
1	A	97	THR	CB-CA-C	-5.85	99.28	109.64
1	A	203	SER	N-CA-C	-5.42	101.90	108.14
1	A	170	ASP	CB-CA-C	-5.38	100.85	110.70
2	B	55	ASP	N-CA-C	5.33	119.74	113.23
1	A	63	SER	CA-C-N	-5.26	118.07	121.65
1	A	63	SER	C-N-CA	-5.26	118.07	121.65
2	B	208	ASP	N-CA-C	-5.25	100.34	108.90
1	A	196	VAL	N-CA-C	5.22	115.42	108.11
2	B	38	ARG	NE-CZ-NH1	5.20	126.70	121.50
1	A	48	ILE	CB-CA-C	-5.06	102.95	110.33

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	86	TYR	Sidechain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1644	0	1588	56	0
2	B	1705	0	1721	60	0
3	C	61	0	50	0	0
4	A	86	0	0	1	0
4	B	91	0	0	1	0
4	C	4	0	0	0	0
All	All	3591	0	3359	109	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 16.

All (109) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:84:PRO:O	2:B:87:THR:HG23	1.65	0.96
1:A:90:GLN:HE22	1:A:93:PHE:H	1.01	0.93
1:A:20:THR:HG22	1:A:74:THR:HB	1.50	0.91
2:B:86:ASP:HB2	2:B:111:ILE:HD12	1.59	0.84
1:A:106:VAL:H	1:A:166:GLN:HE22	1.26	0.81
2:B:38:ARG:HD2	2:B:48:LEU:HD21	1.60	0.80
1:A:90:GLN:NE2	1:A:93:PHE:H	1.81	0.76
1:A:90:GLN:NE2	1:A:92:HIS:H	1.84	0.75
1:A:162:SER:HB2	2:B:167:PRO:HG2	1.71	0.70
2:B:210:ARG:NH1	2:B:212:GLU:OE1	2.25	0.70
2:B:29:LEU:O	2:B:53:SER:HB2	1.93	0.69
1:A:133:VAL:HG21	2:B:124:LEU:HD22	1.75	0.67
1:A:118:PHE:HB2	1:A:133:VAL:HG22	1.77	0.67
1:A:90:GLN:HE21	1:A:92:HIS:H	1.42	0.64
1:A:190:LYS:HG3	1:A:191:VAL:HG23	1.77	0.64
1:A:118:PHE:HD2	1:A:133:VAL:HG23	1.61	0.64
2:B:213:PRO:C	2:B:214:LYS:HE3	2.23	0.64
2:B:214:LYS:HE3	2:B:214:LYS:N	2.13	0.64
2:B:82(C):VAL:HB	2:B:111:ILE:HD13	1.79	0.63
1:A:8:PRO:O	1:A:102:THR:HB	2.01	0.61
1:A:79:ARG:HB3	1:A:81:GLU:OE2	2.01	0.60
1:A:167:ASP:HB3	1:A:170:ASP:HB2	1.84	0.60
2:B:82(C):VAL:CG2	2:B:111:ILE:HD13	2.33	0.59
1:A:29:VAL:HG11	1:A:90:GLN:HG3	1.84	0.59
2:B:213:PRO:HA	2:B:214:LYS:HE3	1.84	0.58
1:A:12:SER:HB3	1:A:140:TYR:OH	2.04	0.58
1:A:29:VAL:HG11	1:A:90:GLN:CG	2.35	0.57
1:A:90:GLN:NE2	1:A:92:HIS:N	2.51	0.57
1:A:81:GLU:H	1:A:81:GLU:CD	2.13	0.57
2:B:185:PRO:C	2:B:187:SER:H	2.13	0.57
1:A:183:LYS:O	1:A:187:GLU:HG2	2.05	0.56
2:B:135:ALA:HB1	2:B:183:THR:HG23	1.88	0.56
2:B:38:ARG:HD3	2:B:46:GLU:OE1	2.06	0.55
1:A:133:VAL:CG2	2:B:124:LEU:HD22	2.37	0.55
2:B:86:ASP:HB2	2:B:111:ILE:CD1	2.34	0.55
2:B:121:VAL:O	2:B:209:LYS:HE3	2.07	0.55
2:B:96:ARG:HD3	2:B:101:ASP:OD2	2.06	0.54
2:B:38:ARG:HB3	2:B:48:LEU:HD11	1.87	0.54
1:A:4:LEU:HD11	1:A:90:GLN:HB2	1.89	0.54
2:B:185:PRO:O	2:B:187:SER:N	2.40	0.54

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:21:ILE:HG23	1:A:102:THR:HG21	1.90	0.53
2:B:100(B):PHE:O	2:B:100(D):VAL:HG23	2.08	0.53
1:A:6:GLN:HE21	1:A:99:GLY:HA3	1.73	0.53
1:A:147:GLN:CD	1:A:154:LEU:HD11	2.34	0.52
1:A:124:GLN:HE22	1:A:131:SER:HB2	1.75	0.52
1:A:2:LEU:C	1:A:2:LEU:HD23	2.34	0.52
1:A:124:GLN:NE2	1:A:131:SER:H	2.07	0.52
1:A:147:GLN:NE2	1:A:154:LEU:HG	2.25	0.52
1:A:39:LYS:HE3	4:A:248:HOH:O	2.09	0.52
2:B:188:SER:HB3	2:B:194:TYR:OH	2.10	0.51
1:A:94:TYR:CZ	2:B:58:ARG:HG3	2.46	0.51
1:A:158:ASN:ND2	1:A:158:ASN:H	2.09	0.51
2:B:87:THR:HG22	2:B:111:ILE:H	1.75	0.51
2:B:193:THR:HG22	2:B:210:ARG:HH21	1.76	0.51
1:A:9:SER:O	1:A:102:THR:HA	2.10	0.51
1:A:155:GLN:HB3	1:A:158:ASN:HD21	1.76	0.51
1:A:118:PHE:HB2	1:A:133:VAL:CG2	2.41	0.50
2:B:84:PRO:O	2:B:87:THR:CG2	2.51	0.50
2:B:213:PRO:C	2:B:214:LYS:NZ	2.69	0.49
2:B:213:PRO:C	2:B:214:LYS:CE	2.85	0.49
1:A:143:GLU:CD	1:A:143:GLU:H	2.20	0.49
2:B:12:VAL:O	2:B:111:ILE:HA	2.13	0.49
2:B:114:THR:HG21	2:B:146:PHE:CB	2.43	0.49
1:A:175:LEU:C	1:A:175:LEU:HD23	2.38	0.48
2:B:87:THR:HB	2:B:110:THR:HA	1.95	0.48
1:A:170:ASP:HB3	1:A:172:THR:HG23	1.95	0.48
2:B:82(C):VAL:CB	2:B:111:ILE:HD13	2.43	0.48
2:B:82(C):VAL:HB	2:B:111:ILE:CD1	2.42	0.48
2:B:201:LYS:HB2	2:B:202:PRO:HD3	1.96	0.47
1:A:158:ASN:H	1:A:158:ASN:HD22	1.62	0.47
2:B:178:LEU:HD12	2:B:178:LEU:C	2.39	0.47
1:A:37:ARG:HB2	1:A:47:LEU:HD11	1.96	0.47
1:A:118:PHE:CD2	1:A:133:VAL:HG23	2.47	0.47
1:A:90:GLN:HE22	1:A:93:PHE:N	1.87	0.46
2:B:29:LEU:O	2:B:53:SER:CB	2.62	0.46
1:A:138:ASN:OD1	2:B:164:HIS:HE1	1.97	0.46
2:B:185:PRO:C	2:B:187:SER:N	2.73	0.46
2:B:13:LYS:H	2:B:16:GLN:NE2	2.13	0.46
1:A:150:VAL:O	1:A:151:ASP:C	2.60	0.45
1:A:147:GLN:CG	1:A:154:LEU:HD11	2.47	0.45
2:B:35(B):GLY:HA2	2:B:50:ILE:HA	1.98	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:213:PRO:C	2:B:214:LYS:HZ2	2.25	0.45
1:A:44:PRO:HD2	2:B:103:TRP:CE3	2.52	0.44
2:B:114:THR:CG2	2:B:115:SER:N	2.81	0.44
2:B:114:THR:CG2	2:B:115:SER:O	2.65	0.44
1:A:124:GLN:HG2	1:A:129:THR:O	2.16	0.44
1:A:201:LEU:HD13	1:A:205:VAL:HG12	1.99	0.44
1:A:206:THR:O	1:A:207:LYS:HD2	2.18	0.44
2:B:114:THR:HG21	2:B:175:LEU:HD23	1.99	0.43
1:A:201:LEU:CD1	1:A:205:VAL:HG12	2.49	0.43
1:A:133:VAL:HG21	2:B:124:LEU:CD2	2.44	0.43
1:A:158:ASN:ND2	1:A:158:ASN:N	2.66	0.43
2:B:114:THR:HG22	2:B:115:SER:O	2.19	0.42
2:B:100(H):ARG:O	2:B:100(K):VAL:HG22	2.20	0.42
2:B:28:SER:C	2:B:30:SER:H	2.27	0.42
2:B:70:THR:CG2	2:B:79:VAL:HB	2.50	0.42
2:B:100(A):LEU:HG	2:B:100(B):PHE:CD1	2.55	0.41
1:A:6:GLN:NE2	1:A:88:CYS:H	2.18	0.41
2:B:138:LEU:CD2	2:B:138:LEU:N	2.83	0.41
1:A:159:SER:HA	1:A:178:THR:O	2.21	0.41
2:B:100:THR:HA	2:B:100(D):VAL:O	2.21	0.41
2:B:87:THR:CG2	2:B:111:ILE:H	2.34	0.41
2:B:138:LEU:N	2:B:138:LEU:HD23	2.36	0.41
2:B:76:ASN:ND2	4:B:217:HOH:O	2.51	0.41
1:A:29:VAL:HG11	1:A:90:GLN:HG2	2.03	0.41
2:B:66:ARG:NH2	2:B:86:ASP:OD2	2.53	0.41
1:A:24:ARG:HA	1:A:69:THR:O	2.21	0.40
2:B:36:TRP:C	2:B:37:ILE:HD12	2.47	0.40
2:B:213:PRO:CA	2:B:214:LYS:HE3	2.48	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	212/214 (99%)	204 (96%)	8 (4%)	0	100	100
2	B	221/235 (94%)	212 (96%)	7 (3%)	2 (1%)	14	11
3	C	5/7 (71%)	5 (100%)	0	0	100	100
All	All	438/456 (96%)	421 (96%)	15 (3%)	2 (0%)	24	23

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	214	LYS
2	B	186	SER

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	187/187 (100%)	178 (95%)	9 (5%)	23	23
2	B	197/203 (97%)	186 (94%)	11 (6%)	19	18
3	C	6/6 (100%)	6 (100%)	0	100	100
All	All	390/396 (98%)	370 (95%)	20 (5%)	21	21

All (20) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	22	THR
1	A	54	LEU
1	A	74	THR
1	A	81	GLU
1	A	97	THR
1	A	102	THR
1	A	143	GLU
1	A	158	ASN
1	A	202	SER
2	B	29	LEU
2	B	87	THR

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	96	ARG
2	B	138	LEU
2	B	149	PRO
2	B	159	LEU
2	B	170	LEU
2	B	178	LEU
2	B	183	THR
2	B	195	THR
2	B	214	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (16) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	6	GLN
1	A	38	GLN
1	A	90	GLN
1	A	124	GLN
1	A	137	ASN
1	A	147	GLN
1	A	155	GLN
1	A	158	ASN
1	A	166	GLN
1	A	199	GLN
1	A	210	ASN
2	B	16	GLN
2	B	39	GLN
2	B	164	HIS
2	B	192	GLN
2	B	199	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2			OWAB(Å ²)	Q < 0.9
1	A	214/214 (100%)	0.21	9 (4%)	40	39	25, 37, 52, 58	0
2	B	227/235 (96%)	0.25	17 (7%)	20	18	23, 35, 57, 78	0
3	C	7/7 (100%)	0.23	1 (14%)	6	5	30, 33, 43, 58	0
All	All	448/456 (98%)	0.23	27 (6%)	27	25	23, 36, 56, 78	0

All (27) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	134	ALA	8.6
2	B	216	CYS	7.6
2	B	126	PRO	6.8
2	B	215	SER	6.0
2	B	133	GLY	4.7
2	B	100(B)	PHE	3.8
1	A	193	GLU	3.8
2	B	114	THR	3.5
2	B	214	LYS	3.5
1	A	170	ASP	3.4
3	C	7	SER	3.3
2	B	185	PRO	3.2
2	B	186	SER	3.2
2	B	187	SER	3.1
1	A	1	ALA	2.7
2	B	135	ALA	2.6
2	B	213	PRO	2.5
2	B	189	LEU	2.4
1	A	202	SER	2.3
1	A	214	CYS	2.3
1	A	126	LYS	2.3
1	A	154	LEU	2.2
2	B	100(C)	GLY	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	100(D)	VAL	2.2
2	B	161	SER	2.1
1	A	169	LYS	2.1
1	A	203	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.