



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 04:32 PM UTC

PDB ID : 1UA7 / pdb_00001ua7
Title : Crystal Structure Analysis of Alpha-Amylase from Bacillus Subtilis complexed with Acarbose
Authors : Kagawa, M.; Fujimoto, Z.; Momma, M.; Takase, K.; Mizuno, H.
Deposited on : 2003-03-03
Resolution : 2.21 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.21 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7682 (2.24-2.20)
Clashscore	190562	8402 (2.24-2.20)
Ramachandran outliers	187476	8303 (2.24-2.20)
Sidechain outliers	187428	8304 (2.24-2.20)
RSRZ outliers	180081	7683 (2.24-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	422	 76% 22% .
2	B	2	 100%
3	C	2	 50% 50%

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 3808 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Alpha-amylase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	422	Total	C	N	O	S	0	0	0
			3303	2054	581	660	8			

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	105	PHE	SER	SEE REMARK 999	UNP P00691
A	163	GLU	ASP	SEE REMARK 999	UNP P00691
A	356	GLN	ASN	engineered mutation	UNP P00691

- Molecule 2 is an oligosaccharide called 4,6-dideoxy-alpha-D-xylo-hexopyranose-(1-4)-alpha-D-glucopyranose.




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
2	B	2	Total	C	O	0	0	0
			21	12	9			

- Molecule 3 is an oligosaccharide called alpha-D-quinovopyranose-(1-4)-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
3	C	2	Total	C	O	0	0	0
			21	12	9			

- Molecule 4 is 6-AMINO-4-HYDROXYMETHYL-CYCLOHEX-4-ENE-1,2,3-TRIOL (CCD ID: ACI) (formula: C₇H₁₃NO₄).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			12	7	1	4		
4	A	1	Total	C	N	O	0	0
			11	7	1	3		

- Molecule 5 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	3	Total	Ca	0	0
			3	3		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	437	Total	O	0	0
			437	437		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Alpha-amylase



• Molecule 2: 4,6-dideoxy-alpha-D-xylo-hexopyranose-(1-4)-alpha-D-glucopyranose



• Molecule 3: alpha-D-quinovopyranose-(1-4)-beta-D-glucopyranose



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	70.32Å 74.20Å 115.72Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.97 – 2.21 19.97 – 2.21	Depositor EDS
% Data completeness (in resolution range)	95.5 (19.97-2.21) 95.5 (19.97-2.21)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.38 (at 2.20Å)	Xtriage
Refinement program	CNS 1.0	Depositor
R, R_{free}	0.208 , 0.265 0.201 , 0.258	Depositor DCC
R_{free} test set	2976 reflections (9.99%)	wwPDB-VP
Wilson B-factor (Å ²)	16.0	Xtriage
Anisotropy	0.964	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 66.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.44$, $\langle L^2 \rangle = 0.26$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	3808	wwPDB-VP
Average B, all atoms (Å ²)	21.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.25% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: BGC, G6D, GLC, CA, GLD, ACI

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.37	0/3383	0.90	9/4596 (0.2%)

There are no bond length outliers.

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	68	GLN	N-CA-C	-8.10	98.50	110.48
1	A	265	VAL	N-CA-C	-7.47	103.66	111.58
1	A	211	GLN	N-CA-C	6.96	120.59	110.28
1	A	65	THR	N-CA-C	-6.14	106.76	114.56
1	A	121	THR	N-CA-C	5.68	118.04	109.23
1	A	140	SER	N-CA-C	5.62	119.25	110.32
1	A	293	ILE	N-CA-C	5.58	116.36	110.72
1	A	59	TYR	N-CA-C	5.33	119.04	112.54
1	A	100	ILE	N-CA-C	5.26	117.16	112.17

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3303	0	3087	68	0
2	B	21	0	19	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	C	21	0	12	1	0
4	A	23	0	22	1	0
5	A	3	0	0	0	0
6	A	437	0	0	8	0
All	All	3808	0	3140	69	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

All (69) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:53:LYS:HE2	6:A:611:HOH:O	1.87	0.74
1:A:391:ASP:HA	1:A:405:VAL:HG23	1.76	0.68
1:A:197:ILE:C	1:A:197:ILE:HD12	2.23	0.64
1:A:127:ILE:HG13	1:A:140:SER:O	1.98	0.64
1:A:151:ASN:O	1:A:154:VAL:HG22	2.00	0.62
1:A:78:GLN:O	1:A:82:GLU:HG3	2.00	0.61
1:A:373:LEU:HB2	1:A:420:ALA:HB3	1.85	0.59
1:A:383:ILE:HG23	1:A:412:GLY:O	2.03	0.58
1:A:344:VAL:HG23	1:A:345:MET:HE2	1.87	0.57
1:A:250:SER:O	1:A:296:ARG:NH2	2.36	0.57
1:A:392:GLY:N	1:A:405:VAL:HG23	2.20	0.57
1:A:185:ASP:CG	1:A:216:ARG:HH22	2.14	0.56
1:A:221:ALA:HA	1:A:224:MET:O	2.05	0.56
1:A:14:ALA:CB	1:A:22:LEU:HD21	2.37	0.55
1:A:228:ALA:HB2	1:A:261:LEU:HG	1.90	0.54
1:A:362:PHE:HB3	1:A:374:ALA:HB3	1.90	0.53
6:A:960:HOH:O	3:C:1:BGC:H6C1	2.08	0.53
1:A:13:HIS:HB2	1:A:302:LEU:HD22	1.89	0.53
1:A:162:LEU:O	1:A:166:LEU:HD22	2.09	0.53
1:A:352:LEU:N	1:A:352:LEU:HD23	2.24	0.53
1:A:7:LYS:HE2	1:A:259:ASP:O	2.10	0.52
1:A:247:SER:HB2	6:A:705:HOH:O	2.10	0.51
1:A:390:PRO:O	1:A:405:VAL:HG21	2.11	0.50
1:A:332:ASP:OD2	1:A:334:ALA:HB3	2.11	0.50
1:A:414:ILE:HG23	1:A:414:ILE:O	2.12	0.50
1:A:394:TYR:HE2	1:A:405:VAL:HG21	1.76	0.50
1:A:384:ASN:C	1:A:384:ASN:HD22	2.20	0.49
1:A:251:HIS:CD2	1:A:253:ALA:H	2.31	0.49
1:A:259:ASP:OD1	1:A:260:LYS:HG3	2.13	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:37:ILE:HD12	1:A:37:ILE:C	2.36	0.49
1:A:340:ARG:O	1:A:344:VAL:HG22	2.13	0.48
1:A:348:GLN:HG3	1:A:367:GLY:HA3	1.96	0.48
1:A:395:ASP:HB2	6:A:880:HOH:O	2.13	0.48
1:A:195:PRO:HG3	1:A:223:TYR:CE1	2.50	0.47
1:A:391:ASP:CA	1:A:405:VAL:HG23	2.44	0.47
1:A:151:ASN:HB3	1:A:154:VAL:HG13	1.97	0.47
1:A:102:HIS:CD2	1:A:144:LEU:HD13	2.51	0.46
1:A:270:THR:HB	1:A:277:GLU:HG3	1.98	0.45
1:A:348:GLN:HG3	1:A:349:PRO:HD2	1.98	0.45
1:A:421:VAL:C	1:A:422:LEU:HD13	2.41	0.45
1:A:345:MET:O	1:A:346:ALA:C	2.60	0.45
1:A:25:ASN:O	1:A:29:ILE:HG13	2.17	0.44
1:A:271:TYR:CD1	1:A:271:TYR:C	2.94	0.44
1:A:345:MET:O	1:A:348:GLN:HB2	2.16	0.44
1:A:348:GLN:HE22	1:A:369:HIS:CE1	2.36	0.44
1:A:209:ILE:HD12	1:A:226:VAL:HG12	2.00	0.44
1:A:122:HIS:HB2	1:A:146:ASP:HB3	2.00	0.44
1:A:336:THR:O	1:A:340:ARG:HG3	2.18	0.43
1:A:405:VAL:HG23	1:A:405:VAL:O	2.18	0.43
1:A:158:LEU:O	1:A:161:PHE:HB3	2.18	0.43
1:A:192:GLN:C	1:A:195:PRO:HD2	2.43	0.43
1:A:62:TYR:HB3	4:A:504:ACI:O6	2.19	0.43
1:A:383:ILE:O	1:A:411:THR:HA	2.19	0.43
1:A:344:VAL:CG2	1:A:345:MET:HE2	2.49	0.43
1:A:396:ASN:OD1	1:A:420:ALA:HA	2.19	0.43
1:A:81:LYS:HE2	1:A:168:ASP:OD1	2.19	0.43
1:A:122:HIS:NE2	1:A:148:ASN:HA	2.34	0.42
1:A:383:ILE:HD11	1:A:410:LEU:HD23	2.01	0.42
1:A:7:LYS:HD2	1:A:300:THR:OG1	2.18	0.42
1:A:394:TYR:HE2	1:A:405:VAL:CG2	2.32	0.42
1:A:76:THR:OG1	1:A:79:GLU:HG3	2.20	0.42
1:A:248:ASN:HB2	6:A:1010:HOH:O	2.19	0.42
1:A:6:ILE:HG13	6:A:1004:HOH:O	2.19	0.42
1:A:194:TRP:CZ3	1:A:197:ILE:HD11	2.54	0.42
1:A:349:PRO:O	1:A:366:ARG:HA	2.20	0.41
1:A:377:GLY:HA3	6:A:700:HOH:O	2.20	0.41
1:A:264:TRP:C	1:A:264:TRP:CD1	2.99	0.41
1:A:296:ARG:HD2	1:A:350:GLU:OE1	2.21	0.41
1:A:35:THR:HB	6:A:889:HOH:O	2.19	0.41

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	420/422 (100%)	405 (96%)	15 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	351/351 (100%)	339 (97%)	12 (3%)	32	42

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	66	SER
1	A	166	LEU
1	A	197	ILE
1	A	225	ASP
1	A	242	ARG
1	A	261	LEU
1	A	302	LEU
1	A	348	GLN
1	A	352	LEU
1	A	356	GLN
1	A	383	ILE
1	A	422	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	112	ASN
1	A	126	GLN
1	A	251	HIS
1	A	342	HIS
1	A	348	GLN
1	A	356	GLN
1	A	384	ASN
1	A	406	ASN
1	A	415	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

4 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	GLC	B	1	2,4	12,12,12	1.47	2 (16%)	17,17,17	0.88	0
2	GLD	B	2	2,4	9,9,10	3.16	5 (55%)	11,12,14	1.27	2 (18%)
3	BGC	C	1	3	12,12,12	1.42	1 (8%)	17,17,17	0.87	0
3	G6D	C	2	3,4	9,9,11	2.76	5 (55%)	11,12,16	1.47	1 (9%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.
'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GLC	B	1	2,4	-	0/2/22/22	0/1/1/1
2	GLD	B	2	2,4	-	-	0/1/1/1
3	BGC	C	1	3	-	0/2/22/22	0/1/1/1
3	G6D	C	2	3,4	-	-	0/1/1/1

All (13) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	2	GLD	O5-C5	6.24	1.49	1.44
3	C	2	G6D	O5-C5	5.81	1.49	1.44
2	B	2	GLD	C4-C5	4.21	1.59	1.52
3	C	2	G6D	O5-C1	3.23	1.49	1.43
2	B	2	GLD	C4-C3	3.18	1.58	1.52
2	B	2	GLD	O5-C1	3.11	1.48	1.43
2	B	2	GLD	C3-C2	3.07	1.56	1.52
3	C	1	BGC	O5-C1	2.93	1.50	1.42
2	B	1	GLC	C4-C5	2.75	1.58	1.53
3	C	2	G6D	C4-C3	2.62	1.57	1.52
3	C	2	G6D	C4-C5	2.57	1.56	1.52
3	C	2	G6D	C3-C2	2.53	1.55	1.52
2	B	1	GLC	C4-C3	2.23	1.58	1.52

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	2	G6D	C1-O5-C5	3.69	117.22	111.80
2	B	2	GLD	C1-O5-C5	2.69	115.75	111.80
2	B	2	GLD	C3-C4-C5	-2.14	107.81	111.21

There are no chirality outliers.


There are no torsion outliers.


There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	C	1	BGC	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 3 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	ACI	A	501	2	12,12,12	5.63	6 (50%)	11,17,17	0.98	0
4	ACI	A	504	3,2	11,11,12	3.27	6 (54%)	8,15,17	1.54	1 (12%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.

'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	ACI	A	501	2	-	2/2/22/22	0/1/1/1
4	ACI	A	504	3,2	-	1/2/18/22	0/1/1/1

All (12) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	501	ACI	C7-C5	16.78	1.56	1.32
4	A	501	ACI	C4-C5	7.38	1.57	1.51
4	A	504	ACI	C7-C5	6.65	1.53	1.33
4	A	504	ACI	C1-C7	4.18	1.59	1.50
4	A	504	ACI	C3-C2	4.08	1.58	1.52
4	A	501	ACI	C1-C7	3.83	1.58	1.50
4	A	501	ACI	C3-C4	3.77	1.58	1.53
4	A	504	ACI	C2-C1	3.66	1.57	1.52
4	A	504	ACI	C4-C3	3.52	1.58	1.52
4	A	504	ACI	C4-C5	2.82	1.56	1.50
4	A	501	ACI	C3-C2	2.40	1.58	1.52
4	A	501	ACI	C2-C1	2.37	1.56	1.52

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	504	ACI	O6-C6-C5	-3.22	103.67	112.21

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	501	ACI	C4-C5-C6-O6
4	A	501	ACI	C7-C5-C6-O6
4	A	504	ACI	C7-C5-C6-O6

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	504	ACI	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	422/422 (100%)	-0.38	2 (0%) 87 86	9, 19, 34, 56	0

All (2) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	405	VAL	3.1
1	A	4	PRO	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	BGC	C	1	12/12	0.85	0.13	21,29,41,50	0
2	GLD	B	2	9/10	0.94	0.07	19,22,24,26	0
2	GLC	B	1	12/12	0.96	0.06	10,17,21,21	0
3	G6D	C	2	9/11	0.97	0.05	16,19,23,24	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around Chain B:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	ACI	A	501	12/12	0.94	0.07	18,24,29,34	0
4	ACI	A	504	11/12	0.94	0.07	12,15,18,19	0
5	CA	A	601	1/1	0.99	0.02	13,13,13,13	0
5	CA	A	602	1/1	0.99	0.02	19,19,19,19	0
5	CA	A	603	1/1	0.99	0.03	11,11,11,11	0

6.5 Other polymers [i](#)

There are no such residues in this entry.