



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 04:50 AM UTC

PDB ID : 4UA2 / pdb_00004ua2
Title : Crystal structure of dual function transcriptional regulator MerR from *Bacillus megaterium* MB1
Authors : Lin, L.Y.; Chang, C.C.; Zou, X.W.; Huang, C.C.; Chan, N.L.
Deposited on : 2014-08-07
Resolution : 2.61 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.61 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4951 (2.64-2.60)
Clashscore	190562	5303 (2.64-2.60)
Ramachandran outliers	187476	5217 (2.64-2.60)
Sidechain outliers	187428	5217 (2.64-2.60)
RSRZ outliers	180081	4950 (2.64-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	132	8% 80% 12% 8%
1	B	132	4% 75% 12% 12%
1	C	132	14% 62% 11% 27%
1	D	132	9% 70% 15% 14%
1	E	132	13% 73% 17% 10%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	132	11% 76% 14% 11%
1	G	132	8% 70% 16% • 13%
1	H	132	11% 67% 14% • 18%

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 7263 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Regulatory protein.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	121	Total	C	N	O	S	Se	0	0	0
			939	596	156	177	4	6			
1	G	115	Total	C	N	O	S	Se	0	0	0
			905	575	150	171	4	5			
1	E	119	Total	C	N	O	S	Se	0	0	0
			944	598	160	176	4	6			
1	C	96	Total	C	N	O	S	Se	0	0	0
			733	465	123	137	3	5			
1	F	118	Total	C	N	O	S	Se	0	0	0
			942	597	160	176	4	5			
1	B	116	Total	C	N	O	S	Se	0	0	0
			943	595	161	179	3	5			
1	D	113	Total	C	N	O	S	Se	0	0	0
			895	569	151	167	3	5			
1	H	108	Total	C	N	O	S	Se	0	0	0
			865	552	142	163	3	5			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MSE	-	initiating methionine	UNP Q799U3
G	1	MSE	-	initiating methionine	UNP Q799U3
E	1	MSE	-	initiating methionine	UNP Q799U3
C	1	MSE	-	initiating methionine	UNP Q799U3
F	1	MSE	-	initiating methionine	UNP Q799U3
B	1	MSE	-	initiating methionine	UNP Q799U3
D	1	MSE	-	initiating methionine	UNP Q799U3
H	1	MSE	-	initiating methionine	UNP Q799U3


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	14	Total 14	O 14	0	0
2	G	16	Total 16	O 16	0	0
2	E	17	Total 17	O 17	0	0
2	C	7	Total 7	O 7	0	0
2	F	16	Total 16	O 16	0	0
2	B	13	Total 13	O 13	0	0
2	D	5	Total 5	O 5	0	0
2	H	9	Total 9	O 9	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Regulatory protein




- Molecule 1: Regulatory protein




- Molecule 1: Regulatory protein




- Molecule 1: Regulatory protein




- Molecule 1: Regulatory protein




- Molecule 1: Regulatory protein



- Molecule 1: Regulatory protein



- Molecule 1: Regulatory protein



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	80.54Å 94.81Å 87.06Å 90.00° 101.83° 90.00°	Depositor
Resolution (Å)	28.41 – 2.61 28.41 – 2.62	Depositor EDS
% Data completeness (in resolution range)	99.1 (28.41-2.61) 98.9 (28.41-2.62)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.22 (at 2.61Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.9_1692)	Depositor
R, R_{free}	0.225 , 0.257 0.230 , 0.258	Depositor DCC
R_{free} test set	1926 reflections (4.95%)	wwPDB-VP
Wilson B-factor (Å ²)	48.0	Xtriage
Anisotropy	0.045	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 56.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	7263	wwPDB-VP
Average B, all atoms (Å ²)	58.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.07% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.26	0/945	0.70	0/1263
1	B	0.28	0/946	0.70	0/1256
1	C	0.30	0/733	0.75	0/976
1	D	0.26	0/898	0.73	0/1194
1	E	0.28	0/949	0.80	1/1264 (0.1%)
1	F	0.29	0/948	0.74	0/1265
1	G	0.32	0/911	0.83	3/1216 (0.2%)
1	H	0.30	0/868	0.71	0/1156
All	All	0.28	0/7198	0.74	4/9590 (0.0%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	E	31	PRO	N-CA-CB	8.64	110.22	103.30
1	G	80	ARG	CA-C-N	7.99	136.08	121.70
1	G	80	ARG	C-N-CA	7.99	136.08	121.70
1	G	44	GLN	N-CA-C	-5.88	105.60	112.89

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	939	0	900	12	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	943	0	942	13	0
1	C	733	0	695	13	0
1	D	895	0	883	15	0
1	E	944	0	926	21	0
1	F	942	0	922	13	0
1	G	905	0	873	18	0
1	H	865	0	852	15	0
2	A	14	0	0	0	0
2	B	13	0	0	0	0
2	C	7	0	0	1	0
2	D	5	0	0	0	0
2	E	17	0	0	1	0
2	F	16	0	0	0	0
2	G	16	0	0	0	0
2	H	9	0	0	0	0
All	All	7263	0	6993	92	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (92) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:G:80:ARG:HB3	1:G:81:ASP:HB2	1.60	0.82
1:A:98:GLU:OE2	1:C:102:ARG:NH1	2.20	0.75
1:C:47:ASP:OD2	1:F:130:MSE:HG2	1.86	0.74
1:B:113:ARG:NH2	1:H:83:TYR:OH	2.25	0.69
1:G:22:TYR:HB3	1:G:27:LEU:HB2	1.72	0.69
1:C:99:ASP:OD1	1:C:102:ARG:NH2	2.26	0.68
1:A:12:CYS:HB2	1:A:13:GLY:HA2	1.76	0.68
1:G:60:PHE:HZ	1:G:93:ILE:HD11	1.59	0.66
1:F:60:PHE:HZ	1:F:93:ILE:HD11	1.61	0.65
1:G:99:ASP:OD1	1:G:102:ARG:NH2	2.31	0.64
1:E:80:ARG:HD2	1:E:81:ASP:H	1.63	0.64
1:E:40:MSE:HA	1:E:42:SER:H	1.63	0.63
1:D:99:ASP:OD1	1:D:102:ARG:NH2	2.33	0.62
1:C:104:GLU:OE2	1:D:94:GLN:NE2	2.33	0.61
1:G:80:ARG:HB3	1:G:81:ASP:CB	2.29	0.60
1:B:83:TYR:OH	1:H:113:ARG:NH2	2.33	0.60
1:F:99:ASP:OD1	1:F:102:ARG:NH2	2.36	0.59
1:C:94:GLN:NE2	1:D:104:GLU:OE2	2.36	0.59

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:H:99:ASP:OD1	1:H:102:ARG:NH2	2.37	0.58
1:A:96:LYS:HE3	1:B:100:LEU:HD11	1.84	0.58
1:E:40:MSE:HA	1:E:42:SER:N	2.19	0.57
1:E:51:PHE:HB2	1:H:126:ILE:HG23	1.87	0.56
1:E:102:ARG:NH1	1:D:98:GLU:OE1	2.38	0.56
1:H:60:PHE:HZ	1:H:93:ILE:HD11	1.71	0.56
1:E:7:GLU:HA	1:E:10:ASP:HB3	1.88	0.56
1:A:99:ASP:OD1	1:A:102:ARG:NH2	2.39	0.55
1:E:99:ASP:OD1	1:E:102:ARG:NH2	2.38	0.55
1:E:83:TYR:OH	1:D:113:ARG:NH2	2.33	0.55
1:A:82:MSE:HE1	1:B:110:LEU:HD13	1.87	0.55
1:E:80:ARG:HH11	1:E:81:ASP:N	2.04	0.54
1:E:102:ARG:HH12	1:D:98:GLU:CD	2.14	0.54
1:G:83:TYR:OH	1:F:113:ARG:NH2	2.38	0.54
1:A:94:GLN:NE2	1:B:104:GLU:OE2	2.39	0.53
1:E:87:ILE:HD12	1:F:124:PRO:HG3	1.89	0.53
1:D:83:TYR:O	1:D:86:THR:OG1	2.26	0.52
1:G:12:CYS:HA	1:G:15:ASN:OD1	2.11	0.50
1:A:54:ARG:HB3	1:G:129:LEU:HD13	1.92	0.50
1:G:80:ARG:HB3	1:G:81:ASP:C	2.37	0.50
1:H:10:ASP:O	1:H:11:LYS:HG2	2.12	0.50
1:C:16:LYS:N	2:C:205:HOH:O	2.45	0.50
1:C:124:PRO:HG3	1:D:87:ILE:HD12	1.94	0.50
1:F:10:ASP:CG	1:F:11:LYS:N	2.69	0.49
1:B:109:ASP:OD1	1:B:113:ARG:NH1	2.45	0.49
1:A:87:ILE:HD13	1:B:124:PRO:HB3	1.95	0.49
1:G:28:ILE:HD11	1:G:49:LEU:HD21	1.95	0.48
1:G:80:ARG:HB3	1:G:81:ASP:CA	2.42	0.48
1:G:99:ASP:OD2	1:H:54:ARG:NH2	2.38	0.48
1:E:80:ARG:HD2	1:E:81:ASP:N	2.28	0.48
1:F:127:GLU:HA	1:F:130:MSE:HG3	1.95	0.48
1:B:54:ARG:HB3	1:D:129:LEU:HD13	1.95	0.48
1:H:8:LEU:HD12	1:H:8:LEU:O	2.12	0.48
1:E:67:LYS:HE3	1:E:82:MSE:SE	2.64	0.48
1:A:108:MSE:O	1:A:112:GLU:HG3	2.13	0.47
1:D:20:ARG:O	1:D:21:TYR:HB3	2.13	0.47
1:D:48:ARG:NH2	1:D:73:ASP:OD1	2.47	0.47
1:C:67:LYS:HE3	1:C:78:LYS:O	2.16	0.46
1:F:10:ASP:OD2	1:F:11:LYS:N	2.48	0.46
1:C:87:ILE:HD13	1:D:124:PRO:HB3	1.96	0.46
1:C:130:MSE:HE1	1:F:50:HIS:HB3	1.99	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:G:41:TYR:HD2	1:G:41:TYR:O	1.99	0.45
1:H:8:LEU:CD1	1:H:49:LEU:HD22	2.47	0.45
1:A:28:ILE:HG22	1:A:48:ARG:HD3	1.99	0.44
1:B:83:TYR:O	1:B:86:THR:OG1	2.28	0.44
1:G:89:LYS:HA	1:G:89:LYS:HE2	1.99	0.44
1:E:12:CYS:HA	1:E:13:GLY:HA2	1.67	0.44
1:G:42:SER:C	1:G:44:GLN:H	2.25	0.44
1:E:83:TYR:O	1:E:86:THR:OG1	2.30	0.44
1:H:60:PHE:CZ	1:H:93:ILE:HD11	2.51	0.43
1:G:113:ARG:NH2	1:F:83:TYR:OH	2.47	0.43
1:E:41:TYR:O	1:E:42:SER:HB3	2.18	0.43
1:H:80:ARG:HA	1:H:81:ASP:HA	1.68	0.43
1:E:121:TYR:OH	1:H:76:GLU:OE1	2.36	0.43
1:B:10:ASP:O	1:B:11:LYS:HB2	2.18	0.43
1:A:129:LEU:HD13	1:G:54:ARG:HB3	2.01	0.42
1:C:82:MSE:HE2	1:D:110:LEU:HD23	1.99	0.42
1:F:10:ASP:C	1:F:12:CYS:H	2.27	0.42
1:H:79:CYS:O	1:H:82:MSE:HG2	2.20	0.42
1:E:40:MSE:N	2:E:201:HOH:O	2.52	0.42
1:D:79:CYS:O	1:D:82:MSE:HB2	2.20	0.42
1:G:79:CYS:HA	1:G:82:MSE:HE2	1.99	0.42
1:E:14:VAL:HG23	1:E:53:LYS:HD2	2.02	0.42
1:E:40:MSE:HA	1:E:41:TYR:HB2	2.02	0.41
1:H:83:TYR:O	1:H:87:ILE:HG12	2.21	0.41
1:C:129:LEU:HD23	1:C:129:LEU:HA	1.75	0.41
1:H:83:TYR:O	1:H:86:THR:OG1	2.29	0.41
1:B:123:CYS:O	1:B:125:ILE:HD12	2.20	0.41
1:D:27:LEU:O	1:D:48:ARG:NH1	2.50	0.41
1:A:82:MSE:HE2	1:B:110:LEU:HD22	2.02	0.41
1:B:4:ARG:HE	1:B:41:TYR:HA	1.85	0.41
1:E:114:CYS:HA	1:E:115:PRO:HD3	1.94	0.41
1:F:28:ILE:HD13	1:F:48:ARG:HD3	2.04	0.40
1:C:130:MSE:HE3	1:F:47:ASP:HB3	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	117/132 (89%)	112 (96%)	5 (4%)	0	100	100
1	B	110/132 (83%)	107 (97%)	2 (2%)	1 (1%)	14	28
1	C	92/132 (70%)	91 (99%)	1 (1%)	0	100	100
1	D	107/132 (81%)	104 (97%)	3 (3%)	0	100	100
1	E	115/132 (87%)	110 (96%)	5 (4%)	0	100	100
1	F	114/132 (86%)	109 (96%)	5 (4%)	0	100	100
1	G	111/132 (84%)	105 (95%)	6 (5%)	0	100	100
1	H	102/132 (77%)	97 (95%)	5 (5%)	0	100	100
All	All	868/1056 (82%)	835 (96%)	32 (4%)	1 (0%)	48	69

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	11	LYS

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	95/117 (81%)	94 (99%)	1 (1%)	65	83
1	B	102/117 (87%)	100 (98%)	2 (2%)	48	73
1	C	71/117 (61%)	70 (99%)	1 (1%)	59	80
1	D	93/117 (80%)	92 (99%)	1 (1%)	65	83

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	E	98/117 (84%)	98 (100%)	0	100	100
1	F	98/117 (84%)	96 (98%)	2 (2%)	48	73
1	G	93/117 (80%)	92 (99%)	1 (1%)	65	83
1	H	91/117 (78%)	89 (98%)	2 (2%)	45	70
All	All	741/936 (79%)	731 (99%)	10 (1%)	61	81

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	7	GLU
1	G	120	ILE
1	C	58	LEU
1	F	44	GLN
1	F	106	MSE
1	B	17	GLU
1	B	106	MSE
1	D	76	GLU
1	H	10	ASP
1	H	95	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	50	HIS
1	C	94	GLN
1	F	50	HIS
1	D	43	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	115/132 (87%)	0.48	10 (8%) 16 12	30, 52, 89, 103	0
1	B	110/132 (83%)	0.38	5 (4%) 38 33	32, 50, 81, 100	0
1	C	91/132 (68%)	0.98	18 (19%) 3 2	34, 64, 125, 144	0
1	D	108/132 (81%)	0.69	12 (11%) 10 8	34, 54, 100, 121	0
1	E	113/132 (85%)	0.71	17 (15%) 5 4	32, 54, 93, 129	0
1	F	112/132 (84%)	0.78	15 (13%) 7 5	30, 56, 103, 131	0
1	G	109/132 (82%)	0.53	10 (9%) 14 11	31, 50, 103, 128	0
1	H	103/132 (78%)	0.69	14 (13%) 7 5	30, 51, 96, 118	0
All	All	861/1056 (81%)	0.65	101 (11%) 9 7	30, 53, 101, 144	0

All (101) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	F	43	GLN	7.8
1	F	42	SER	6.4
1	B	41	TYR	6.4
1	E	6	GLY	6.4
1	G	21	TYR	6.1
1	H	17	GLU	5.9
1	E	12	CYS	5.6
1	C	18	THR	5.6
1	C	47	ASP	5.1
1	F	9	ALA	4.8
1	G	41	TYR	4.6
1	E	32	GLU	4.5
1	H	9	ALA	4.5
1	H	16	LYS	4.5
1	D	23	GLU	4.2
1	F	11	LYS	4.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	12	CYS	4.0
1	C	17	GLU	4.0
1	E	33	ARG	4.0
1	C	22	TYR	4.0
1	H	8	LEU	3.9
1	D	42	SER	3.9
1	F	5	ILE	3.9
1	A	6	GLY	3.9
1	A	8	LEU	3.8
1	E	75	ASP	3.8
1	H	80	ARG	3.8
1	C	44	GLN	3.8
1	D	15	ASN	3.5
1	A	4	ARG	3.5
1	F	8	LEU	3.4
1	E	31	PRO	3.4
1	G	19	ILE	3.3
1	B	13	CYS	3.3
1	E	41	TYR	3.2
1	E	8	LEU	3.2
1	G	12	CYS	3.2
1	C	66	ASP	3.1
1	G	13	GLY	3.1
1	C	24	ARG	3.1
1	D	4	ARG	3.1
1	C	75	ASP	3.0
1	H	12	CYS	3.0
1	E	29	PRO	3.0
1	D	10	ASP	3.0
1	F	10	ASP	2.9
1	B	10	ASP	2.9
1	A	92	ASP	2.9
1	G	81	ASP	2.9
1	F	41	TYR	2.9
1	H	131	LYS	2.8
1	F	44	GLN	2.8
1	H	10	ASP	2.8
1	H	18	THR	2.8
1	F	15	ASN	2.7
1	A	20	ARG	2.7
1	B	119	ASP	2.7
1	D	18	THR	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	49	LEU	2.6
1	C	78	LYS	2.6
1	C	50	HIS	2.6
1	G	9	ALA	2.6
1	E	42	SER	2.6
1	C	19	ILE	2.6
1	H	49	LEU	2.6
1	F	75	ASP	2.5
1	E	13	GLY	2.5
1	G	24	ARG	2.5
1	D	8	LEU	2.5
1	E	47	ASP	2.5
1	D	30	GLU	2.5
1	G	20	ARG	2.4
1	C	53	LYS	2.4
1	F	45	THR	2.4
1	C	84	ASP	2.4
1	C	69	LEU	2.4
1	H	30	GLU	2.4
1	H	19	ILE	2.3
1	C	45	THR	2.3
1	A	33	ARG	2.3
1	D	9	ALA	2.2
1	C	23	GLU	2.2
1	E	119	ASP	2.2
1	B	42	SER	2.2
1	D	29	PRO	2.2
1	D	28	ILE	2.2
1	H	81	ASP	2.2
1	E	80	ARG	2.2
1	G	10	ASP	2.2
1	C	46	VAL	2.2
1	A	18	THR	2.1
1	F	12	CYS	2.1
1	E	9	ALA	2.1
1	F	16	LYS	2.1
1	E	44	GLN	2.1
1	H	21	TYR	2.1
1	A	10	ASP	2.0
1	E	81	ASP	2.0
1	A	14	VAL	2.0
1	F	80	ARG	2.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	71	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.