



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2026 – 01:48 AM UTC

PDB ID : 4UAH / pdb_00004uah
Title : Structure of the Ssl1 laccase mutant H99N with depleted type-2 copper ion
Authors : Gunne, M.; Hoepfner, A.; Jaeger, V.D.; Urlacher, V.B.
Deposited on : 2014-08-09
Resolution : 1.73 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.73 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1187 (1.74-1.74)
Clashscore	190562	1207 (1.74-1.74)
Ramachandran outliers	187476	1200 (1.74-1.74)
Sidechain outliers	187428	1200 (1.74-1.74)
RSRZ outliers	180081	1188 (1.74-1.74)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	293	78% 14% • 7%
1	B	293	2% 79% 12% • 8%
1	C	293	2% 80% 14% • 6%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 6978 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Copper oxidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	272	Total	C	N	O	S	0	4	0
			2135	1334	387	404	10			
1	B	271	Total	C	N	O	S	0	4	0
			2126	1328	389	399	10			
1	C	276	Total	C	N	O	S	0	6	0
			2167	1356	392	409	10			

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	33	MET	-	initiating methionine	UNP B5HSR1
A	34	HIS	-	expression tag	UNP B5HSR1
A	35	HIS	-	expression tag	UNP B5HSR1
A	36	HIS	-	expression tag	UNP B5HSR1
A	37	HIS	-	expression tag	UNP B5HSR1
A	38	HIS	-	expression tag	UNP B5HSR1
A	39	HIS	-	expression tag	UNP B5HSR1
A	99	ASN	HIS	engineered mutation	UNP B5HSR1
B	33	MET	-	initiating methionine	UNP B5HSR1
B	34	HIS	-	expression tag	UNP B5HSR1
B	35	HIS	-	expression tag	UNP B5HSR1
B	36	HIS	-	expression tag	UNP B5HSR1
B	37	HIS	-	expression tag	UNP B5HSR1
B	38	HIS	-	expression tag	UNP B5HSR1
B	39	HIS	-	expression tag	UNP B5HSR1
B	99	ASN	HIS	engineered mutation	UNP B5HSR1
C	33	MET	-	initiating methionine	UNP B5HSR1
C	34	HIS	-	expression tag	UNP B5HSR1
C	35	HIS	-	expression tag	UNP B5HSR1
C	36	HIS	-	expression tag	UNP B5HSR1
C	37	HIS	-	expression tag	UNP B5HSR1
C	38	HIS	-	expression tag	UNP B5HSR1
C	39	HIS	-	expression tag	UNP B5HSR1

Continued on next page...


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	99	ASN	HIS	engineered mutation	UNP B5HSR1

- Molecule 2 is COPPER (II) ION (CCD ID: CU) (formula: Cu).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	5	Total Cu 5 5	0	0
2	B	3	Total Cu 3 3	0	0
2	C	1	Total Cu 1 1	0	0

- Molecule 3 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	B	1	Total C O 6 3 3	0	0
3	B	1	Total C O 6 3 3	0	0
3	C	1	Total C O 6 3 3	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	186	Total 186	O 186	0	0
4	B	149	Total 149	O 149	0	0
4	C	188	Total 188	O 188	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Copper oxidase



• Molecule 1: Copper oxidase



• Molecule 1: Copper oxidase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	51.59Å 104.33Å 163.64Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	54.55 – 1.73 54.55 – 1.73	Depositor EDS
% Data completeness (in resolution range)	99.8 (54.55-1.73) 99.8 (54.55-1.73)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.10 (at 1.73Å)	Xtriage
Refinement program	REFMAC 5.8.0071	Depositor
R, R_{free}	0.170 , 0.207 (Not available) , 0.195	Depositor DCC
R_{free} test set	4562 reflections (4.90%)	wwPDB-VP
Wilson B-factor (Å ²)	12.9	Xtriage
Anisotropy	0.017	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 36.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	6978	wwPDB-VP
Average B, all atoms (Å ²)	15.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.37% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, CU

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.74	18/2206 (0.8%)	1.53	14/2993 (0.5%)
1	B	1.48	16/2196 (0.7%)	1.39	7/2978 (0.2%)
1	C	1.60	19/2245 (0.8%)	1.40	12/3048 (0.4%)
All	All	1.61	53/6647 (0.8%)	1.44	33/9019 (0.4%)

All (53) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	215	ARG	C-N	31.44	1.73	1.33
1	A	202	ALA	C-N	-15.38	1.12	1.33
1	A	56	ALA	C-N	11.86	1.50	1.33
1	A	44	GLU	C-N	-10.30	1.19	1.33
1	A	297	GLY	N-CA	7.92	1.52	1.45
1	B	200	ARG	C-N	-7.13	1.24	1.33
1	A	293	MET	C-N	-7.11	1.23	1.33
1	C	221	ILE	C-O	6.93	1.31	1.24
1	A	231	HIS	N-CA	6.89	1.54	1.46
1	A	292	ASP	C-N	-6.83	1.23	1.33
1	C	279	GLY	N-CA	-6.68	1.38	1.45
1	A	266	PHE	C-O	6.62	1.32	1.23
1	B	159	THR	C-N	-6.58	1.23	1.33
1	B	71	ILE	N-CA	6.50	1.50	1.45
1	B	251	PRO	CA-C	6.33	1.58	1.52
1	C	258	LYS	C-O	-6.26	1.17	1.23
1	B	77	GLU	N-CA	6.15	1.53	1.46
1	C	75	LEU	C-O	6.11	1.31	1.23
1	B	292	ASP	N-CA	6.04	1.53	1.46
1	C	283	TYR	C-O	5.88	1.31	1.23
1	B	238	ALA	CA-C	-5.86	1.45	1.52
1	C	276	VAL	N-CA	-5.84	1.41	1.46
1	C	239	ASP	CA-C	-5.68	1.47	1.53

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	169	GLY	C-O	5.65	1.31	1.23
1	A	96	ALA	N-CA	5.62	1.53	1.45
1	A	310	GLY	C-O	-5.57	1.18	1.23
1	B	136[A]	ARG	C-O	5.55	1.30	1.23
1	B	136[B]	ARG	C-O	5.55	1.30	1.23
1	B	267	GLY	N-CA	5.53	1.50	1.45
1	C	300	LEU	N-CA	5.53	1.53	1.46
1	C	189	THR	C-O	5.48	1.30	1.24
1	A	307	THR	N-CA	5.47	1.52	1.45
1	B	218	PHE	N-CA	5.45	1.52	1.46
1	A	259	ILE	C-O	5.44	1.30	1.24
1	B	103	LEU	C-O	5.39	1.31	1.23
1	C	311	TYR	C-N	5.37	1.40	1.33
1	C	198	ASN	CA-C	5.37	1.60	1.53
1	C	295	MET	C-O	5.35	1.29	1.23
1	A	86	ILE	N-CA	5.35	1.52	1.46
1	C	114	LEU	C-O	5.30	1.30	1.24
1	C	111	GLY	N-CA	5.23	1.52	1.45
1	C	238	ALA	C-O	-5.23	1.17	1.23
1	B	199	ASN	C-N	-5.19	1.25	1.33
1	A	285	CYS	C-O	5.17	1.30	1.23
1	A	198	ASN	N-CA	5.14	1.54	1.46
1	B	236	ARG	C-O	5.09	1.30	1.23
1	B	251	PRO	C-O	5.09	1.28	1.23
1	C	276	VAL	CA-C	5.08	1.59	1.52
1	C	114	LEU	N-CA	5.06	1.52	1.46
1	C	184	PRO	C-O	5.05	1.29	1.23
1	A	128	THR	CA-C	5.04	1.58	1.52
1	C	71	ILE	N-CA	5.04	1.49	1.45
1	A	135	GLY	N-CA	5.03	1.51	1.45

All (33) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	215	ARG	O-C-N	28.01	155.15	123.22
1	B	200	ARG	O-C-N	-18.38	100.84	121.43
1	A	215	ARG	CA-C-N	-14.83	103.71	123.14
1	A	215	ARG	C-N-CA	-14.83	103.71	123.14
1	B	200	ARG	CA-C-N	12.09	132.22	119.89
1	B	200	ARG	C-N-CA	12.09	132.22	119.89
1	C	130	ARG	NE-CZ-NH2	-11.46	108.88	119.20
1	C	130	ARG	NE-CZ-NH1	8.69	130.19	121.50

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	199	ASN	CA-C-N	7.32	130.88	120.49
1	B	199	ASN	C-N-CA	7.32	130.88	120.49
1	C	130	ARG	CD-NE-CZ	7.16	134.42	124.40
1	C	247	GLY	CA-C-N	6.99	127.14	119.87
1	C	247	GLY	C-N-CA	6.99	127.14	119.87
1	A	43	GLY	CA-C-N	6.62	132.39	121.39
1	A	43	GLY	C-N-CA	6.62	132.39	121.39
1	A	214	ASP	CA-C-N	6.51	131.43	122.77
1	A	214	ASP	C-N-CA	6.51	131.43	122.77
1	C	174	VAL	CB-CA-C	-6.45	100.10	110.28
1	C	271	ILE	N-CA-C	-6.38	98.67	107.99
1	A	44	GLU	CA-C-N	-6.25	114.90	122.90
1	A	44	GLU	C-N-CA	-6.25	114.90	122.90
1	A	308	ILE	CB-CA-C	6.14	115.34	109.33
1	B	241	ARG	N-CA-C	6.09	117.59	111.07
1	A	174	VAL	CB-CA-C	-6.00	101.58	110.33
1	B	195	MET	CG-SD-CE	-5.98	87.74	100.90
1	C	161	HIS	CA-CB-CG	-5.77	108.03	113.80
1	C	228	HIS	CA-C-N	-5.60	114.91	122.86
1	C	228	HIS	C-N-CA	-5.60	114.91	122.86
1	A	43	GLY	O-C-N	-5.42	114.33	123.00
1	C	312	GLU	O-C-N	5.28	125.51	121.23
1	A	56	ALA	CA-C-N	-5.27	113.49	122.56
1	A	56	ALA	C-N-CA	-5.27	113.49	122.56
1	C	130	ARG	CG-CD-NE	-5.08	100.81	112.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2135	0	2015	26	0
1	B	2126	0	2021	20	0
1	C	2167	0	2056	14	0
2	A	5	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	3	0	0	0	0
2	C	1	0	0	0	0
3	B	12	0	16	0	0
3	C	6	0	8	3	0
4	A	186	0	0	10	1
4	B	149	0	0	3	0
4	C	188	0	0	7	0
All	All	6978	0	6116	62	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (62) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:215:ARG:C	1:A:216:VAL:N	1.73	1.44
1:A:215:ARG:CA	1:A:216:VAL:N	2.15	1.09
1:A:215:ARG:HA	1:A:216:VAL:N	1.82	0.94
1:A:312:GLU:OE1	4:A:501:HOH:O	1.88	0.91
1:B:200:ARG:HD3	1:B:206:PRO:HD3	1.54	0.90
1:A:215:ARG:C	1:A:216:VAL:CA	2.52	0.82
1:A:253[A]:GLN:HG3	4:A:676:HOH:O	1.81	0.80
4:A:666:HOH:O	1:B:244:MET:HG2	1.84	0.78
1:A:215:ARG:HD3	1:A:269:GLN:OE1	1.82	0.78
3:C:401:GOL:O3	4:C:674:HOH:O	2.02	0.77
1:C:124:THR:HB	4:C:665:HOH:O	1.86	0.74
1:A:203:HIS:CE1	1:A:294:GLY:HA2	2.24	0.71
3:C:401:GOL:H12	4:C:688:HOH:O	1.90	0.70
1:A:185:ASP:N	1:A:215:ARG:O	2.25	0.69
3:C:401:GOL:C1	4:C:688:HOH:O	2.40	0.69
1:A:168:LYS:HG2	4:A:652:HOH:O	1.94	0.66
1:A:225[B]:GLU:OE2	4:A:670:HOH:O	2.15	0.60
1:A:195:MET:HE1	1:A:293:MET:HE3	1.85	0.58
1:B:200:ARG:NH2	1:B:204:SER:O	2.33	0.58
1:B:47:ARG:CG	4:B:606:HOH:O	2.52	0.57
1:A:195:MET:CE	1:A:293:MET:HE3	2.35	0.56
1:A:65:GLU:HB2	1:A:68:ARG:HG3	1.87	0.55
1:A:47:ARG:HD3	1:A:87:GLU:CD	2.32	0.55
1:B:47:ARG:HG2	4:B:606:HOH:O	2.07	0.53
1:B:65[B]:GLU:HB2	1:B:68[B]:ARG:HB2	1.91	0.53
1:A:215:ARG:C	1:A:216:VAL:HA	2.34	0.52

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:168:LYS:HD3	4:A:652:HOH:O	2.11	0.50
1:A:282:MET:HG3	1:C:114:LEU:HD22	1.93	0.50
1:B:102:GLY:HA3	1:B:150:TRP:CD2	2.46	0.50
1:A:244:MET:HE2	4:A:684:HOH:O	2.12	0.49
1:B:113:THR:OG1	1:B:160:GLU:OE1	2.31	0.49
1:C:130:ARG:HD3	4:C:668:HOH:O	2.13	0.49
4:A:666:HOH:O	1:B:244:MET:CG	2.53	0.48
1:C:284:HIS:HB3	1:C:296[B]:VAL:HG12	1.95	0.48
1:A:244:MET:CE	4:A:684:HOH:O	2.62	0.47
1:B:200:ARG:NE	1:B:204:SER:O	2.48	0.46
1:B:200:ARG:CD	1:B:206:PRO:HD3	2.35	0.46
1:C:75:LEU:HA	1:C:173:PRO:HG2	1.98	0.45
1:C:203:HIS:CE1	1:C:294:GLY:HA2	2.51	0.45
1:C:229:THR:O	1:C:285:CYS:HA	2.17	0.45
1:A:229:THR:O	1:A:285:CYS:HA	2.16	0.44
1:B:241:ARG:HB2	4:C:640:HOH:O	2.16	0.44
1:A:242:THR:HB	1:A:244:MET:HE3	1.99	0.44
1:B:179:LYS:HE3	1:B:179:LYS:HB2	1.75	0.42
1:B:217:GLU:HB2	1:B:269:GLN:HG2	2.01	0.42
1:A:185:ASP:HB2	1:A:215:ARG:O	2.19	0.42
1:B:229:THR:O	1:B:285:CYS:HA	2.18	0.42
1:B:47:ARG:HG3	4:B:606:HOH:O	2.18	0.42
1:B:75:LEU:HA	1:B:173:PRO:HG2	2.01	0.42
1:C:75:LEU:HG	1:C:77[B]:GLU:HG3	2.02	0.42
1:C:284:HIS:CB	1:C:296[B]:VAL:HG12	2.50	0.42
1:B:65[B]:GLU:HA	1:B:65[B]:GLU:OE1	2.19	0.42
1:C:289:SER:O	1:C:293:MET:HG3	2.19	0.41
1:B:283:TYR:CZ	1:B:297:GLY:HA3	2.55	0.41
1:A:50:LEU:O	1:A:88:PHE:HA	2.21	0.41
1:C:215:ARG:HD3	1:C:269:GLN:OE1	2.20	0.41
1:A:289:SER:HB2	1:A:293:MET:HE2	2.02	0.41
1:B:238:ALA:O	1:B:243:GLY:HA2	2.21	0.41
1:C:200:ARG:HD3	1:C:206:PRO:HD3	2.03	0.41
1:A:168:LYS:CD	4:A:652:HOH:O	2.69	0.40
1:C:253:GLN:HA	4:C:640:HOH:O	2.21	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:527:HOH:O	4:A:529:HOH:O[4_445]	2.12	0.08

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	274/293 (94%)	266 (97%)	8 (3%)	0	100	100
1	B	273/293 (93%)	263 (96%)	10 (4%)	0	100	100
1	C	280/293 (96%)	276 (99%)	4 (1%)	0	100	100
All	All	827/879 (94%)	805 (97%)	22 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	223/236 (94%)	222 (100%)	1 (0%)	84	79
1	B	222/236 (94%)	216 (97%)	6 (3%)	39	16
1	C	228/236 (97%)	225 (99%)	3 (1%)	61	43
All	All	673/708 (95%)	663 (98%)	10 (2%)	63	38

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	251	PRO
1	B	136[A]	ARG
1	B	136[B]	ARG
1	B	167	ARG
1	B	179	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	209	GLU
1	B	220	MET
1	C	212[A]	VAL
1	C	212[B]	VAL
1	C	220	MET

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	203	HIS
1	B	253	GLN
1	C	203	HIS
1	C	288	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 12 ligands modelled in this entry, 9 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	GOL	B	401	-	5,5,5	1.09	0	5,5,5	0.74	0
3	GOL	C	401	-	5,5,5	0.94	0	5,5,5	1.39	1 (20%)
3	GOL	B	402	-	5,5,5	0.82	0	5,5,5	1.49	1 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	B	401	-	-	2/4/4/4	-
3	GOL	C	401	-	-	0/4/4/4	-
3	GOL	B	402	-	-	0/4/4/4	-

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	401	GOL	O2-C2-C1	2.56	119.76	109.18
3	B	402	GOL	O2-C2-C1	2.25	118.51	109.18

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	401	GOL	C1-C2-C3-O3
3	B	401	GOL	O2-C2-C3-O3

There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	C	401	GOL	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	A	3

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	A	215:ARG	C	216:VAL	N	1.73
1	A	44:GLU	C	45:VAL	N	1.19
1	A	202:ALA	C	203:HIS	N	1.12

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å²)	Q<0.9
1	A	272/293 (92%)	-0.23	4 (1%) 72 79	7, 14, 29, 48	4 (1%)
1	B	271/293 (92%)	-0.05	7 (2%) 57 65	8, 15, 33, 55	4 (1%)
1	C	276/293 (94%)	-0.47	6 (2%) 62 70	6, 11, 28, 44	6 (2%)
All	All	819/879 (93%)	-0.25	17 (2%) 63 70	6, 13, 31, 55	14 (1%)

All (17) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	56	ALA	4.4
1	B	313	PRO	3.9
1	B	56	ALA	3.3
1	A	43	GLY	3.2
1	C	313	PRO	2.8
1	B	312	GLU	2.8
1	B	293	MET	2.7
1	B	311	TYR	2.7
1	C	249	ASP	2.7
1	B	43	GLY	2.6
1	C	318	GLY	2.4
1	A	293	MET	2.3
1	B	308	ILE	2.3
1	C	312	GLU	2.1
1	C	311	TYR	2.1
1	C	314	HIS	2.1
1	A	314	HIS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	CU	B	404	1/1	0.93	0.07	19,19,19,19	1
2	CU	A	405	1/1	0.94	0.05	16,16,16,16	1
2	CU	A	403	1/1	0.94	0.07	22,22,22,22	1
3	GOL	B	401	6/6	0.95	0.09	13,15,18,24	0
3	GOL	C	401	6/6	0.95	0.08	15,17,23,35	0
3	GOL	B	402	6/6	0.96	0.08	18,22,25,30	0
2	CU	B	405	1/1	0.97	0.05	20,20,20,20	1
2	CU	A	404	1/1	0.98	0.07	30,30,30,30	1
2	CU	B	403	1/1	0.99	0.02	12,12,12,12	0
2	CU	A	402	1/1	0.99	0.04	20,20,20,20	1
2	CU	C	402	1/1	1.00	0.01	8,8,8,8	0
2	CU	A	401	1/1	1.00	0.02	7,7,7,7	1

6.5 Other polymers [i](#)

There are no such residues in this entry.