



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 08:01 PM UTC

PDB ID : 5UAX / pdb_00005uax
Title : Structure of apo human PYCR-1 crystallized in space group C2
Authors : Tanner, J.J.
Deposited on : 2016-12-20
Resolution : 1.85 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3428 (1.86-1.86)
Clashscore	190562	3579 (1.86-1.86)
Ramachandran outliers	187476	3553 (1.86-1.86)
Sidechain outliers	187428	3553 (1.86-1.86)
RSRZ outliers	180081	3429 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	322	6% 79% 6% 14%
1	B	322	2% 80% • 15%
1	C	322	5% 77% 8% 15%
1	D	322	4% 81% • 16%
1	E	322	19% 74% 6% 21%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 10002 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Pyrroline-5-carboxylate reductase 1, mitochondrial.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	278	Total	C	N	O	S	0	3	0
			1992	1256	349	375	12			
1	B	273	Total	C	N	O	S	0	2	0
			1946	1228	340	366	12			
1	C	274	Total	C	N	O	S	0	2	0
			1932	1220	335	365	12			
1	D	270	Total	C	N	O	S	0	1	0
			1912	1206	336	359	11			
1	E	255	Total	C	N	O	S	0	0	0
			1713	1078	304	322	9			

There are 110 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-21	MET	-	initiating methionine	UNP P32322
A	-20	HIS	-	expression tag	UNP P32322
A	-19	HIS	-	expression tag	UNP P32322
A	-18	HIS	-	expression tag	UNP P32322
A	-17	HIS	-	expression tag	UNP P32322
A	-16	HIS	-	expression tag	UNP P32322
A	-15	HIS	-	expression tag	UNP P32322
A	-14	SER	-	expression tag	UNP P32322
A	-13	SER	-	expression tag	UNP P32322
A	-12	GLY	-	expression tag	UNP P32322
A	-11	VAL	-	expression tag	UNP P32322
A	-10	ASP	-	expression tag	UNP P32322
A	-9	LEU	-	expression tag	UNP P32322
A	-8	GLY	-	expression tag	UNP P32322
A	-7	THR	-	expression tag	UNP P32322
A	-6	GLU	-	expression tag	UNP P32322
A	-5	ASN	-	expression tag	UNP P32322
A	-4	LEU	-	expression tag	UNP P32322
A	-3	TYR	-	expression tag	UNP P32322

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	PHE	-	expression tag	UNP P32322
A	-1	GLN	-	expression tag	UNP P32322
A	0	SER	-	expression tag	UNP P32322
B	-21	MET	-	initiating methionine	UNP P32322
B	-20	HIS	-	expression tag	UNP P32322
B	-19	HIS	-	expression tag	UNP P32322
B	-18	HIS	-	expression tag	UNP P32322
B	-17	HIS	-	expression tag	UNP P32322
B	-16	HIS	-	expression tag	UNP P32322
B	-15	HIS	-	expression tag	UNP P32322
B	-14	SER	-	expression tag	UNP P32322
B	-13	SER	-	expression tag	UNP P32322
B	-12	GLY	-	expression tag	UNP P32322
B	-11	VAL	-	expression tag	UNP P32322
B	-10	ASP	-	expression tag	UNP P32322
B	-9	LEU	-	expression tag	UNP P32322
B	-8	GLY	-	expression tag	UNP P32322
B	-7	THR	-	expression tag	UNP P32322
B	-6	GLU	-	expression tag	UNP P32322
B	-5	ASN	-	expression tag	UNP P32322
B	-4	LEU	-	expression tag	UNP P32322
B	-3	TYR	-	expression tag	UNP P32322
B	-2	PHE	-	expression tag	UNP P32322
B	-1	GLN	-	expression tag	UNP P32322
B	0	SER	-	expression tag	UNP P32322
C	-21	MET	-	initiating methionine	UNP P32322
C	-20	HIS	-	expression tag	UNP P32322
C	-19	HIS	-	expression tag	UNP P32322
C	-18	HIS	-	expression tag	UNP P32322
C	-17	HIS	-	expression tag	UNP P32322
C	-16	HIS	-	expression tag	UNP P32322
C	-15	HIS	-	expression tag	UNP P32322
C	-14	SER	-	expression tag	UNP P32322
C	-13	SER	-	expression tag	UNP P32322
C	-12	GLY	-	expression tag	UNP P32322
C	-11	VAL	-	expression tag	UNP P32322
C	-10	ASP	-	expression tag	UNP P32322
C	-9	LEU	-	expression tag	UNP P32322
C	-8	GLY	-	expression tag	UNP P32322
C	-7	THR	-	expression tag	UNP P32322
C	-6	GLU	-	expression tag	UNP P32322
C	-5	ASN	-	expression tag	UNP P32322

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	-4	LEU	-	expression tag	UNP P32322
C	-3	TYR	-	expression tag	UNP P32322
C	-2	PHE	-	expression tag	UNP P32322
C	-1	GLN	-	expression tag	UNP P32322
C	0	SER	-	expression tag	UNP P32322
D	-21	MET	-	initiating methionine	UNP P32322
D	-20	HIS	-	expression tag	UNP P32322
D	-19	HIS	-	expression tag	UNP P32322
D	-18	HIS	-	expression tag	UNP P32322
D	-17	HIS	-	expression tag	UNP P32322
D	-16	HIS	-	expression tag	UNP P32322
D	-15	HIS	-	expression tag	UNP P32322
D	-14	SER	-	expression tag	UNP P32322
D	-13	SER	-	expression tag	UNP P32322
D	-12	GLY	-	expression tag	UNP P32322
D	-11	VAL	-	expression tag	UNP P32322
D	-10	ASP	-	expression tag	UNP P32322
D	-9	LEU	-	expression tag	UNP P32322
D	-8	GLY	-	expression tag	UNP P32322
D	-7	THR	-	expression tag	UNP P32322
D	-6	GLU	-	expression tag	UNP P32322
D	-5	ASN	-	expression tag	UNP P32322
D	-4	LEU	-	expression tag	UNP P32322
D	-3	TYR	-	expression tag	UNP P32322
D	-2	PHE	-	expression tag	UNP P32322
D	-1	GLN	-	expression tag	UNP P32322
D	0	SER	-	expression tag	UNP P32322
E	-21	MET	-	initiating methionine	UNP P32322
E	-20	HIS	-	expression tag	UNP P32322
E	-19	HIS	-	expression tag	UNP P32322
E	-18	HIS	-	expression tag	UNP P32322
E	-17	HIS	-	expression tag	UNP P32322
E	-16	HIS	-	expression tag	UNP P32322
E	-15	HIS	-	expression tag	UNP P32322
E	-14	SER	-	expression tag	UNP P32322
E	-13	SER	-	expression tag	UNP P32322
E	-12	GLY	-	expression tag	UNP P32322
E	-11	VAL	-	expression tag	UNP P32322
E	-10	ASP	-	expression tag	UNP P32322
E	-9	LEU	-	expression tag	UNP P32322
E	-8	GLY	-	expression tag	UNP P32322
E	-7	THR	-	expression tag	UNP P32322

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
E	-6	GLU	-	expression tag	UNP P32322
E	-5	ASN	-	expression tag	UNP P32322
E	-4	LEU	-	expression tag	UNP P32322
E	-3	TYR	-	expression tag	UNP P32322
E	-2	PHE	-	expression tag	UNP P32322
E	-1	GLN	-	expression tag	UNP P32322
E	0	SER	-	expression tag	UNP P32322

- Molecule 2 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total Cl 1 1	0	0
2	B	1	Total Cl 1 1	0	0
2	C	1	Total Cl 1 1	0	0
2	D	1	Total Cl 1 1	0	0
2	E	1	Total Cl 1 1	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	109	Total O 109 109	0	0
3	B	118	Total O 118 118	0	0
3	C	93	Total O 93 93	0	0
3	D	116	Total O 116 116	0	0
3	E	66	Total O 66 66	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Pyrroline-5-carboxylate reductase 1, mitochondrial




- Molecule 1: Pyrroline-5-carboxylate reductase 1, mitochondrial



- Molecule 1: Pyrroline-5-carboxylate reductase 1, mitochondrial



- Molecule 1: Pyrroline-5-carboxylate reductase 1, mitochondrial



SER
PRO
ALA
ALA
ILE
LYS
LYS
ILE
THR
THR
ASP
LYS
VAL
LYS
LEU
ASP
SER
PRO
ALA
GLY
THR
ALA
LEU

● Molecule 1: Pyrroline-5-carboxylate reductase 1, mitochondrial



MET
HIS
HIS
HIS
HIS
HIS
SER
SER
GLY
VAL
ASP
LEU
GLY
THR
GLU
ASN
LEU
TYR
PHE
GLN
SER
M1
S2
V3
G4
F5
I6
G7
A8
G9
Q10
L11
A12
F13
A14
F19
T20
A21
A22
G23
V24
L25
A26
A27
H28
K29
I30
M31
A32
S33
S34
PRO
ASP
MET
ASP
LEU
ALA
THR

VAL
SER
ALA
LEU
ARG
LYS
MET
GLY
V50
K51
L52
T53
P54
H55
H56
K57
E58
T59
V60
Q61
H62
S63
D64
V65
L66
F67
L68
A69
V70
K71
P72
H73
I74
I75
F76
F77
I78
L79
D80
E81
I82
D85
D89
I91
S94
V99
F111
R112
P113
A114
P115
R119
C120

R129
T133
H140
A141
Q142
V143
E144
C159
G177
Y180
M216
Q222
K228
L268
Q269
S270
MET
ALA
ASP
GLN
GLU
GLN
VAL
SER
PRO
PRO
ALA
ALA
ILE
LYS
LYS
THR
ILE
LEU
ASP
LYS
VAL
LYS
LEU
ASP
SER
PRO
ALA
GLY
THR
ALA
LEU

4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	183.85Å 120.69Å 88.14Å 90.00° 109.25° 90.00°	Depositor
Resolution (Å)	60.34 – 1.85 60.34 – 1.85	Depositor EDS
% Data completeness (in resolution range)	99.3 (60.34-1.85) 99.3 (60.34-1.85)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.71 (at 1.86Å)	Xtriage
Refinement program	PHENIX (1.10.1_2155)	Depositor
R, R_{free}	0.172 , 0.200 0.175 , 0.202	Depositor DCC
R_{free} test set	7616 reflections (4.93%)	wwPDB-VP
Wilson B-factor (Å ²)	24.2	Xtriage
Anisotropy	0.522	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 56.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	10002	wwPDB-VP
Average B, all atoms (Å ²)	35.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.76% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.69	0/2033	0.75	0/2768
1	B	0.69	0/1983	0.77	0/2700
1	C	0.63	0/1968	0.73	0/2682
1	D	0.69	0/1945	0.77	0/2647
1	E	0.54	0/1737	0.67	0/2373
All	All	0.65	0/9666	0.74	0/13170

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1992	0	1973	14	0
1	B	1946	0	1935	8	0
1	C	1932	0	1888	11	0
1	D	1912	0	1899	3	0
1	E	1713	0	1612	7	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	D	1	0	0	0	0
2	E	1	0	0	0	0
3	A	109	0	0	0	0
3	B	118	0	0	0	0
3	C	93	0	0	0	0
3	D	116	0	0	0	0
3	E	66	0	0	0	0
All	All	10002	0	9307	40	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (40) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:231:VAL:HG12	1:B:124[B]:THR:HG21	1.87	0.57
1:A:204[A]:ARG:NH1	1:A:208:GLN:HB2	2.21	0.56
1:A:93:VAL:HG21	1:A:149[A]:MET:HE2	1.89	0.53
1:C:1:MET:HE2	1:C:148:LEU:HD21	1.89	0.52
1:E:33:SER:HA	1:E:53:THR:O	2.10	0.51
1:A:204[A]:ARG:HH12	1:A:208:GLN:HB2	1.75	0.51
1:A:8:ALA:HB3	1:A:34:SER:HB2	1.92	0.51
1:C:185:LEU:HD21	1:C:210:LEU:HG	1.94	0.49
1:B:33:SER:HA	1:B:53:THR:O	2.13	0.48
1:D:114:ALA:HA	1:D:140:HIS:CD2	2.49	0.48
1:D:8:ALA:HB3	1:D:34:SER:HB2	1.97	0.47
1:A:185:LEU:HD21	1:A:210:LEU:HG	1.97	0.47
1:A:230:ASN:O	1:B:10[B]:GLN:HG3	2.15	0.47
1:B:48:MET:HE2	1:B:48:MET:HB3	1.87	0.45
1:C:33:SER:HA	1:C:53:THR:O	2.18	0.44
1:D:124:THR:N	1:D:125:PRO:CD	2.81	0.44
1:E:22:ALA:HA	1:E:129:ARG:CZ	2.48	0.44
1:A:108:LEU:HD23	1:A:108:LEU:HA	1.84	0.44
1:E:114:ALA:N	1:E:115:PRO:HD3	2.32	0.43
1:C:204:ARG:HH12	1:C:208:GLN:HB2	1.83	0.43
1:E:133:THR:O	1:E:159:CYS:HA	2.19	0.43
1:A:1:MET:HE2	1:A:148:LEU:HD21	2.01	0.43
1:C:93:VAL:HG21	1:C:149[A]:MET:HE2	2.00	0.42
1:C:109:SER:HA	1:C:112:ARG:O	2.19	0.42
1:E:114:ALA:HA	1:E:140:HIS:CG	2.54	0.42
1:B:172:GLY:HA2	1:B:261:SER:OG	2.19	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:133:THR:O	1:C:159:CYS:HA	2.20	0.41
1:C:264:ARG:O	1:C:268:LEU:HG	2.19	0.41
1:E:99:VAL:O	1:E:119:ARG:HD2	2.20	0.41
1:C:13:PHE:HA	1:C:45:LEU:HD21	2.03	0.41
1:E:177:GLY:HA2	1:E:180:TYR:CD2	2.56	0.41
1:A:48:MET:HE2	1:A:48:MET:HB3	1.79	0.41
1:B:185:LEU:HD21	1:B:210:LEU:HG	2.02	0.41
1:A:230:ASN:HB3	1:B:10[A]:GLN:NE2	2.36	0.41
1:C:114:ALA:HA	1:C:140:HIS:CD2	2.56	0.41
1:A:109:SER:HA	1:A:112:ARG:O	2.19	0.41
1:A:172:GLY:HA2	1:A:261:SER:OG	2.21	0.41
1:B:133:THR:O	1:B:159:CYS:HA	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	279/322 (87%)	271 (97%)	8 (3%)	0	100	100
1	B	273/322 (85%)	269 (98%)	4 (2%)	0	100	100
1	C	272/322 (84%)	263 (97%)	9 (3%)	0	100	100
1	D	269/322 (84%)	264 (98%)	5 (2%)	0	100	100
1	E	251/322 (78%)	243 (97%)	8 (3%)	0	100	100
All	All	1344/1610 (84%)	1310 (98%)	34 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	200/254 (79%)	193 (96%)	7 (4%)	32	16
1	B	196/254 (77%)	192 (98%)	4 (2%)	48	36
1	C	190/254 (75%)	186 (98%)	4 (2%)	47	33
1	D	190/254 (75%)	188 (99%)	2 (1%)	65	57
1	E	152/254 (60%)	147 (97%)	5 (3%)	33	18
All	All	928/1270 (73%)	906 (98%)	22 (2%)	43	28

All (22) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	43	SER
1	A	112	ARG
1	A	120	CYS
1	A	125	PRO
1	A	210	LEU
1	A	222	GLN
1	A	228	LYS
1	B	37	MET
1	B	210	LEU
1	B	216	MET
1	B	222	GLN
1	C	0	SER
1	C	39	LEU
1	C	120	CYS
1	C	222	GLN
1	D	120	CYS
1	D	142	GLN
1	E	94	SER
1	E	120	CYS
1	E	216	MET
1	E	222	GLN
1	E	228	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (13) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	28	HIS
1	A	140	HIS
1	A	226	GLN
1	A	230	ASN
1	B	222	GLN
1	C	140	HIS
1	C	222	GLN
1	C	226	GLN
1	D	62	HIS
1	E	56	ASN
1	E	222	GLN
1	E	226	GLN
1	E	230	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 5 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	278/322 (86%)	0.16	20 (7%) 21 23	13, 31, 58, 76	3 (1%)
1	B	273/322 (84%)	-0.06	7 (2%) 57 61	13, 29, 55, 85	2 (0%)
1	C	274/322 (85%)	0.18	17 (6%) 26 29	14, 33, 61, 85	2 (0%)
1	D	270/322 (83%)	0.00	12 (4%) 39 42	15, 30, 56, 87	1 (0%)
1	E	255/322 (79%)	0.89	60 (23%) 2 2	17, 45, 80, 88	0
All	All	1350/1610 (83%)	0.23	116 (8%) 16 16	13, 32, 69, 88	8 (0%)

All (116) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	37	MET	5.9
1	D	269	GLN	5.5
1	B	-2	PHE	5.5
1	E	55	HIS	5.3
1	E	74	ILE	5.2
1	E	72	PRO	5.1
1	C	42	VAL	5.1
1	E	143	VAL	4.6
1	E	34	SER	4.6
1	E	30	ILE	4.5
1	E	73	HIS	4.3
1	E	82	ILE	4.2
1	E	27	ALA	4.2
1	E	50	VAL	4.1
1	D	39	LEU	4.1
1	A	37	MET	4.0
1	E	64	ASP	4.0
1	E	26	ALA	3.9
1	D	143	VAL	3.9
1	C	74	ILE	3.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	E	270	SER	3.8
1	A	42	VAL	3.8
1	C	143	VAL	3.7
1	E	28	HIS	3.7
1	E	54	PRO	3.7
1	C	38	ASP	3.7
1	E	75	ILE	3.7
1	E	52	LEU	3.6
1	C	34	SER	3.6
1	C	43	SER	3.6
1	E	11	LEU	3.6
1	D	37	MET	3.5
1	E	1	MET	3.5
1	E	69	ALA	3.5
1	B	37	MET	3.5
1	D	40	ALA	3.5
1	E	24	VAL	3.4
1	E	70	VAL	3.4
1	E	13	PHE	3.3
1	B	35	PRO	3.3
1	C	39	LEU	3.2
1	E	25	LEU	3.2
1	C	269	GLN	3.1
1	E	14	ALA	3.1
1	B	143	VAL	3.1
1	E	8	ALA	3.1
1	E	10	GLN	3.0
1	E	58	GLU	3.0
1	E	85	ASP	3.0
1	E	31	MET	3.0
1	A	270	SER	3.0
1	E	3	VAL	2.9
1	E	77	PHE	2.9
1	C	8	ALA	2.9
1	C	40	ALA	2.9
1	A	8	ALA	2.9
1	E	7	GLY	2.9
1	B	77	PHE	2.9
1	E	80	ASP	2.9
1	D	38	ASP	2.8
1	A	39	LEU	2.8
1	E	142	GLN	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	E	68	LEU	2.8
1	B	38	ASP	2.8
1	A	35	PRO	2.7
1	A	-7	THR	2.7
1	A	75	ILE	2.7
1	E	23	GLY	2.7
1	E	111	PHE	2.6
1	B	270	SER	2.6
1	A	77	PHE	2.6
1	E	99	VAL	2.6
1	A	44	ALA	2.6
1	E	62	HIS	2.5
1	E	19	PHE	2.5
1	E	66	LEU	2.5
1	D	24	VAL	2.5
1	E	113	PRO	2.5
1	E	67	PHE	2.5
1	C	11	LEU	2.5
1	A	7	GLY	2.5
1	E	78	ILE	2.5
1	E	56	ASN	2.5
1	E	2	SER	2.5
1	E	63	SER	2.5
1	E	53	THR	2.4
1	C	270	SER	2.4
1	E	88	ASP	2.4
1	C	54	PRO	2.4
1	E	60	VAL	2.4
1	E	20	THR	2.3
1	C	69	ALA	2.3
1	A	143	VAL	2.3
1	E	269	GLN	2.3
1	D	43	SER	2.3
1	D	41	THR	2.3
1	D	268	LEU	2.3
1	E	268	LEU	2.3
1	C	7	GLY	2.2
1	A	43	SER	2.2
1	E	29	LYS	2.2
1	D	36	ASP	2.2
1	A	38	ASP	2.1
1	E	112	ARG	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	40	ALA	2.1
1	E	21	ALA	2.1
1	C	52	LEU	2.1
1	A	-3	TYR	2.1
1	E	91	ILE	2.1
1	E	144	GLU	2.1
1	A	54	PRO	2.0
1	D	35	PRO	2.0
1	A	11	LEU	2.0
1	A	97	ALA	2.0
1	E	5	PHE	2.0
1	A	72	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	CL	B	401	1/1	0.97	0.10	34,34,34,34	0
2	CL	E	401	1/1	0.97	0.08	36,36,36,36	0
2	CL	D	401	1/1	0.98	0.07	29,29,29,29	0
2	CL	C	401	1/1	0.98	0.05	28,28,28,28	0
2	CL	A	401	1/1	0.99	0.04	25,25,25,25	0

6.5 Other polymers [i](#)

There are no such residues in this entry.